

Review: The effect and possible mitigation of UV radiation on baculoviruses as bioinsecticides

ERISE ANGGRAINI^{1,2,*}, WEI HONG LAU^{1,♥}, GANESAN VADAMALAI¹, LIH LING KONG³, MAZIDAH MAT⁴

¹Department of Plant Protection, Faculty of Agriculture, Universiti Putra Malaysia. 43400 UPM Serdang, Selangor, Malaysia.

Tel./fax.: +60-3-9769 1000, ♥email: lauwei@upm.edu.my

²Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya. Jl. Raya Palembang-Prabumulih Km. 32, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia. ♥email: erise.anggraini@unsri.ac.id

³Institute of Plantation Studies, Universiti Putra Malaysia. 43400 UPM Serdang, Selangor, Malaysia

⁴Malaysian Agricultural Research and Development Institute. Persiaran MARDI-UPM, 43400 Serdang, Selangor, Malaysia

Manuscript received: 13 December 2021. Revision accepted: 28 June 2022.

Abstract. Anggraini E, Lau WH, Vadamalai G, Kong LL, Mat M. 2022. Review: The effect and possible mitigation of UV radiation on baculoviruses as bioinsecticides. *Biodiversitas* 23: 3721-3735. Indiscriminate use of pesticides has raised concerns over environmental and human health issues. A more environmentally-friendly biological approach to control insect pests is gaining importance. For decades, bioinsecticides based on viruses have been used to protect crops and forests. The viruses belonging to the *Baculoviridae* family are known to infect insect pests, and hence can be used as biopesticides. Baculoviruses are mostly specific to the host and are not lethal to the non-target organism. However, their usage is limited for field application due to a slow kill speed and sensitivity towards UV irradiation. This review paper discusses the life cycle of baculoviruses, their mode of infection and transmission as well as the influence of UV radiation on the effectiveness of baculoviral insecticides for field application.

Keywords: Baculovirus, insect pest, irradiated virus, UV radiation

INTRODUCTION

Chemical pesticides are commonly used to control insect pests in the field, but the indiscriminate use of chemical pesticides has raised concern over the pollution of natural resources and the development of resistant target pests, as well as detrimental effects on human health, useful insects and non-targeted organisms (Beas-Catena et al. 2014). There is substantial documentation of the adverse effects of synthetic chemical insecticides on beneficial insects of vegetable crops (Smagghe et al. 2013; Fernandes et al. 2016). Alternatively, biological control methods are becoming popular, particularly in controlling the insect pest population (Barratt et al. 2018). The use of biocontrol agents not only benefits the economy and the life of animals and users but also reduces the exposure of harmful chemicals to the environment (Chidawanyika et al. 2012). Microbial pesticides are less costly and environmentally friendly (Usta 2013; Kumari et al. 2014; Ruiu 2018). Entomopathogenic microbes such as viruses, bacteria and fungi are the common microbial biopesticides used in the field (Kvakkestad et al. 2020).

Entomopathogenic fungi (EPF) are claimed to be effective in controlling insect pests where they can infect disease-causing insects under the right conditions (Trizelia et al. 2015; Bamisile et al. 2021). EPF directly colonise the insect's body by penetrating the cuticle (Bamisile et al. 2021). To date, more than 700 species from about 90 different genera have been designated as insect pathogenic

fungi (Khachatourians and Qazi 2008), and 20 out of 31 orders of insects are infected by EPF (Araújo and Hughes 2019). However, EPFs require 2-3 weeks to kill their insect hosts (Singh et al. 2017). In addition, entomopathogenic bacteria, such as the commercial biocontrol agent *Bacillus thuringiensis* (Bt), are known as the most successful microbial insecticides against different orders of insect pests in agriculture (Jouzani et al. 2017). During sporulation, Bt releases insecticidal crystalline inclusions called Cry or Cyt toxins, which demonstrate a highly selective spectrum of insecticidal activity and kill insects within a confined area (Bravo et al. 2011). Nowadays, the crystal toxins of Bt can be expressed in plants to create crops resistant to certain insect pests (Abbas 2018). However, insects can develop resistance to Bt transgenic crops (Tabashnik et al. 2013; Tay et al. 2015) and bacteria toxins (Tetreau 2018).

Entomopathogenic viruses are well-known as very specific to target insects (Kalha et al. 2014), and they can cause epizootics in the natural population of insects (Sun 2015). For example, the *Lymantria xyliana* multiple nucleopolyhedrovirus (LyxyMNPV) is highly virulent to the casuarina moth *Lymantria xyliana* Swinehoe and has been investigated as a possible biopesticide for controlling outbreaks of *L. xyliana* Swinehoe (Nai et al. 2010). Among viruses that are known to infect insects, the *Baculoviridae* family is considered a potential pesticide for field application because they are host-specific and typically confined to a single or a small number of closely

related insect species (Haase et al. 2015). Moreover, they do not produce any toxins or metabolites that could harm human health (Landwehr 2021) and are not pathogenic to vertebrates (Amalfi et al. 2020). In addition, baculoviruses have polyhedral occlusive bodies (OBs) that confer resistance to adverse environmental conditions (Harrison and Hoover 2012a; Susurluk et al. 2013; Sajjan and Hinchigeri 2016). Furthermore, baculoviruses have been successfully controlled using the classical biological control methods (Ibarra and Del Rincón-Castro 2008). Introduction of the viruses into an insect population that has never been treated previously, and once inoculated and established, the virus will persist in the insect population without further application (Podgwaite 1985; Ibarra and Del Rincón-Castro 2008). Moreover, based on the efficacy, Nucleopolyhedrovirus (NPV), a member of baculoviruses, has been used to control pests biologically and has caused almost 90% larval mortality (Cuartas et al. 2014). NPVs have tremendous potential as bioinsecticides compared to other viral family members.

Baculovirus usage as a potential bioinsecticide may have potential limitations. For instance, UV radiation can affect the efficacy of baculoviruses (Zhang et al. 2016). Although they do not rely on the temperature and humidity to infect their host, their stability could be challenged by solar irradiation under field conditions in the ultraviolet spectrum at 290-400 nm (Bullock et al. 1970) particularly in the UV-B range (280-320 nm) (Lasa et al. 2007). Another study has revealed that extending the duration of UV exposure resulted in a decline in both virus concentration and insect mortality (Salama et al. 2017). UV radiation is also a challenge to other biological control agents such as EPFs and Bt. The mortality of *Spodoptera litura* decreased significantly when exposed to *Beauveria bassiana* irradiated under UV-C (Herlinda et al. 2020). *Beauveria bassiana* was unable to adjust to direct sunlight, and its conidia were swiftly destroyed by ultraviolet (UV) radiation (Jackson et al. 2010; Kaiser et al. 2019). Sunlight exposure could affect the efficacy of *Bacillus thuringiensis israelensis* (Bti) and *Bacillus sphaericus* (Bs) against *Anopheles* larvae (Zogo et al. 2019). This review paper will give insights into the limitation of baculovirus as biocontrol agents, specifically the effects of UV radiation on the infection cycle of baculoviruses and the preventive measures against UV inactivation of baculoviruses.

Baculovirus

Baculoviruses are the major group of arthropod viruses that are well known as biocontrol agents against pests of agricultural crops. They are divided into four genera based on their target host (ICTV 2020). The genera *Alphabaculovirus* (baculoviruses are called nucleopolyhedrovirus or NPV) and *Betabaculovirus* (baculoviruses are called granulovirus or GV) infect lepidopteran insects while the genera *Gammabaculovirus* (NPV) and *Deltabaculovirus* infect hymenopteran and dipteran insects, respectively (Jehle et al. 2006; van Oers 2021). The name of baculovirus is derived from its rod-shaped nucleocapsids. Baculovirus contains double-stranded DNA (dsDNA), which is circular, supercoiled and

protected from the environmental hazards due to the presence of a polyhedral crystalline matrix (Rohrmann 2019). Its genome size ranges from 80-200 kb (Szewczyk et al. 2006), which encodes 70-180 genes. Of these, 31 core genes are well identified in all baculovirus genomes. These core genes are involved in various functions like infection, cell cycle arrest, replication, transduction, virus assembly, packaging and release (Miele et al. 2011).

The infection process of baculoviruses

During the life cycle of a baculovirus, two phenotypic virions, namely budded virions (BV) and ocular-derived virions (ODV), are produced at distinct chronological stages. BV is produced during the early stage of infection. It buds off from the cytoplasmic membrane as a single nucleocapsid per envelope and is involved in the secondary infection. ODV is produced during the latter phase of infection and embedded in a protein matrix, which is crystalline in nature and referred to as occlusion bodies (OB). Even though these two kinds of virions display similarities in their genome, they are different in their origins, envelope compositions and functions in the viral life cycle (Haase et al. 2015). The major proteins of the crystalline matrix in the OBs are called polyhedron (NPV) and granule (GV). This protein is very stable outside the host. Therefore, OBs can withstand different environmental conditions and prevail in the infectious state indefinitely.

The primary infection starts when OBs are ingested by susceptible lepidopteran larvae *per os*. The gastrointestinal tract of insects is the site of baculoviral infection. It can be divided into three categories: foregut, midgut and hindgut (Saxena et al. 2018). After ingestion, due to the extreme alkaline pH of the midgut, the OB is degraded to release ODVs. The ODVs are responsible for primary infection of the midgut epithelial cells. They pass through the peritrophic membrane (PM), which is made up of chitin and glycoprotein, lines the midgut and protects the cells from abrasion, cut, or insect pathogens (Terra 2001). Some baculoviruses encode an enzyme (metalloproteinase) containing divalent cations as an integral part of their structure, which helps them to cross PM. The enhancin is a metalloproteinase, which is believed to digest mucin, a component of PM, thereby allowing virus access to the epithelial cell surface (Wang and Hu 2019). The enhancin of GV is believed to increase the infectivity of some NPVs (Slavicek and Popham 2005). After crossing the PM, ODV will bind to the ODV-specific receptors and fuse to the brush border of the microvilli of the columnar epithelial cells, releasing nucleocapsids (Figure 1). Nucleocapsids travel through the cytoplasm to the nucleus which is the site for baculovirus replication (Harrison and Hoover 2012b). Transcription and translation of early genes then take place. Some proteins are translocated into the nucleus where nucleocapsids and virions are assembled. These nucleocapsids bud through the basal membrane of the cell as BV. It is presumed that during this stage, they obtain an envelope containing GP64 protein which can recognize a target cell membrane. The GP64 protein is encoded by the *GP64* gene and it plays a role in the attachment of BVs to the cell membrane (Hefferon et al. 1999). These BVs

disperse in the haemolymph to infect other cells during the systemic phase of infection (Becnel and White 2007; Rohrmann 2019). Most BVs escape the midgut basal lamina to infect larval tissues (Clem and Passarelli 2013). These secondary infections by the BVs attack all cells in the larval body such as tracheal cells, haemolymph, body fat and epidermis cells. Through the process of receptor mediated-endocytosis, they enter the cells and replicate in the nucleus (D'Amico and Slavicek 2012; Khorramnejad et al. 2021). During the late stage of infection, nucleocapsids are enveloped to form ODVs, which will be occluded in the polyhedral crystalline matrix as OBs. These OBs accumulate in the nucleus (Sajap et al. 2000). As a result of the accumulation of viral proteins and virions in the nucleus and the nucleus becomes swollen. Later, the host cells lyse, the tissues liquefy and the cuticle weakens and ruptures (Monteiro et al. 2012). This liquification of the insect host will contaminate the foliage, and consequently continue the viral transmission cycle to other insect hosts (Harrison and Hoover 2012; Elder 2013), which could induce a self-propagating and positive-feedback loop (Elder 2013). In some cases where infected insects only showed symptoms in their midgut and fat body while other tissues remain uninfected (Senthil-Nathan 2015). Some insects showed sub-lethal infection in which they do not die following the ingestion of viral OBs (Williams et al. 2017).

During the process of infection, the integument of larvae changes color, and the development, feeding and mobility of larvae decrease (Harrison and Hoover 2012b).

Spodoptera litura larvae infected with *Spodoptera litura* nucleopolyhedrovirus showed characteristics of polyhedrosis symptoms (Nazli-Huda et al. 2012). The integument becomes flaccid and pink, and the infected larvae stop feeding after three days of post-inoculation (Javier-Hila and Caoili 2020). Dorsal, lateral, and the ventral sides of larvae changed to a pinkish color compared to those of healthy larvae after five days post-inoculation. The larvae ruptured after seven days post-inoculation with single dose of 2.4×10^7 OBs, releasing white body fluids filled with OBs (Nazli-Huda et al. 2012). Shortly before death, the infected larvae migrate to the upper part of the plant. They died by hanging themselves on pseudopods, a behaviour of the tree top disease induced by baculovirus (Goulson 1997). *Spodoptera exigua* larvae infected with baculoviruses died in the top 10% of the leaves of chili plants and closer to the stems of healthy conspecific species of the same stage (Rebolledo et al. 2015). This was clear evidence that baculovirus had induced climbing behaviour in the susceptible hosts. Recent research has revealed that the *ecdysteroid UDP-glucosyl transferase (egt)* gene of *Lymantria dispar* multiple nucleopolyhedrovirus had caused this behaviour in the larvae of the gypsy moth *L. dispar* (Hoover et al. 2011). On the other hand, *Trichoplusia ni* and *S. exigua* infected with a mutant AcMNPV (with a deletion of the *egt* gene) still exhibited tree-top disease after infection (Ros et al. 2015). The tree-top disease may not be solely induced by *egt* gene which may not be applicable for all baculovirus-host interactions.




Figure 1. Replication and transmission of nucleopolyhedroviruses. (1) OBs are ingested by insects during the primary infection. (2) OBs are dissolved in the midgut of the insect, and (3) ODVs are released, passing through the peritrophic membrane and (4) fused with the microvilli of midgut epithelial cells to release the nucleocapsids. (5) The nucleocapsids enter into the nucleus and release the viral DNA, which allows replication to begin. Nucleocapsid progenies form in the nucleus, bud through the nuclear membrane and form budded viruses (BV). (6) During the secondary infection, these BVs disperse in the hemolymph and cause infection to other tissues. (7) Endocytosis allows BVs enter into the cells and replicate in the nucleus. (8) Newly formed nucleocapsids can either form BVs to infect other cells or (9) be enveloped to form ODVs that are occluded within OBs. (10) During late stage of infection, fully formed OBs are seen in the nucleus. (11) Upon death, the larvae hang on the uppermost leaves of the host plant and the integument ruptures, releasing the OBs into the environment. If other insects consume the contaminated leaves, the infection cycle will begin again




Figure 2. Horizontal and vertical transmission of baculoviruses (Adapted with permission from Williams et al. 2017). (1) After larvae ingest the OBs on the contaminated leaves, (2) some infected individuals develop a fatal condition and (3) release OBs on the host plant where they can be transmitted to other susceptible insects. (4) Rainfall will wash the OBs from the leaves into the soil, where they might be transferred back to plants via biotic and abiotic mechanisms. (5) The infected insects may be symptomless and continue to complete their lifecycle. (6) These adults might disperse, (7) mate and (8) lay eggs on other host plants. (9) The viruses are transmitted from the infected parents to their progenies. (10) Vertical transmission can be sustained for multiple generations until an elicitor or stressor transforms a latent infection into (11) a deadly disease, resuming the horizontal transmission cycle.

Transmission modes of baculoviruses

Insect baculoviruses have two types of transmission mode: horizontal and vertical (Figure 2). In horizontal transmission, viruses are transmitted from infected hosts to susceptible hosts, which are not in a parent-progeny relationship. During this stage of transmission, viruses sit and wait until they encounter a susceptible host and infect them. The viruses that retain virulence outside the host can infect a host of similar or next generation. In vertical transmission, the viruses are transmitted from the infected parents to their progenies. Vertical transmission is a mechanistic transmission of viruses across several generations (Williams et al. 2017). Infection of larvae with a viral infection does not always result in death. Some insects survive, reproduce and generate progenies. It has been suggested that the development of the disease can be overcome by metamorphosis (Burand and Park 1992; Kukan 1999).

UV radiation and its potential effect on baculovirus

Although the use of baculoviruses presents several advantages, they have their limitations in achieving their optimum goal of controlling pests in the field. Besides their limited host range and slow lethality, baculoviruses can be inactivated by UV radiation. The ultra-violet spectrum of solar radiation is the primary environmental parameter that

affects the field persistence of these viruses (Beas-Catena et al. 2014).

Solar radiation consists of ultraviolet or UV (200-390 nm), visible (390-780 nm) and infrared (more than 780 nm) radiation (El-Sharkawey et al. 2009). Around 56% of all electromagnetic radiation is infrared photons, 39% is visible light, and 5% is ultraviolet (UVA+UVB) radiation (290-400 nm) that reaches the Earth's surface (Lyons et al. 2020). The sun emits ultraviolet radiation in the form of UVA (320-390 nm), UVB (280-320 nm) and UVC (200-280 nm) bands (Hockberger 2002). UVC may be inhibited by the ozone layer however UVA and UVB reach the earth's surface after atmospheric filtration (Lubin and Jensen 1995). UVB rays inactivate insect pathogens and viruses under normal circumstances (Ignoffo et al. 1977; Asano 2005; Lasa et al. 2007). As obligate pathogens, baculoviruses have to remain viable outside their host to cause an effective infection inside their host (Redman et al. 2016). When directly exposing baculoviruses to sunlight in the field, the baculoviruses could lose their infectivity within 24 hours (Petrik et al. 2003). In general, direct exposure to solar radiation may impair the viability and infectivity of microorganisms owing to DNA denaturation and the synthesis of extremely reactive oxygen radicals (Tamez-Guerra et al. 2005). Solar radiation, particularly ultraviolet (UV) rays (300-400 nm), induces mutagenic and fatal alterations in nucleic acids (Bull et al. 1976). In

prokaryotic and eukaryotic microorganisms, ultraviolet radiation has become the primary genotoxic agent (Douki et al. 2017).

Single-stranded DNA viruses are more prone to UV radiation compared to dsDNA viruses (Gerba et al. 2002). Owing to the large size of nucleic acid and the structure of the capsid, dsDNA viruses are more resistant to UV inactivation. In addition, RNA viruses are more vulnerable to UV exposure because they have no genome repair systems (Wigginton et al. 2012; Elveborg et al. 2022). In contrast to DNA-based life forms, the majority of RNA viruses lack systems for detecting and repairing mismatches, therefore, the polymerase error was not corrected (Steinhauer et al. 1992; Elveborg et al. 2022). If a virus is inactivated by UV radiation, it will lose its biological activity. In many cases, pathogens will lose their efficacy in a few days after exposure to UVB (Martin and Domek 2002). Hence, the application of baculoviruses on foliage is hindered by UVB (Killick 1990).

UV radiation disrupts normal processes and leads to the formation of dimers. UV-induced dipyrimidine photoproducts such as cyclobutene pyrimidine dimers (CPDs), pyrimidine-pyrimidone 6-4 photoproducts (64PPs) (Mouret et al. 2006) and Dewar Valence isomers will be induced after exposure to UV radiation (Douki et al. 2017). CPDs occur from the [2+2] cycloaddition reaction between the C5-C6 double bonds of two adjacent pyrimidine bases, thymine-thymine (TT) or cytosine-cytosine (CC); however, the TT photoproduct that is a well-known thymine dimer is the hallmark of UV-induced DNA damage (Douki et al. 2017). Meanwhile, the 64PPs are generated from a single covalent bond formed between the 5' end of C6 and 3' end of C4 of adjacent pyrimidines (Rastogi et al. 2010). The dewar pyrimidinone is formed by a reversible isomerization of the 64PP upon further exposure

to UVA wavelengths around 325 nm (Matsunaga et al. 1991). Predominantly, DNA damage occurs in the bi-pyrimidine sequences upon excitation of thymine, the most photoreactive base, or, to a lesser extent, of cytosine (Cadet et al. 2014). CPDs, 64PPs and Dewar Valence isomers block the replicative DNA polymerases from passing through them when they reside on a template strand, and the transcription will be terminated (Ikehata and Ono 2011). Although DNA has a repair mechanism called nucleotide excision repair (NER), double-stranded DNA can be degraded due to failure in the repair (Ikehata and Ono 2011). This can result in death of the microorganisms.

DNA sensitivity to UV radiation depends on the purine and pyrimidine composition (Smith 1966). Due to the presence of a double-ring structure in purine, their resistance towards photoreaction increases ten folds in comparison with pyrimidine (Smith and Hanawalt 2013). Hence, more pyrimidine dimers are formed. The possibility of photodimerization is relative to the occurrence of pyrimidine sequences in the neighbourhood (Kleinerhanns et al. 2013). Moreover, suitable orientation and positioning of the pyrimidine bases may result in dimerization. The precise order of adjoining base pairs and frequent occurrence of thymine determines the sensitivity of the virus to UV radiation (Figure 3). The precise mechanism of the inactivation of the virus due to UV radiation is still not completely understood. However, the hydrogen peroxides from one or more amino acids produced by UV radiation can reduce the vitality of baculovirus (Arora and Shera 2014). In viruses containing the dsRNA genome, UV radiation may result in the accumulation of uracil dimers (Williams 2017). UV radiation damages the viral genomes and capsid proteins (Eischeid and Linden 2011; Sigstam et al. 2013).




Figure 3. UV radiation damages dsDNA of baculovirus by forming thymine dimer which caused failure in DNA replication

Radiation at higher wavelengths and higher energy, such as UVA rays and visible light, can cause the inactivation of the virus if exposed for a longer period (Martin and Domek 2002). This exposure period is high in tropical regions and at higher altitudes, whereas it decreases with an increase in latitude. In temperate regions, there are variations in UV light due to cloud cover and precipitation (Williams 2017). The amount of shade provided by upper layers of plant parts or adjacent plants could reduce the exposure of OBs to solar UV rays (Maramorosch 2012). OBs present at the center part or bottom part of the plant will obtain a lesser amount of ultraviolet radiation and are likely to maintain integrity for longer than those present at the upper part. For example, *Heliothis* nuclear polyhedrosis virus was found to be rapidly inactivated on the upper leaf surface of cotton, soybean, and tomato when grown in field condition (Young and Yearian 1974). In addition, bioassay infection of *Panolis flammea* at 3 positions on the lodgepole pine tree: the upper, middle, and lower trees that were sprayed using purified *P. flammea* NPV; the results showed that the infected larvae were higher in the lower position of the tree with treatment (Killick and Warden 1991).

Methods to prevent UV inactivation of baculovirus

If baculoviruses are inactivated by UV radiation, their usage in the field will be hampered. Therefore, methods to enhance/prolong the persistence of the virus are required for effective utilization of this biocontrol agent against target pests.

Use of UV protectants

Several researchers have conducted studies to improve the effectiveness of entomopathogenic viruses and prevent them from UV inactivation. UV protectants used to protect the virus include dyes, optical brighteners, lignin derivatives or natural ingredients (Table 1). Oxidative enzymes such as dilodin, riboflavin, folic acids, inols, pyridoxine and vitamins were also claimed to be potential agents to protect entomopathogenic viruses from UV radiations (El-Helaly 2013). Several natural plant products have antioxidative potential to provide capacity to counteract UV-induced adverse effects on human skin (Dunaway et al. 2018). The composition of the customary sunscreens contain inorganic compounds, such as titanium dioxide (TiO₂), zinc oxide (ZnO) and minerals, which help to reflect and scatter UVA and UVB. Meanwhile, the mode of action of ZnO and TiO₂ is to absorb UV radiation (Cole et al. 2016). TiO₂ is more effective for UVB protection while ZnO is effective for UVA protection. The combination of these particles could enhance UV protection (Smijs and Pavel 2011). ZnO and TiO₂ are good UV protectants for *Cydia pomonella* GV isolate ZY (CpGV-ZY) (Wu et al. 2015). A recent study claimed that a novel formulation technology by encapsulating baculovirus in an ENTOSTAT-TiO₂ waxy coat has effectively safeguarded the efficacy of the sensitive viral DNA from damage by the UV radiation in the sunlight, and at the same time not phytotoxic (Wilson et al. 2020). However, the use of such metal-based particles for protecting human

skin has raised concern regarding their safety and potential toxicity.

Tinopal has been used as a UV protectant in the formulation of baculovirus-based biopesticide. It is an optical brightener derived from the stilbene compounds (Goulson et al. 2000). These compounds can absorb UV rays and reflect the light in the visible range (Martínez et al. 2009). The unformulated and formulated *Spodoptera frugiperda* NPV with Tinopal was examined. The research findings revealed that SfMNPV alone was more sensitive to UV than in mixture with 1.25 or 0.1% Tinopal C1101 (Mondragon et al. 2007). Tinopal could retain the efficacy of *S. frugiperda* NPV even after 4 hours of UV exposure (Mondragon et al. 2007). In the field studies conducted in Mexico and the United Kingdom on three distinct flower species, 0.1% or 1% concentrations of Tinopal CBS had significantly inhibited bees to flowers. A total of 64.6% rejection rate recorded on the flowers treated with Tinopal CBS compared to 5.1% on the control flowers (Goulson et al. 2000). Therefore, Tinopal is able to protect the baculovirus from UV radiation as well as to inhibit insects to the plants.

Some researchers have used natural compounds derived from leaves and fruits (referred to as herbal cosmetics) to protect entomopathogenic viruses from UV radiation (Table 1). Flavonoids are the most studied sunlight protectant which includes quercetin (3,40-dihydroxyflavonol) (Cefali et al. 2016). This compound is present in fruits (strawberries, grapefruits and apples), vegetables and beverages (tea and red wine) (Lakhanpal and Rai 2007). Quercetin has functioned as an anti-inflammatory, antioxidant, immunomodulator (Korać and Khambholja 2011), inhibits skin tumour formation (Nishino et al. 1984) and has a potential topical sunscreen (Saewan and Jimtaisong 2013). Due to the presence of aromatic rings in their molecular structure, flavonoids absorb UV rays between 200 and 400 nm, making them appropriate for use as a sunscreen agent (Cefali et al. 2016).

The use of inexpensive natural additives such as moringa, rice bran, cacao and green tea to sustain the effectiveness of *Spodoptera littoralis* multiple nucleopolyhedrovirus (*SpliMNPV*) against UV lights was analysed (El-Helaly 2013). Moringa exhibited the highest protection rate compared to rice bran, cacao and green tea. The moringa additive at 1% concentration could retain the activity of *SpliMNPV* (mortality of 93.24%) even after exposure to UV radiation. This could be due to the natural occurrence of high antioxidant content and anti-inflammatory compounds in moringa (Yong-Bing et al. 2019). Tea (polyphenol) has photo protective effect on human skin, reducing the number of sunburned cells and alleviating DNA damage (Li et al. 2014). Among the tea tested against the beet armyworm NPV (HaNPV), black tea was the best UV protectant with the highest larval mortality recorded at 90%. Black tea and lignin showed good result in protecting *Spodoptera exigua* MNPV (Salamouny et al. 2009). The combination of both black tea and lignin as UV protectant in biopesticides is safe and can be obtained easily. Plant phenolics such as caffeic acid, gallic acid and

chlorogenic acid were able to protect SeMNPV at concentration as low as 0.5 mM. The addition of a UV protectant in the baculovirus formulation has been proven effective in improving viral stability either in the laboratory or in field application. UV protectants derived from natural ingredients will be a great candidate in the biopesticide industry. However, in cases where chemicals are used as UV protectants in the viral formulation, its concentration must be kept as minimal as possible. It is important to study the harmful effect, if any, of chemicals used as UV protectant.

Selection of insect baculovirus resistant to UV irradiation

Insect baculoviruses with resistant capability to UV irradiation and maximal vertical transmission rate were selected. One strain of GV from a codling moth (CpGV) exhibited resistance to UV radiation under natural sunlight and in laboratory conditions (Brassel and Benz 1979). The resistance capacity of the altered strain was increased 5.6-fold compared to the original isolate, and its infectivity period was double that of the original isolate, i.e. it could stay for double the period in comparison with the original isolate (Brassel and Benz 1979). Witt and Stairs (1975) demonstrated that within a population of *Galleria mellonella* NPV (GmNPV) active against *Galleria mellonella*, one subpopulation was susceptible to low UV doses, while another subpopulation was susceptible only to high UV doses, equating to a nearly 1,000-fold difference in UV susceptibility. They hypothesised that this variety in UV reaction could be due to genetic variability and that it would be able to select UV-tolerant virus strains. Akhanaev et al. (2017) measured the relative rate of inactivation and the half-life of the virus to assess the UV tolerance of two LdMNPV strains from Western Siberia (LdMNPV-27/0) and North America (LdMNPV-45/0). The North American strain, previously demonstrated to be more virulent against *Lymantria dispar* larvae, was more sensitive to UV after 15 minutes of exposure to sunlight and lost its potency more quickly than the Siberian strain. Therefore, it is not recommended to utilise LdMNPV-45/0 in locations with high UV loads or at densities of forest stands. Another study reported the use of *Cryptophlebia leucotreta* granulovirus-SA (CrleGV-SA) as a commercial biopesticide against the false codling moth, *Thaumatotibia leucotreta*, in citrus and other crops (Mwanza et al. 2022). This virus strain was isolated from infected *T. leucotreta* larvae and subjected to UV irradiation in a climate chamber with simulating field settings under laboratory condition. This UV-resistant CrleGV-SA strain was thousand-fold more virulence than the wild-type isolate (Mwanza et al. 2022). Single Nucleotide Polymorphisms (SNPs) was

discovered in the 5th generation of UV-resistant CrleGV-SA which is believed to have improved the UV tolerance and pathogenicity of the virus.

Enhancing photostability of baculoviruses by genetic engineering

Isolation, insertion and expression of genes responsible for the DNA repair system in the genome of baculovirus will result in the production of UV-resistant baculovirus (Zhao et al. 2003; Ferrelli et al. 2018). Genetic modification research for improving the baculovirus resistance to UV inactivation is very much lacking. Petrik et al. (2003) had constructed a recombinant *Autographa californica* M nucleopolyhedrovirus (AcMNPV) that encoded cv-PDG, an algal virus-specific pyrimidine dimer glycosylase. The expression of cv-PDG resulted in a threefold reduction in the UV inactivation of budded virus compared to the wild type AcMNPV in plaque test using *Spodoptera frugiperda* Sf21 cells. However, the occluded viruses expressing cv-PDG did not become more resistant to UV inactivation than the wild-type AcMNPV when fed to *S. frugiperda* or *Trichoplusia ni* neonates. Several explanations for the lack of protection against UV inactivation in occluded virus were explained by Petrik et al. (2003). They assumed that the cv-PDG expression occurred in the midgut epithelial cells while the DNA repair pathways required to complete the repair process by cv-PDG were not operational in the midgut epithelial cells, therefore there was insufficient time for the viral DNA to be repaired. The insects periodically shed their midgut lining during moulting process into next instars; therefore the recombinant viruses were unable to initiate the replication in the midgut cells.

Baculoviruses as bioinsecticides in Asian countries

The effectiveness of baculoviruses as a bioinsecticide has been tested in several countries in Asia (Table 2). Research on the efficacy of wild baculoviruses is fundamental to finding out how effective the wild baculoviruses are and how they react to other active ingredients of bioinsecticides, such as bacteria or botanical pesticides, and synthetic pesticides when used in a combined application (Samada and Tambunan 2020; Landwehr 2021). The difference in the effectiveness of imported isolates and local isolates has been studied to ensure the efficacy and stability of both isolates for use in the local area (Akhanaev et al. 2017). It has been widely shown that the indigenous natural resources have provided a good control of insect pests (Grzywacz et al. 2014; Rathore et al. 2021).

Table 1. Research related to UV protectants used to prevent inactivation of baculoviruses during the past 10 years

Insect target	Virus	UV protectants (plant-based ingredients)	UV experiment	Result	Reference
<i>Spodoptera littoralis</i>	<i>SpliMNPV</i>	Orange oil 5% Tea tree oil 5% Marjoram 5%	UV lamp (Utalix; 23 sun UV, Germany)	Orange oil gave the best protection to the virus.	Sayed et al. (2020)
<i>Helicoverpa armigera</i>	HaNPV	Black tea Green tea Pepper Green cabbage Red cabbage Mango leaves Guava leaves Basil Dill Peppermint	Bacterial lamp 312 nm (Phillips TUV 30 W)	Black tea, dill, green cabbage, pepper provided the highest level of protection.	Ibrahim et al. (2019)
<i>Spodoptera littoralis</i>	<i>SpliNPV</i>	Moringa Cacao Green tea	Field experiment	Moringa could prolong the activity of the virus under field conditions.	El-Helaly (2020)
<i>Spodoptera exigua</i> First and second instar	<i>SpliMNPV</i>	Henna Green tea leave Clove flower	Field experiment	The addition of clove extract as a natural UV protectant had provided higher persistence of virus in the field.	Sukirno et al. (2018)
<i>Diaphania pulverulentalis</i> (Hampson)	DpNPV	DpNPV formulation (10% starch + 0.2% Tinopal + 1% Tween 80 + DpNPV)	UV lamp & sunlight	Tinopal showed significant protection to DpNPV when exposed to UV lamps and sunlight.	Prabhu and Mahalingam (2017)
<i>Spodoptera littoralis</i>	<i>SpliMNPV</i>	Henna whole Dates fruit (var. 'Sukari') without seed Black grape whole fruit Kiwi whole fruit, Olive fruit without seed Lemon whole fruit Pomegranate whole fruit without skin Red beetroot whole tuber Green tea whole leaves Clove whole flower	Under laboratory condition (UVB lamps 302 nm (FotodyneInc, New Berlin, WI, Germany) Field trials	Cloves, henna, green tea, pomegranate, and grape extracts exhibited a higher rate of virus protection compared to unprotected virus. In field conditions, clove and henna-based formulations had increased the persistence of the virus.	Sutanto et al. (2017)
<i>Spodoptera littoralis</i> neonate larvae	<i>SpliMNPV</i>	Cacao (5% and 10%) Green coffee (5% and 10%) Fresh green cabbage (5% and 10%) Fresh red cabbage (5% and 10%)	Natural sunlight in field	Cacao was the best protective additive.	El-Helaly et al. (2013)

<i>Spodoptera littoralis</i>	SpliNPV	Moringa Rice bran Cacao Green coffee	UV lamps with an absorbance under 300 nm	Moringa was the best candidate.	El-Helaly (2013)
<i>Spodoptera exigua</i>	SeNPV	Dust formulation: Husk charcoal Coconut shell charcoal Soot Powder Yam flour Liquid formula: Molasses Green tea filtrate Turmeric filtrate Yam filtrate	Sunlight (3000-4000 $\mu\text{W}/\text{cm}^2$)	The best dust formulations: coconut shell charcoal soot husk charcoal The best liquid formulations: molasses, turmeric filtrate yam filtrate	Samsudin et al. (2011)
<i>Chrysodeixis chalcites</i> (Esper)	ChchNPV-TF1	The photoprotective substances were divided into three groups: low protection (0-8%; kaolinite), intermediate protection (48-62%; green tea, moringa, bentonite and cacao) and high protection (87-100%; charcoal, iron oxide)	under laboratory conditions using a Crosslinker as UV light source at 200 J cm^{-2} field conditions	1% cacao and 1% charcoal were the best result for laboratory conditions. 1% charcoal treatment exhibited the highest protection.	Cakmak et al. (2021)
<i>Spodoptera littoralis</i> (neonate larvae on squash plants)	SpliMNPV	Nano aluminium oxide Nano zinc oxide Cacao (positive control)	Simulated UV light A set of four UV lamps	Nano zinc oxide was the best formulation.	El-Helaly and Sayed (2015)
<i>Helicoverpa armigera</i> third instar	HaNPV	Microencapsulations using Gelatin 3% and 5% Sodium alginate 3% and 5% Starch 3% and 5%	385 nm UVA 254 UVC With Entela lamp model UVGL 25 of 4 W	Gelatin was the best and the most stable formulation.	Gifani et al. (2015)
<i>Cydia pomonella</i> L	CpGV	ZnO (Hengxing [®] zinc oxide) 20 mg mL^{-1} TiO (Dengfeng [®] titanium dioxide) 20 mg mL^{-1}	UVB light (20W) UV lamps model K5 B20H	ZnO and TiO ₂ were good UV protectants to CpGV-ZY isolate.	Wu et al. (2015)
<i>Pieris brassicae</i>	PbGV	Iron oxide (black) 1% and 2%) Robin blue (1% and 2%) Tinopal (1% and 2%) Congo (1% and 2%)	1. UV lamps (15 W Philips G15T8) 2. Field study	Congo red was more effective. In field conditions, Congo red and robin blue gave the best protection to virus.	Sood et al. (2013)
<i>Spodoptera litura</i>	SINPV	a) Sodium 2,5-bis(4-(2,4-disulfonate) phenyl)-1,3,4-oxadiazole with solvent: (1) DMF, (2) Acetone, (3) THF b) Sodium 2,5-bis(4-(2-sulfonate) phenyl)-1,3,4-oxadiazole with solvent: (1) DMF, (2) Acetone, (3) THF	a) 1. DMF 364 nm 2. Acetone 348 nm 3. THF 349 nm b) 1. DMF 348 nm 2. Acetone 344 nm 3. THF 346 nm	Sodium 2,5-bis(4-(2-sulfonate) phenyl)-1,3,4-oxadiazole compound showed greater protection to SINPV.	Zhu et al. (2013)
<i>Spodoptera frugiperda</i>	SfMNPV	Lignin encapsulated (sodium lignin, calcium chloride dehydrates)	1. Laboratory assay using a Suntest CPS (Hereus, Germany) 2. Field application	The spray-dried lignin encapsulated formulation protected virus.	Behle and Popham (2012)
<i>Spodoptera littoralis</i>	SINPV	Starch, clay, glycerine, extract of black tea	Natural light in field application	Clay and black tea were the best UV protectants.	El-Husseini et al. (2012)

Table 2. Efficacy test of baculoviruses as bioinsecticides in Asian countries during the past 10 years

Country	Host	Virus	Efficacy	References
Russia	<i>Lymantria dispar</i>	LdMNPV-27/0 isolated from Western Siberia, Russia LdMNPV-45/0 isolated from North America	The strain isolated in North America (LdMNPV-45/0) diminished its potency significantly and more rapidly than the strain isolated in Western Siberia (LdMNPV-27/0) when exposed to sunlight.	Akhanaev et al. (2017)
Turkey	<i>Malacosoma americanum</i> (F) <i>Malacosoma neustria</i>	<i>Malacosoma neustria</i> NPV Turkish isolate 3	The virulence of ManeNPV-T3 was determined under laboratory conditions using third instar <i>M. americanum</i> and <i>M. neustria</i> larvae, with 0.6×10^5 occlusion bodies per larva inducing 100% mortality after 10 days.	Demir et al. (2014)
Japan	<i>Adoxophyes honmai</i> (Lepidoptera: Tortricidae)	<i>A. orana</i> (AdorNPV) <i>A. honmai</i> (AdhoNPV)	AdhoNPV had a lower field effectiveness in reducing damage to tea plants caused by <i>A. honmai</i> larvae than the fast-killing AdorNPV. AdhoNPV transmitted horizontally at a substantially greater rate than AdorNPV.	Takahashi et al. (2015)
India	<i>Spilosoma obliqua</i>	<i>Spilosoma obliqua</i> MNPV	SpobMNPV recorded LC_{50} of 4.46×10^5 POBs mL^{-1} and ST_{50} at 72 h, respectively, against 3rd instar <i>S. obliqua</i> larvae.	Sujayanand et al. (2019)
Mongolia	<i>Lymantria dispar asiatica</i> Vnukovskij	LdMNPV-H from Heilongjiang Province, China LdMNPV-J from Japan LdMNPV-D from commercial product	LdMNPV-D, a commercial strain accessible from North America, was moderately pathogenic, while LdMNPV-J, a Japanese isolate, was the least harmful. The dose-response slope for the three virus isolates confirmed that the Chinese gypsy moth larvae were more homogeneous susceptible to LdMNPV-H and LdMNPV-D than to LdMNPV-J.	Duan et al. (2012)
Pakistan	<i>Spodoptera litura</i>	SpltNPV-Pak-BNG isolate Pakistan	The dose-to-death response and mortality rate of various instar larvae of <i>S. litura</i> were determined. Second and third instar larvae exhibited similar dose-response to mortality, whereas 4th instar larvae were less susceptible. The mean survival time followed a similar trend to the mean death time values and was identical for 2nd and 3rd instar larvae but was significantly prolonged for 4th instar larvae.	Ali et al. (2018)
Pakistan	<i>Spodoptera litura</i> F	<i>S. litura</i> NPV (V-SpltNPV) was isolated from infected larvae on a cotton crop	Spinosad combined with V-SpltNPV (1×10^8 OBs mL^{-1}) caused 100% mortality of 2nd instar larvae in a greenhouse experiment. A single application of V-SpltNPV (1×10^8 OBs mL^{-1}) had resulted in 52.63% of examined larvae.	Ayyub et al. (2019)
Pakistan	<i>Spodoptera litura</i> F	Nuclear polyhedrosis virus (NPV) and flubendiamide	For pupation and adult emergence of <i>S. litura</i> , combining NPV and flubendiamide showed to be more lethal at higher concentrations than applying them separately at lower concentrations.	Maqsood et al. (2017)
Pakistan	<i>Plutella xylostella</i>	<i>Plutella xylostella</i> granulovirus (PxGV) and <i>Azadirachta indica</i> (AZA)	Sublethal dose rate of PxGV and AZA mixture under laboratory condition caused higher larval mortality of <i>P. xylostella</i> .	Malik et al. (2020)
Pakistan	<i>Helicoverpa armigera</i> (Hubner)	Combination NPV and 0.1% spinosad	<i>H. armigera</i> was given various concentrations of NPV and spinosad individually, as well as a combination of NPV and 0.1% Spinosad. These results showed that effectiveness and tenacity of mixture caused significant mortality of larvae.	Nawaz et al. (2019)
Indonesia	<i>Spodoptera litura</i>	<i>Spodoptera litura</i> NPV (SINPV)	The investigation demonstrated that the interaction between the SINPV isolate and the virus concentration had a substantial effect on the feeding, mortality, LC_{50} and LT_{50} . Additionally, these potential characteristics varied across the <i>S. litura</i> larval instars. SINPV JTM 97c and SINPV JTM 02-5 was potential in controlling <i>S. litura</i> on soybean.	Bedjo (2017)
Indonesia	<i>Spodoptera exigua</i> Hubn	<i>Spodoptera exigua</i> NPV (SeNPV)	Within five days of infection, SeNPV killed 77.5% larvae in a laboratory trial. Additionally, SeNPV decreased the feeding capability of <i>S. exigua</i> by 0.43 g per day. Additionally, SeNPV reduced pupae weight by 0.063 g. In a greenhouse experiment, the application of SeNPV resulted in 100% larval mortality and a 10.43% reduction in the intensity of shallot damage caused by <i>S. exigua</i> .	Supyani et al. (2014)
Philippines	<i>Spodoptera litura</i> F	<i>Spodoptera picta</i> NPV (SppiNPV)	The SppiNPV produced signs and symptoms of baculoviral infection in <i>S. litura</i> larvae, indicating SppiNPV could cross-infect the <i>S. litura</i> larvae.	Javier-Hila and Caoili (2020)

Synchronisation of viral bioinsecticide application and bioecology of target insect

Knowledge of the insect pest behaviour and biology involved are prerequisites for successful bioinsecticide application. Insect behaviour encompasses a vast range of behaviours, such as movement, grooming, feeding, communication, reproduction, dispersal, flight, learning, migration, host or prey selection, diapause, and numerous reactions to environmental threats like temperature, humidity, parasites, and toxins (Hoy 2019). Understanding the behaviour of insect pest could improve pest management programs (Campbell 2006). Insect life cycles vary depending on the species. The complete and incomplete metamorphosis are the two most typical types. The life cycle is essential for controlling insects because, depending on the type of life cycle, an insect's habitat, behaviours, and appearance can change significantly over the course of its lifetime (Wilbur 1980).

For example, the newly hatched larvae of the tobacco armyworm, *Spodoptera litura* scrape the chlorophyll content and feed voraciously (Krisnawati et al. 2021). This will cause the leaf to appear yellowish-white. As the caterpillar grows, it consumes the leaves entirely, leaving behind only the stalks and midribs. Additionally, larval feeding becomes more aggressive as the larvae mature. The larvae eat in fragile pods initially, then burrow through the thick pod and consume the grains (Sunil et al. 2019). During the daytime, the larvae hide under the plants, cracks and crevices of soil debris, and during the night time, they feed on the plants (Natarikar and Balikai 2017). The intensity of the sun is low in the evening, and nocturnal insects become active at night. Since *S. litura* activity was observed in the dark, the application of the virus can be made late in the afternoon, thereby protecting the viral bioinsecticide from UV rays. According to Dezianian et al. (2010), younger larvae of *Plutella xylostella* L. were more susceptible to PxGV; however, the older larvae were less tolerant. Thus, the application of PxGV should be targeted at the early stages of larval development to obtain full control of the insect hosts.

Attention must be paid to the local climate as well when applying baculoviruses to the field. Indonesia has a tropical climate with two seasons throughout the year. In 2017, the average solar exposure was 4.1 to 8 hours per day. Less than 10% of days in the year 2017 achieved 12 hours of solar exposure per day (Hamdi 2019). Incoming solar radiation depends on many factors like altitude and latitude, aerosols, season, and cloud cover (Hillel et al. 2008). It is known that the average sunlight decreases sharply according to the height of the place. Moreover, solar UV radiation increased with altitude, which was mainly due to a decrease in the number of air molecules, ozone, aerosols and clouds in the atmosphere, in addition to surfaces being covered with snow (Gröbner 2013). The altitude effect plays an important role in the understanding of the UV radiation field in mountainous terrain (Schmucki and Philipona 2002).

Baculoviruses have polyhedral occlusive bodies (OBs) that shield the viruses from harsh environmental conditions

(Susurluk et al. 2013). Formulation of baculoviruses should focus on whether the carrier could affect the efficacy of OBs. Detergents such as sodium dodecyl sulphate (SDS), sodium deoxycholate and Triton X were often utilised in laboratory studies to extract OBs from the dead baculovirus-infected larvae or cells (Kato et al. 2011), and SDS could affect the efficacy of the baculoviruses (Nazli-Huda et al. 2012). As a result, the composition of bioinsecticides should be free of detergents. Knowledge of formulation is particularly essential to learn how to improve the stability of bioinsecticides before they are deployed in the field; also, the lifespan of microbial insecticides must be evaluated before commercialization (Ramanujam et al. 2014).

The knowledge regarding the synchronization between the target insect bioecology, virus characteristics and the weather or climate of the agroecosystem is essential for the effective use of baculoviruses to control the susceptible hosts in the field. In addition, the application time is crucial for successful insect control. The life cycle of an insect host varies in different agroecological zones. Therefore, knowledge of the life cycle of an insect host should be part of the training program for farmers using biopesticides.

In conclusion, baculoviruses have great potential to be used as bioinsecticides. The vulnerability of baculoviruses to UV radiation could be decreased through the use of various types of protective agents. Natural ingredients must be considered as the primary option for UV protectors since the purpose of using bioinsecticides is to produce green agricultural produce and products for the consumer, and the natural ingredients must be cheap and easy to get. If chemical UV protectants are used in the viral formulations, the quantity of the chemicals in the formulation must follow the safety standards and guidelines, so that the formulations are safe for the environment and non-target organisms. Research on the recombinant UV-resistant baculoviruses is still limited. The virulence and photostability of the wild-type viruses are the important criteria for selection. The target of insects, habitat, indigenous virus used, and time of applications of bioinsecticides are also crucial as these factors will affect the efficacy of the viral bioinsecticides in the field. Quality assurance is important in the production of baculoviruses for bioinsecticide industry as to ensure a sustainable production of functional virus product for insect pest control.

ACKNOWLEDGEMENTS

The authors thank the Southeast Asia Ministers of Education Organisation (SEAMEO)-The Southeast Asian Regional Center for Graduate Study and Research in Agriculture (SEARCA) for supporting Ms. Erise Anggraini with a PhD Scholarship at Universiti Putra Malaysia under the SEARCA Scholarship. Also, we thank the anonymous reviewers for their careful reading of our paper and their many insightful comments and suggestions.

REFERENCES

- Abbas MST. 2018. Genetically engineered (Modified) crops (*Bacillus thuringiensis* crops) and the world controversy on their safety. *Egypt J Biol Pest Control* 28: 52. DOI: 10.1186/s41938-018-0051-2.
- Akhanaev YB, Belousova IA, Ershov NI, Nakai M, Martemyanov VV, Glupov VV. 2017. Comparison of tolerance to sunlight between spatially distant and genetically different strains of *Lymantria dispar* nucleopolyhedrovirus. *PLoS ONE* 12 (12): e0189992. DOI: 10.1371/journal.pone.0189992.
- Ali G, Vlak JM, van der Werf W. 2018. Biological activity of Pakistani isolate SpltNPV-Pak-BNG in second, third and fourth instar larvae of the leafworm *Spodoptera litura*. *Biocontrol Sci Technol* 28 (5): 521-527. DOI: 10.1080/09583157.2018.1461197.
- Amalfi S, Molina GN, Bevacqua RJ, López MG, Taboga O, Alfonso V. 2020. Baculovirus transduction in mammalian cells is affected by the production of type I and III interferons, which is mediated mainly by the cGAS-STING pathway. *J Virol* 94 (21): e01555. DOI: 10.1128/jvi.01555-20.
- Araújo JP, Hughes DP. 2019. Zombie-ant fungi emerged from non-manipulating, beetle-infecting ancestors. *Curr Biol* 29 (21): 3735-3738. DOI: 10.1016/j.cub.2019.09.004.
- Arora R, Shera P. 2014. Genetic improvement of biocontrol agents for sustainable pest management. *Basic and Applied Aspects of Biopesticides*. Springer, New Delhi. DOI: 10.1007/978-81-322-1877-7_15.
- Asano S. 2005. Ultraviolet protection of a granulovirus product using iron oxide. *Appl Entomol Zool* 40 (2): 359-364. DOI: 10.1303/aez.2005.359.
- Ayyub MB, Nawaz A, Arif MJ, Amrao L. 2019. Individual and combined impact of nuclear polyhedrosis virus and spinosad to control the tropical armyworm, *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae), in cotton in Pakistan. *Egypt J Biol Pest Control* 29: 67. DOI: 10.1186/s41938-019-0170-4.
- Bamisile BS, Akutse KS, Siddiqui JA, Xu Y. 2021. Model application of entomopathogenic fungi as alternatives to chemical pesticides: prospects, challenges, and insights for next-generation sustainable agriculture. *Front Plant Sci* 12: 741804. DOI: 10.3389/fpls.2021.741804.
- Barratt BI, Moran VC, Bigler F, Van Lenteren JC. 2018. The status of biological control and recommendations for improving uptake for the future. *BioControl* 63 (1): 155-167. DOI: 10.1007/s10526-017-9831-y.
- Beas-Catena A, Sánchez-Mirón A, García-Camacho F, Contreras-Gómez A, Molina-Grima E. 2014. Baculovirus biopesticides: An overview. *J Anim Plant Sci* 24 (2): 362-373.
- Becnel JJ, White SE. 2007. Mosquito pathogenic viruses - The last 20 years. *J Am Mosq Control Assoc* 23 (2 Suppl): 36-49. DOI: 10.2987/8756-971X(2007)23[36:MPVLY]2.0.CO;2
- Bedjo. 2017. The potential of various isolates of *Spodoptera litura* nuclear polyhedrosis viruses from east java (Indonesia) to control *Spodoptera litura* on soybean. *Biodiversitas* 18 (2): 582-588. DOI: 10.13057/biodiv/d180219.
- Behle RW, Popham HJR. 2012. Laboratory and field evaluations of the efficacy of a fast-killing baculovirus isolate from *Spodoptera frugiperda*. *J Invertebr Pathol* 109 (2): 194-200. DOI: 10.1016/j.jip.2011.11.002.
- Brassel J, Benz G. 1979. Selection of a strain of the granulosis virus of the codling moth with improved resistance against artificial ultraviolet radiation and sunlight. *J Invertebr Pathol* 33 (3): 358-363. DOI: 10.1016/0022-2011(79)90038-7.
- Bravo A, Likitvivatanavong S, Gill SS, Soberón M. 2011. *Bacillus thuringiensis*: A story of a successful bioinsecticide. *Insect Biochem Mol Biol* 41 (7): 423-431. DOI: 10.1016/j.ibmb.2011.02.006.
- Bull DL, Ridgway RL, House VS, Pryor NW. 1976. Improved formulations of the heliothis nuclear Polyhedrosis Virus1. *J Econ Entomol* 69 (6): 731-736. DOI: 10.1093/jee/69.6.731.
- Bullock HR, Hollingsworth JP, Hartstack Jr AW. 1970. Virulence of Heliothis Nuclear Polyhedrosis Virus to Monochromatic Ultraviolet Irradiation. *J Invertebr Pathol* 16 (3): 419-422. DOI: 10.1016/0022-2011(70)90161-8.
- Burand JP, Park EJ. 1992. Effect of nuclear polyhedrosis virus infection on the development and pupation of gypsy moth larvae. *J Invertebr Pathol* 60 (2): 171-175. DOI: 10.1016/0022-2011(92)90092-I.
- Cadet J, Grand A, Douki T. 2014. Solar UV Radiation-Induced DNA Bipyrimidine Photoproducts: Formation and Mechanistic Insights. *Photoinduced phenomena in nucleic acids II* Springer, Cham. DOI: 10.1007/128_2014_553.
- Çakmak T, Simón O, Kaydan MB, Tange DA, González Rodríguez AM, Piedra-Buena Díaz A, Caballero Murillo P, Hernández Suárez E. 2021. Effects of several UV-protective substances on the persistence of the insecticidal activity of the Alphabaculovirus of *Chrysodeixis chalcites* (ChcNPV-TF1) on banana (*Musa acuminata*, Musaceae, Colla) under laboratory and openfield conditions. *PLoS ONE* 16 (5): e0250217. DOI: 10.1371/journal.pone.0250217.
- Campbell JF. 2006. *Stored-Product Insect Behavior*. Second Edition, Insect Management for Food Storage and Processing: Second Edition. Second Edition. AACC International. DOI: 10.1016/B978-1-891127-46-5.50012-X.
- Cefali LC, Ataide JA, Moriel P, Foglio MA, Mazzola PG. 2016. Plant-based active photoprotectants for sunscreens. *Intl J Cosmet Sci* 38 (4): 346-353. DOI: 10.1111/ics.12316.
- Chidawanyika F, Mudavanhu P, Nyamukondiwa C. 2012. Biologically based methods for pest management in agriculture under changing climates: Challenges and future directions. *Insects* 3 (4): 1171-1189. DOI: 10.3390/insects3041171.
- Clem RJ, Passarelli AL. 2013. Baculoviruses: Sophisticated pathogens of insects. *PLoS Pathog* 9 (11): e1003729. DOI: 10.1371/journal.ppat.1003729.
- Cole C, Shyr T, Ou-Yang H. 2016. Metal oxide sunscreens protect skin by absorption, not by reflection or scattering. *Photodermatol Photoimmunol Photomed* 32 (1): 5-10. DOI: 10.1111/php.12214.
- Cuartas P, Barrera G, Barreto E, Villamizar L. 2014. Characterisation of a *Colombian granulovirus* (Baculoviridae: Betabaculovirus) isolated from *Spodoptera frugiperda* (Lepidoptera: Noctuidae) larvae. *Biocontrol Sci Technol* 24 (11): 1265-1285. DOI: 10.1080/09583157.2014.933312.
- D'Amico V, Slavicek J. 2012. Interactions Between Nucleopolyhedroviruses and Polydnviruses in Larval Lepidoptera. In: Adoga MP (eds). *Molecular Virology*. IntechOpen, London. DOI: 10.5772/35180.
- Demir I, Nalçacıoğlu R, Gholizad LM, Demirbağ Z. 2014. A highly effective nucleopolyhedrovirus against *Malacosoma* spp. (Lepidoptera: Lasiocampidae) from Turkey: Isolation, characterization, phylogeny, and virulence. *Turk J Agric For* 38 (4): 462-470. DOI: 10.3906/tar-1307-32.
- Dezianian A, Sajap AS, Lau WH, Omar D, Kadir HA, Mohamed R, Yusoh MR. 2010. Morphological characteristics of *P. xylostella* Granulovirus and effects on its larval host diamondback moth *Plutella xylostella* L. (Lepidoptera, Plutellidae). *Am J Agric Biol Sci* 5 (1): 43-49.
- Douki T, von Koschimbahr A, Cadet J. 2017. Insight in DNA repair of UV-induced pyrimidine dimers by chromatographic methods. *Photochem Photobiol* 93 (1): 207-215. DOI: 10.1111/php.12685.
- Duan LQ, Otvos IS, Xu LB, Conder N, Wang Y2012. Field testing chinese and japanese gypsy moth nucleopolyhedrovirus and disparvirus against a Chinese population of *lymantria dispar asiatica* in Huhhot, Inner Mongolia, People's Republic of China. *J Econ Entomol* 105 (2): 344-353. DOI: 10.1603/ec11294.
- Dunaway S, Odin R, Zhou L, Ji L, Zhang Y, Kadekaro AL. 2018. Natural antioxidants: Multiple mechanisms to protect skin from solar radiation. *Front Pharmacol* 24 (9): 392. DOI: 10.3389/fphar.2018.00392.
- Eischeid AC, Linden KG. 2011. Molecular indications of protein damage in adenoviruses after UV disinfection. *Appl Environ Microbiol* 77 (3): 1145-1147. DOI: 10.1128/AEM.00403-10.
- El-Helaly A. 2013. Moringa, rice bran as a new additives for a Baculovirus against ultraviolet effect. *Glob J Mul App Sci* 1 (1): 19-25. DOI: 10.4172/2161-0983.S1.004.
- El-Helaly A, Khatib M, El-Salamouny S, El-Sheikh M, Elnagar S. 2013. Promising additives to protect the activity of baculovirus biocontrol agent under field-sunlight conditions in Egypt. *J Life Sci* 7 (5): 495-500.
- El-Helaly A. 2020. Moringa water extract promising additive to prolong the activity of baculovirus under field-sunlight conditions in Egypt. *Braz J Biol* 80 (4): 891-896. DOI: 10.1590/1519-6984.225485.
- El-Helaly A, Sayed WAA. 2015. New approach based on nanotechnology in baculovirus protection. *Egypt Acad J Biol Sci F Toxicol Pest Control* 7 (1): 163-175. DOI: 10.21608/eajbsf.2015.17251.

- El-Sharkawey AZ, Ragaei M, Sabbour MM, Afaf AA, Mohamed HA, Samy R. 2009. Laboratory evaluation of antioxidants as UV-protectants for *Bacillus thuringiensis* against potato tuber moth larvae. *Aust J Basic Appl Sci* 3 (2): 358-370.
- Elder BD. 2013. Developing models of disease transmission: insights from ecological studies of insects and their baculoviruses. *PLoS Pathog* 9 (6): e1003372. DOI: 10.1371/journal.ppat.1003372.
- Elveborg S, Monteil V, Mirazimi A. 2022. Methods of inactivation of highly pathogenic viruses for molecular, serology or vaccine development purposes. *Pathogens* 11 (2): 271. DOI: 10.3390/pathogens11020271.
- Fernandes ME, Alves FM, Pereira RC, Aquino LA, Fernandes FL, Zanuncio JC. 2016. Lethal and sublethal effects of seven insecticides on three beneficial insects in laboratory assays and field trials. *Chemosphere* 156: 45-55. DOI: 10.1016/j.chemosphere.2016.04.115.
- Ferrelli ML, Pidre ML, Ghiringhelli PD, Torres S, Fabre ML, Masson T, Cédola MT, Sciocco-Cap A, Romanowski V. 2018. Genomic analysis of an Argentinean isolate of *Spodoptera frugiperda* granulovirus reveals that various baculoviruses code for Lef-7 proteins with three F-box domains. *PLoS ONE* 13 (8): e0202598. DOI: 10.1371/journal.pone.0202598.
- Gerba CP, Gramos DM, Nwachuku N. 2002. Comparative inactivation of enteroviruses and adenovirus 2 by UV light. *Appl Environ Microbiol* 68 (10): 5167-5169. DOI: 10.1128/AEM.68.10.5167-5169.2002.
- Gifani A, Marzban R, Safekordi A, Ardjmand M, Dezianian A. 2015. Ultraviolet protection of nucleopolyhedrovirus through microencapsulation with different polymers. *Biocontrol Sci Technol* 25 (7): 814-827. DOI: 10.1080/09583157.2015.1018814.
- Goulson D. 1997. Wipfelkrankheit: Modification of host behaviour during baculoviral infection. *Oecologia* 109 (2): 219-228. DOI: 10.1007/s004420050076.
- Goulson D, Martinez AM, Hughes WO, Williams T. 2000. Effects of optical brighteners used in biopesticide formulations on the behavior of pollinators. *Biol Control* 19 (3): 232-236. DOI: 10.1006/bcon.2000.0861.
- Gröbner J. 2013. Ultraviolet radiation: Distribution and variability. In *Environmental Toxicology. Encyclopedia of Sustainability Science and Technology*. Springer, New York, NY.
- Grzywacz D, Stevenson PC, Mushobozi WL, Belmain S, Wilson K. 2014. The use of indigenous ecological resources for pest control in Africa. *Food Secur* 6 (1): 71-86. DOI: 10.1007/s12571-013-0313-5.
- Haase S, Sciocco-Cap A, Romanowski V. 2015. Baculovirus insecticides in Latin America: Historical overview, current status and future perspectives. *Viruses* 7 (5): 2230-2267. DOI: 10.3390/v7052230.
- Hamdi S. 2019. Analysis of ultraviolet index, ultraviolet B insolation, and sunshine duration at Bandung in year 2017. *IOP Conf Ser: Earth Environ Sci* 303 (1): 012018. DOI: 10.1088/1755-1315/303/1/012018.
- Harrison R, Hoover K. 2012a. Baculoviruses and other occluded insect viruses. *Insect Pathology*. In: Vega FE, Kaya HK (eds). *Insect Pathology* (Second Edition). Academic Press, Camb. DOI: 10.1016/B978-0-12-384984-7.00004-X.
- Harrison R, Hoover K. 2012b. Baculoviruses and other occluded insect viruses. Second Edi, *Insect Pathology*. Second Edi. Elsevier Inc., Amsterdam. DOI: 10.1016/B978-0-12-384984-7.00004-X.
- Hefferon KL, Oomens AG, Monsma SA, Finnerty CM, Blissard GW. 1999. Host cell receptor binding by baculovirus GP64 and kinetics of virion entry. *Virology* 258 (2): 455-468. DOI: 10.1006/viro.1999.9758.
- Herlinda S, Oktareni SS, Suparman S, Anggarini E, Elfita E, Setiawan A, Verawaty M, Hasbi H, Lakitan B. 2020. Effect of application of UV irradiated *Beauveria bassiana* and *Metarhizium anisopliae* on larval weight and mortality of *Spodoptera litura*. *Adv Biol Sci Res* 8 (Iccesi 2019): 64-70. DOI: 10.2991/absr.k.200513.011.
- Hillel D, Hatfield JL, Powlson DS, Rosenzweig C, Scow KM, Singer MJ, Sparks DL. 2008. *Solar Radiation*. Environmental Sciences: A Student's Companion.
- Hockberger PE. 2002. A history of ultraviolet photobiology for humans, animals and microorganisms. *Photochem Photobiol* 76 (6): 561-79. DOI: 10.1562/0031-8655(2002)0760561AHOUFP2.0.CO2.
- Hoover K, Grove M, Gardner M, Hughes DP, McNeil J, Slavicek J. 2011. A gene for an extended phenotype. *Science* 333 (6048): 1401. DOI: 10.1126/science.1209199.
- Hoy CW. 2019. Insect control in the field using temperature extremes. In *Temperature sensitivity in insects and application in integrated pest management*. CRC Press. DOI: 10.1201/9780429308581-10.
- Ibarra JE, Del Rincón-Castro MC. 2008. Insect viruses diversity, biology, and use as bioinsecticides. *Trop Biol Conserv Manag* 5: 1-10.
- Ibrahim SA, El-Banna HMS, El-Kareem SMIA. 2019. Determination of the effect of some plant extracts as the UV-Protectant to the American bollworm, *Helicoverpa armigera* (Hübner) nucleopolyhedrovirus (Ha NPV) virulence under laboratory conditions. *IOSR J Agric Vet Sci* 12 (12): 10-14.
- Ignoffo CM, Hostetter DL, Sikorowski PP, Sutter G, Brooks WM. 1977. Inactivation of representative species of entomopathogenic viruses, a bacterium, fungus, and protozoan by an ultraviolet light source. *Environ Entomol* 6 (3): 411-415. DOI: 10.1093/ee/6.3.411.
- Ikehata H, Ono T. 2011. The mechanisms of UV mutagenesis. *J Radiat Res* 52 (2): 115-125. DOI: 10.1269/jrr.10175.
- Jackson MA, Dunlap CA, Jaronski ST. 2010. Ecological considerations in producing and formulating fungal entomopathogens for use in insect biocontrol. *BioControl* 55 (1): 129-145. DOI: 10.1007/s10526-009-9240-y.
- Javier-Hila AMV, Caoili BL. 2020. Cross-infectivity of a putative *Spodoptera picta* nucleopolyhedrovirus to *Spodoptera litura* fabricius (Lepidoptera: Noctuidae). *Philippine J Sci* 149 (3): 887-896.
- Jehle JA, Blissard GW, Bonning BC, Cory JS, Herniou EA, Rohrmann GF, Theilmann DA, Thiem SM, Vlak JM. 2006. On the classification and nomenclature of baculoviruses: A proposal for revision. *Arch Virol* 151 (7): 1257-1266. DOI: 10.1007/s00705-006-0763-6.
- Jouzani GS, Valijanian E, Sharafi R. 2017. *Bacillus thuringiensis*: a successful insecticide with new environmental features and tidings. *Appl Microbiol Biotechnol* 101 (7): 2691-2711. DOI: 10.1007/s00253-017-8175-y.
- Kaiser D, Bacher S, Mène-Saffrané L, Grabenweger G. 2019. Efficiency of natural substances to protect *Beauveria bassiana* conidia from UV radiation. *Pest Manag Sci* 75 (2): 556-563. DOI: 10.1002/ps.5209.
- Kalha CS, Singh PP, Kang SS, Hunjan MS, Gupta V, Sharma R. 2014. Entomopathogenic viruses and bacteria for insect-pest control, integrated pest management: current concepts and ecological perspective. Elsevier Inc., Amsterdam. DOI: 10.1016/B978-0-12-398529-3.00013-0.
- Kato T, Suzuki F, Park EY. 2011. Purification of functional baculovirus particles from silkworm larval hemolymph and their use as nanoparticles for the detection of human prorenin receptor (PRR) binding. *BMC Biotechnol* 11 (1): 10. DOI: 10.1186/1472-6750-11-60.
- Khachatourians G, Qazi S. 2008. Human and Animal Relationships. *Human and Animal Relationships*. DOI: 10.1007/978-3-540-79307-6.
- Khorramnejad A, Perdomo HD, Palatini U, Bonizzoni M, Gasmi L. 2021. Cross talk between viruses and insect cells cytoskeleton. *Viruses* 13 (8): 1658. DOI: 10.3390/v13081658.
- Killick HJ. 1990. Influence of droplet size, solar ultraviolet light and protectants, and other factors on the efficacy of baculovirus sprays against *Panolis flammea* (Schiff.) (Lepidoptera: Noctuidae). *Crop Protection* 9 (1): 21-28. DOI: 10.1016/0261-2194(90)90041-5.
- Killick HJ, Warden SJ. 1991. Ultraviolet penetration of pine trees and insect virus survival. *Entomophaga* 36 (1): 87-94. DOI: 10.1007/BF02374639.
- Kleinermaans K, Nachtigallova D, de Vries MS. 2013. Excited state dynamics of DNA bases. *Intl Rev Phys Chem* 32 (2): 308-342. DOI: 10.1080/0144235X.2012.760884.
- Korać RR, Khambholja KM. 2011. Potential of herbs in skin protection from ultraviolet radiation. *Pharmacogn Rev* 5 (10): 164-173. DOI: 10.4103/0973-7847.91114.
- Krisnawati A, Nuryati N, Adie MM. 2021. Assessment of soybean resistance to leaf-feeding insect, *Spodoptera litura*. *Biosaintifika J Biol Biol Educ* 13 (1): 92-98. DOI: 10.15294/biosaintifika.v13i1.24794.
- Kukan B. 1999. MINIREVIEW Vertical Transmission of Nucleopolyhedrovirus in Insects. *J Invertebr Pathol* 74 (1999): 103-111. DOI: 10.1006/jipa.1999.4873.
- Kumari S, Chourasia SK, Jha MN. 2014. Microbial Pesticide: A Boom For Sustainable Agriculture. *Intl J Sci Eng Res* 5 (6): 1394-1397.
- Kvakkestad V, Sundbye A, Gwynn R, Klingen I. 2020. Authorization of microbial plant protection products in the Scandinavian countries: A comparative analysis. *Environ Sci Policy* 106: 115-124. DOI: 10.1016/j.envsci.2020.01.017.
- Lakhanpal P, Rai DK. 2007. Quercetin: a versatile flavonoid. *Internet J Med Update - EJOURNAL* 2 (2): 22-37. DOI: 10.4314/ijmu.v2i2.39851.

- Landwehr A. 2021. Benefits of baculovirus use in IPM strategies for open field and protected vegetables. *Front Sustain Food Syst* 4: 593796. DOI: 10.3389/fsufs.2020.593796.
- Lasa R, Ruiz-Portero C, Alcázar MD, Belda JE, Caballero P, Williams T. 2007. Efficacy of optical brightener formulations of *Spodoptera exigua* multiple nucleopolyhedrovirus (SeMNPV) as a biological insecticide in greenhouses in southern Spain. *Biol Control* 40 (1): 89-96. DOI: 10.1016/j.biocontrol.2006.06.015.
- Li N, Deng L, Xiang L, Liang Y. 2014. Photoprotective effect of tea and its extracts against ultraviolet radiation-induced skin disorders. *Trop J Pharm Res* 13 (3): 475-483. DOI: 10.4314/tjpr.v13i3.24.
- Lubin D, Jensen EH. 1995. Effects of clouds and stratospheric ozone depletion on ultraviolet radiation trends. *Nature* 377 (6551): 710-713. DOI: 10.1038/377710a0.
- Lyons AB, Trullas C, Kohli I, Hamzavi IH, Lim HW. 2020. Photoprotection beyond ultraviolet radiation: A review of tinted sunscreens. *J Am Acad Dermatol* 84 (5): 1393-1397. DOI: 10.1016/j.jaad.2020.04.079.
- Malik MA, Ahmad SJ, Arif MJ, Ahmad JN. 2020. Management of diamond back moth. *Pak J Zool* 52 (2): 641-647. DOI: 10.17582/journal.pjz/20190808100805.
- Maqsood S, Afzal M, Aqueel MA, Raza AB, Wakil W, Babar MH. 2017. Efficacy of nuclear polyhedrosis virus and flubendiamide alone and in combination against *Spodoptera litura* F. *Pak J Zool* 49 (5): 1783-1788. DOI: 10.17582/journal.pjz/2017.49.5.1783.1788.
- Maramorosch K. 2012. Viral insecticides for biological control. Elsevier, Amsterdam.
- Martin S, Domek J. 2002. Relative effects of ultraviolet and visible light on the activities of corn earworm and beet armyworm (Lepidoptera: Noctuidae) nucleopolyhedroviruses. *J Econ Entomol* 95 (2): 261-268. DOI: 10.1603/0022-0493.95.2.261.
- Martínez AM, Velasco S, Méndez A, Figueroa JI, España ML, Cárdenas-Navarro R, Pineda S. 2009. Effects of optical brighteners used in biopesticide formulations on crops: reflectance, stomatal conductance, photosynthesis, and growth. *Commun Agric Appl Biol Sci* 74 (1): 117-123.
- Matsunaga T, Hieda K, Nikaido O. 1991. Wavelength dependent formation of thymine dimers and (6-4) photoproducts in DNA by monochromatic ultraviolet light ranging from 150 to 365 nm. *Photochem Photobiol* 54 (3): 403-410. DOI: 10.1111/j.1751-1097.1991.tb02034.x.
- Miele SA, Garavaglia MJ, Belaich MN, Ghiringhelli PD. 2011. Baculovirus: Molecular insights on their diversity and conservation. *Intl J Evol Biol* 2011: 379424. DOI: 10.4061/2011/379424.
- Mondragón G, Pineda S, Martínez A, Martínez AM. 2007. Optical brightener Tinopal C1101 as an ultraviolet protectant for a nucleopolyhedrovirus. *Commun Agric Appl Biol Sci* 72 (3): 543-547.
- Monteiro F, Carinhas N, Carrondo MJ, Bernal V, Alves PM. 2012. Toward system-level understanding of baculovirus-host cell interactions: From molecular fundamental studies to large-scale proteomics approaches. *Front Microbiol* 3: 391. DOI: 10.3389/fmicb.2012.00391.
- Mouret S, et al. 2006. Cyclobutane pyrimidine dimers are predominant DNA lesions in whole human skin exposed to UVA radiation. *Proc Nat Acad Sci U S A* 103 (37): 13765-13770. DOI: 10.1073/pnas.0604213103.
- Mwanza P, Jukes M, Dealtry G, Lee M, Moore S. 2022. Selection for and Analysis of UV-Resistant *Cryptophlebia leucotreta* Granulovirus-SA as a Biopesticide for *Thaumatococcus leucotreta*. *Viruses* 14 (1): 28. DOI: 10.3390/v14010028.
- Nai YS, Wu CY, Wang TC, Chen YR, Lau WH, Lo CF, Tsai MF, Wang CH. 2010. Genomic sequencing and analyses of *Lymantria xylin* multiple nucleopolyhedrovirus. *BMC Genomics* 11: 116. DOI: 10.1186/1471-2164-11-116.
- Natkar PK, Balikai RA. 2017. Present status on bio-ecology and management of tobacco caterpillar, *Spodoptera litura* (Fabricius) - An update. *Intl J Plant Prot* 10 (1): 193-202. DOI: 10.15740/HAS/IJPP/10.1/193-202.
- Nawaz A, Ali H, Sufyan M, Gogi MD, Arif MJ, Ranjha MH, Arshid M, Waseem M, Mustafa T, Qasim M, Rizwan M. 2019. Comparative bio-efficacy of nuclear polyhedrosis virus (NPV) and Spinosad against American bollworm, *Helicoverpa armigera* (Hubner). *Revista Brasileira de Entomologia* 63 (4): 277-282. DOI: 10.1016/j.rbe.2019.09.001.
- Nazli-Huda, Sajap AS, Lau WH. 2012. Stability of *Spodoptera litura* nucleopolyhedrovirus in sodium dodecyl sulphate. *Afr J Biotechnol* 11 (16): 3877-3881. DOI: 10.5897/AJB11.2848.
- Nishino H, Iwashima A, Fujiki H, Sugimura T. 1984. Inhibition by quercetin of the promoting effect of teleocidin on skin papilloma formation in mice initiated with 7, 12- dimethylbenz [a] anthracene. *GANN Jpn J Cancer Res* 75 (2): 113-116.
- van Oers MM. 2021. Baculoviruses: Molecular Biology and Replication (Baculoviridae), Encyclopedia of Virology. Elsevier Ltd., Amsterdam. DOI: 10.1016/b978-0-12-809633-8.21550-1.
- Petrik DT, Iseli A, Montelone BA, Van Etten JL, Clem RJ. 2003. Improving baculovirus resistance to UV inactivation: Increased virulence resulting from expression of a DNA repair enzyme. *J Invertebr Pathol* 82 (1): 50-56. DOI: 10.1016/s0022-2011(02)00197-0.
- Podgwaite JD. 1985. Strategies for field use of baculoviruses. In: Maramorosch K, Sherman KE (eds.) *Viral insecticides for biological control*. Academic Press, Waltham, MA. DOI: 10.1016/B978-0-12-470295-0.50028-8.
- Prabhu S, Mahalingam C. 2017. Effect of sunlight and UV light against DpNPV (Nuclear Polyhedrosis Virus) formulation on larval mortality of mulberry leaf webber, *Diaphania pulverulentalis* Hampson. *Intl J Curr Microbiol Appl Sci* 6 (3): 1897-1905. DOI: 10.20546/ijcmas.2017.603.215.
- Ramanujam B, Rangeshwaran R, Sivakumar G, Mohan M, Yandigeri MS. 2014. Management of insect pests by microorganisms. *Proc Indian Natl Sci Acad* 80 (2): 455-471. DOI: 10.16943/ptinsa/2014/v80i2/3.
- Rastogi RP, Kumar A, Tyagi MB, Sinha RP. 2010. Molecular mechanisms of ultraviolet radiation-induced DNA damage and repair. *J Nucleic Acids* 2010: 592980 DOI: 10.4061/2010/592980.
- Rathore S, Chandola M, Raghuvanshi R, Kaur M, Singh KV. 2021. Indigenous pest management practices of indian hill farmers: Introspecting their rationale and communication pattern for secure ecosystems. *Sustainability (Switzerland)* 13 (21): 11608. DOI: 10.3390/su132111608.
- Rebolledo D, Lasa R, Guevara R, Murillo R, Williams T. 2015. Baculovirus-induced climbing behavior favors intraspecific necrophagy and efficient disease transmission in *Spodoptera exigua*. *PLoS ONE* 10 (9): e0136742. DOI: 10.1371/journal.pone.0136742.
- Redman EM, Wilson K, Cory JS. 2016. Trade-offs and mixed infections in an obligate-killing insect pathogen. *J Anim Ecol* 85 (5): 1200-1209. DOI: 10.1111/1365-2656.12547.
- Rohrmann G. 2019. Structural proteins of baculovirus occlusion bodies and virions. In *Baculovirus Molecular Biology* [Internet]. 4th edition. National Center for Biotechnology Information (US).
- Ros VI, van Houte S, Hemerik L, van Oers MM. 2015. Baculovirus-induced tree-top disease: How extended is the role of egt as a gene for the extended phenotype? *Mol Ecol* 24 (1): 249-258. DOI: 10.1111/mec.13019.
- Ruii L. 2018. Microbial biopesticides in agroecosystems. *Agronomy* 8 (11): 235. DOI: 10.3390/agronomy8110235.
- Saewan N, Jimtaisong A. 2013. Photoprotection of natural flavonoids. *J Appl Pharm Sci* 3 (9): 129-141.
- Sajap AS, Kotulai JR, Bakir MA, Hong LW, Samad MA, Kadir HA. 2000. Pathogenicity and characteristics of *Spodoptera litura* Nucleopolyhedrovirus from Peninsular Malaysia. *Pertanika J Trap Agric Sci* 23 (1): 23-28.
- Sajjan DB, Hinchigeri SB. 2016. Structural organization of baculovirus occlusion bodies and protective role of multilayered polyhedron envelope protein. *Food Environ Virol* 8 (1): 86-100. DOI: 10.1007/s12560-016-9227-7.
- Salama MS, Abd El-Salam AM, Mahmoud DM, Samah MM. 2017. Effect of ultra violet radiations on insecticidal activity of *Spodoptera littoralis* multinucleocapsid nuclear polyhedrosis virus against *Spodoptera littoralis* Bois (Lepidoptera: Noctuidae). *Biosci Res* 14 (3): 645-652.
- Salamouny ES, Shapiro M, Ling KS, Shepard BM. 2009. Black tea and lignin as ultraviolet protectants for the beet armyworm nucleopolyhedrovirus 1. *J Entomol Sci* 44 (1): 50-58. DOI: 10.18474/0749-8004-44.1.50.
- Samada LH, Tambunan USF. 2020. Biopesticides as promising alternatives to chemical pesticides: A review of their current and future status. *OnLine J Biol Sci* 20 (2): 66-76. DOI: 10.3844/ojbsci.2020.66.76.
- Samsudin S, Santoso T, Rauf A, Kusumah YM. 2011. Keefektifan bahan pelindung alami dalam mempertahankan infektifitas *Spodoptera*

- exigua* Nucleopolyhedrovirus (SeNPV). Berita Biologi 10 (6): 689-697. [Indonesian]
- Saxena A, Byram PK, Singh SK, Chakraborty J, Murhammer D, Giri L. 2018. A structured review of baculovirus infection process: Integration of mathematical models and biomolecular information on cell-virus interaction. J Gen Virol 99 (9): 1151-1171. DOI: 10.1099/jgv.0.001108.
- Sayed WAA, El-Bendary HK, El-Helal A. 2020. Increasing the efficacy of the cotton leaf worm *Spodoptera littoralis* nucleopolyhedrosis virus using certain essential oils. Egypt J Biol Pest Control 30 (1): 1-7. DOI: 10.1186/s41938-019-0201-1.
- Schmucki DA, Philipona R. 2002. Ultraviolet radiation in the Alps: the altitude effect. Optical Eng 41 (12): 3090. DOI: 10.1117/1.1516820.
- Senthil-Nathan S. 2015. A review of biopesticides and their mode of action against insect pests. Environ Sustain 49-63. DOI: 10.1007/978-81-322-2056-5_3.
- Sigstam T, Gannon G, Cascella M, Pecson BM, Wigginton KR, Kohn T. 2013. Subtle differences in virus composition affect disinfection kinetics and mechanisms. Appl Environ Microbiol 79 (11): 3455-3467. DOI: 10.1128/AEM.00663-13.
- Singh D, Raina TK, Singh J. 2017. Entomopathogenic fungi: An effective biocontrol agent for management of insect populations naturally. J Pharm Sci Res 9 (6): 830-839.
- Slavicek JM, Popham HJR. 2005. The *Lymantria dispar* Nucleopolyhedrovirus enhancers are components of occlusion-derived virus. J Virol 79 (16): 10578-10588. DOI: 10.1128/JVI.79.16.10578-10588.2005.
- Smaghe G, Deknopper J, Meeus I, Mommaerts V. 2013. Dietary chlorantraniliprole suppresses reproduction in worker bumblebees. Pest Manag Sci 69 (7): 787-791. DOI: 10.1002/ps.3504.
- Smijts TG, Pavel S. 2011. Titanium dioxide and zinc oxide nanoparticles in sunscreens: Focus on their safety and effectiveness. Nanotechnol Sci Appl 4 (1): 95-112. DOI: 10.2147/NSA.S19419.
- Smit, K. 1966. Physical and chemical changes induced in nucleic acids by ultraviolet light. Radiat Res Suppl 6: 54-79. DOI: 10.2307/3583551.
- Smith K, Hanawalt P. 2013. Molecular photobiology: inactivation and recovery. Elsevier.
- Sood P, Mehta PK, Prabhakar CS. 2013. Effect of UV protectants on the efficacy of *Pieris brassicae* granulovirus. Biol Agric Hortic 29 (2): 69-81. DOI: 10.1080/01448765.2013.763449.
- Steinhauer DA, Domingo E, Holland JJ. 1992. Lack of evidence for proofreading mechanisms associated with an RNA virus polymerase. Gene 122 (2): 281-288. DOI: 10.1016/0378-1119(92)90216-C.
- Sujayanand GK, Akram M, Ashish N, Aravind K, Shripad B, Kumar SD, Raja H, Sonika P. 2019. Virulence, cross infectivity and molecular characterization of *Spilosoma obliqua* MNPV from north India for Bihar hairy caterpillar management. Res J Biotechnol 14 (12): 58-70.
- Sukirno S, Tufail M, Rasool KG, El Salamouny S, Sutanto KD, Aldawood AS. 2018. The Efficacy and Persistence of *Spodoptera littoralis* Nucleopolyhedrovirus (SpliMNPV) Applied in UV Protectants against the Beet Armyworm, *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae) under Saudi Field Conditions. Pak J Zool 50 (5). DOI: 10.17582/journal.pjz/2018.50.5.1895.1902.
- Sun X. 2015. History and current status of development and use of viral insecticides in China. Viruses 7 (1): 306-319. DOI: 10.3390/v7010306.
- Sunil K, Katti P, Nadagouda S, Hanchinal SG. 2019. Life table evaluation of *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) on Groundnut. Ann Plant Soil Res 21 (9): 1-12.
- Supyani SS, Noviayanti PN, Wijayanti RW. 2014. Insecticidal properties of *Spodoptera exigua* nuclear polyhedrosis virus local isolate against *Spodoptera exigua* on shallot. Intl J Entomol Res 2 (3): 175-180.
- Susurluk H, Toprak U, Gürkan MO. 2013. Concentration of sodium dodecyl sulfate used in occlusion body extraction affects *Spodoptera littoralis* nucleopolyhedrovirus biological activity. Turk J Biol 37 (2): 171-175. DOI: 10.3906/biy-1204-50.
- Sutanto KD, El Salamouny S, Tufail M, Ghulam Rasool K, Sukirno S, Shepard M, Shapiro M, Saad Aldawood A. 2017. Evaluation of natural additives to enhance the persistence of *Spodoptera littoralis* (Lepidoptera: Noctuidae) Nucleopolyhedrovirus (SpliMNPV) under field conditions in Saudi Arabia. J Econ Entomol 110 (3): 924-930. DOI: 10.1093/jee/tox085.
- Szewczyk B, Hoyos-Carvajal L, Paluszek M, Skrzecz I, De Souza ML. 2006. Baculoviruses - Re-emerging biopesticides. Biotechnol Adv 24 (2): 143-160. DOI: 10.1016/j.biotechadv.2005.09.001.
- Tabashnik BE, Brévault T, Carrière Y. 2013. Insect resistance to Bt crops: Lessons from the first billion acres. Nat Biotechnol 31 (6): 510-521. DOI: 10.1038/nbt.2597.
- Takahashi M, Nakai M, Saito Y, Sato Y, Ishijima C, Kunimi Y. 2015. Field efficacy and transmission of fast- and slow-killing nucleopolyhedroviruses that are infectious to *Adoxophyes honmai* (Lepidoptera: Tortricidae). Viruses 7 (3): 1271-1283. DOI: 10.3390/v7031271.
- Tamez-Guerra P, Rodriguez-Padilla C, Galán-Wong LJ. 2005. Solar radiation (UV) protectants for microbial insecticides. DOI: 10.1021/bk-2005-0906.ch010.
- Tay WT, Mahon RJ, Heckel DG, Walsh TK, Downes S, James WJ, Lee SF, Reineke A, Williams AK, Gordon KH. 2015. Insect Resistance to *Bacillus thuringiensis* Toxin Cry2Ab Is Conferred by Mutations in an ABC Transporter Subfamily A Protein. PLoS Genet 11 (11): e1005534. DOI: 10.1371/journal.pgen.1005534.
- Terra WR. 2001. The origin and functions of the insect peritrophic membrane and peritrophic gel. Arch Insect Biochem Physiol 47 (2): 47-61. DOI: 10.1002/arch.1036.
- Tetreau G. 2018. Interaction between insects, toxins, and bacteria: Have we been wrong so far? Toxins 10 (7): 1-13. DOI: 10.3390/toxins10070281.
- Trizelia, Armon N, Jailani H. 2015. The diversity of entomopathogenic fungi on rhizosphere of various vegetable crops. Pros Sem Nas Masy Biodiv Indon 1: 998-1004. DOI: 10.13057/psnmibi/m010307.
- Usta C. 2013. Microorganisms in biological pest control — a review (Bacterial Toxin Application and Effect of Environmental Factors). Curr Prog Biol Res 13: 287-317. DOI: 10.5772/55786.
- Wang M, Hu Z. 2019. Cross-talking between baculoviruses and host insects towards a successful infection. Philos Trans R Soc B: Biol Sci 374 (1767): 20180324. DOI: 10.1098/rstb.2018.0324.
- Wigginton KR, Pecson BM, Sigstam T, Bosshard F, Kohn T. 2012. Virus inactivation mechanisms: Impact of disinfectants on virus function and structural integrity. Environ Sci Technol 46 (21): 12069-12078. DOI: 10.1021/es3029473.
- Wilbur HM. 1980. Complex Life Cycles. Annu Rev Ecol Syst 11 (1): 67-93. DOI: 10.1146/annurev.es.11.110180.000435.
- Williams T, Virto C, Murillo R, Caballero P. 2017. Covert infection of insects by baculoviruses. Front Microbiol 8: 1-13. DOI: 10.3389/fmicb.2017.01337.
- Williams T. 2017. Viruses. Ecol Invertebr Dis 213-285. DOI: 10.1002/9781119256106.ch7.
- Wilson K, Grzywacz D, Curcic I, Scoates F, Harper K, Rice A, Paul N, Dillon A. 2020. A novel formulation technology for baculoviruses protects biopesticide from degradation by ultraviolet radiation. Sci Rep 10 (1): 1-10. DOI: 10.1038/s41598-020-70293-7.
- Witt DJ, Stairs GR. 1975. The effects of ultraviolet irradiation on a Baculovirus infecting *Galleria mellonella*. J Invertebr Pathol 26 (3): 321-327. DOI: 10.1016/0022-2011(75)90229-3.
- Wu ZW, Fan JB, Yu H, Wang D, Zhang YL. 2015. Ultraviolet protection of the *Cydia pomonella* granulovirus using zinc oxide and titanium dioxide. Biocontrol Sci Technol 25 (1): 97-107. DOI: 10.1080/09583157.2014.951029.
- Yong-Bing X, Gui-Lin C, Ming-Quan G. 2019. Antioxidant and anti-inflammatory activities of the crude extracts of *Moringa oleifera* from Kenya and their correlations with flavonoids. Antioxidants 8 (8). DOI: 10.3390/antiox8080296.
- Young SY, Yearian WC. 1974. Persistence of *Heliothis*. Environ Entomol 3 (2): 253-255. DOI: 10.1093/ee/3.2.253.
- Zhang L, Zhang X, Zhang Y, Wu S, Gelbič I, Xu L, Guan X. 2016. A new formulation of *Bacillus thuringiensis*: UV protection and sustained release mosquito larvae studies. Sci Rep 6 (1): 1-8. DOI: 10.9734/JSRR/2016/11862.
- Zhao Y, Chapman DAG, Jones IM. 2003. Improving baculovirus recombination. Nucleic Acids Res 31 (2). DOI: 10.1093/nar/ngn006.
- Zhu YC, Lu HX, He DH, Yang ZR. 2013. Synthesis, fluorescence properties and applications of two novel oxadiazole-based stilbene optical brighteners as UV protectants for insect baculovirus. J Photochem Photobiol B: Biol 125: 8-12. DOI: 10.1016/j.jphotobiol.2013.04.006.
- Zogo B, Tchikoi BN, Koffi AA, Dahounto A, Ahoua Alou LP, Dabiré RK, Baba-Moussa L, Moiroux N, Penetier C. 2019. Impact of sunlight exposure on the residual efficacy of bioinsecticides *Bacillus thuringiensis* israelensis and *Bacillus sphaericus* against the main malaria vector, *Anopheles gambiae*. Malaria J 18 (1): 1-9. DOI: 10.1186/s12936-019-2687-0.