

Plant diversity after sixteen years post coal mining in East Kalimantan, Indonesia

LIRIS LIS KOMARA, DEVI NANDITA CHOESIN, TATI SURYATI SYAMSUDIN

School of Life Sciences and Technology, Institut Teknologi Bandung, Jalan Ganesa No. 10, Bandung 40132, West Java, Indonesia.

Tel.: +62-22-2511575; +62-22-250 0258, Fax.: +62-22-2534107, email: liskomara@yahoo.co.id

Manuscript received: 19 December 2015. Revision accepted: 25 June 2016.

Abstract. Komara LL, Choesin DN, Syamsudin TS. 2016. Plant diversity after sixteen years post coal mining in East Kalimantan, Indonesia. *Biodiversitas* 17: 531-538. Post coal mining areas need to be rehabilitated through reclamation and revegetation. The objective of this study was to evaluate plant diversity after 16 years of revegetation activities in a coal mining reclamation site in East Kalimantan. In an effort to restore plant diversity, the coal mining company began by planting fast growing species as pioneers, then planting local species after three years. This study compared a 20 hectare reclamation site with conditions in the pre-mining area, which covered 14,988 hectares. Vegetation sampling was conducted in 20 plots measuring 20x20 m² along line transects, with 100 m distance between plots. A total of 104 plant species were found in the reclamation site, consisting of 76 tree species and 28 herbaceous species. Tree species consisted of 35 planted local species (e.g., *Dryobalanops aromatica*, *Eusideroxylon zwageri*, *Macaranga gigantea*), 25 planted non-local species, and 16 local species that grew spontaneously (e.g., *Leucaena glauca*, *Lansium domesticum*, *Shorea laevis*). In comparison, 133 species were found in the pre-mining site, consisting of 132 local tree species, one non-local tree species (*Acacia mangium*) and 52 herbaceous species. Tree species diversity index in the reclamation site after 16 years post mining (i.e., 3.54) was still lower than in the pre-mining area (4.29); while the diversity indices for herbaceous plants were relatively similar (2.97 and 2.67 in the reclamation and pre-mining sites respectively). The slightly higher diversity of herbaceous plants in the reclamation site may be attributed to higher coverage per species in this site, despite lower species richness.

Keywords: Kalimantan, local species, plant diversity, reclamation, revegetation

INTRODUCTION

Plant species composition in an area depends on environmental factors, such as temperature, humidity, nutrition, sunlight, topography, bedrock geology, soil characteristics, canopy structure and land use history (Hutchinson et al. 1999). Tropical rain forests are the most important plant species diversity centers in the world (Turner 2004). Situated in the tropics, the island of Borneo, including within it the Indonesian provinces of Kalimantan, contains a highly diverse flora, due partly to its unique geological and climatic history. The island is known to have 15,000 flowering plant species, and more than 3,000 species of trees, including 267 Dipterocarpaceae species (WWF 2005). However, this high diversity is currently being threatened by human activity and changes in land use.

Coal mining is a major activity that is changing landscapes. Coal mining causes changes in biodiversity (Cooke and Johnson 2002); soil profile (Makineci et al. 2011) and geological structure permanently (Shrestha and Lal 2011) by leaving large overburden areas (Graham and Haynes 2004; Sheoran and Sheoran 2009; Alday et al. 2011). Considering the impact of mining on the environment, post-mining areas need to be rehabilitated by conducting reclamation and revegetation (Hazarika et al. 2006). Reclamation in post coal mining areas involves moving the overburden to its original contour and spreading top soil over it (Shrestha and Lal 2011; Malakar

et al. 2015). After the reclamation process is completed, the reclamation site is then ready to be revegetated (Shrestha and Lal 2011; Wardana 2008). At reclamation sites, soil nutrients are generally limited, soil pH is low, and there are often metal contaminants; therefore, revegetation activity must be carried out with plants selected on the basis of their ability to survive and regenerate or reproduce under severe conditions. Normally the revegetation process is started by selecting plants that are resistant to drought, or fast growing crops or fodder which can grow with limited nutrients (Sheoran et al. 2010).

In East Kalimantan province, Indonesia, coal mining covers an area of almost 3.27 million hectares (Nugroho and Adman 2011). Previous reclamation efforts in Kalimantan have shown that directly planting local tree species in reclamation sites is not successful, compared to planting pioneer plants such as *Acacia mangium*, *Paraserianthes falcataria* (Mansur 2010). According to Indonesian regulation, the success of post mining reclamation is indicated by 90% growth of vegetation and vegetation conditions that are close to pre-mining conditions (Regulation of State Minister for The Environment no. 4/2012). According to Claassen et al. (2008) one of the indicators of reclamation success is the presence of vegetation. Specifically, Perrow and Davy (2002) mention plant species composition and richness as criteria for evaluating the success of restoration. These parameters are easy to measure and are quite sensitive (Dale and Beyeler 2001; Ludwig et al. 2003). The objective

of this study was to evaluate plant diversity after 16 years of revegetation activities in a coal mining reclamation site in East Kalimantan.

MATERIALS AND METHODS

Study area

The study was conducted at a post coal mining area in the lowlands (approximately 58-200 m above sea level) of East Kutai District, East Kalimantan Province (Figure 1). The mining area was previously a production forest, i.e., a forest concession area, loaned from the Ministry of Forestry under condition that at the end of mining activities, the area should be returned to its pre-mining condition. This study was conducted in two sites: 1) a pre-mining area, i.e., a production forest dominated by *Acacia mangium* (mainly for pulp and paper production) which has been abandoned for a period of 16 years, and 2) a reclamation site, i.e., a post mining area which has been prepared for reclamation by forming it following its original contour. The latter site was covered with 30 cm of top soil in order to plant vegetation. The distance between the two locations was approximately 3 km. At the reclamation site, about 48 tree species were planted, including fast growing species (*Acacia mangium*, *Cassia siamea*, *Paraserianthes falcataria* etc.). In the reclamation program, plant survival was monitored every three months for a period of one year. Trees showing unsuccessful growth were replaced using the same species. In order to restore plant diversity to its pre-mining condition, in the third year of revegetation, a total of 56 local species, e.g., *Dryobalanops aromatica* and *Shorea leprosula*, were planted.

Procedures

Plant diversity in both the pre-mining and reclamation sites was studied for a period of six months (from March to September 2013). In each location, vegetation sampling was conducted in 20 plots measuring 20x20 m² along line transects (Soerianegara and Indrawan 1998). The distance between plots was 100 m (Fig.2). The 20x20 m² plots were used to sample tree species within these plots, 5x5 m² subplots were used to sample non-tree woody plants and 1x1 m² subplots to sample herbaceous plants. Non-tree woody plants were later grouped together with herbaceous plants. Plant samples (both tree and herbaceous species) were collected if needed in order to identify unknown species for identification. Plant species were identified using references in Herbarium Bandungense, School of Life Sciences and Technology, Institut Teknologi Bandung, West Java, Indonesia.

Degree of vegetation cover was measured as percentage of area occupied by a plant's crown, stem (basal area) or patch. Tree basal area was determined after measuring diameter at breast height (Phillips 1959). The data collected from plots were analyzed for frequency, density and abundance (Kent and Coker 1992), their relative values were calculated as follows:

$$\text{Relative density} = \frac{\text{Number of individuals of species}}{\text{Total number of individuals}} \times 100$$

$$\text{Relative dominance} = \frac{\text{Dominance of a species}}{\text{Dominance of all species}} \times 100$$

$$\text{Relative frequency} = \frac{\text{Frequency of species}}{\text{Frequency of all individuals}} \times 100$$




Figure 1. Location of the study area in East Kalimantan at 117°12'50"-117°23'30" EL and 00°02'20"-00°13'00" NL. A. Borneo island, B. Red circle 1 is the pre-mining site and red circle 2 is reclamation site location.




Figure 2. A. Plots in line transects in pre-mining and reclamation sites, each plot measuring 20x20 m² with 100 m distance between plots; B. Nested plot measuring 20x20 m² for trees, 5x5 m² for non-tree woody plants and 1x1 m² for herbaceous plants.

Importance value for each species = relative density + relative dominance + relative frequency

Where, dominance is defined as the mean basal area per tree times the number of trees of the species.

The species diversity index (H') was determined following Shannon-Wiener (Kent and Coker 1992; Hazarika et al. 2006; Ekka and Behera 2011), i.e., as follows:

$$H' = -\sum_{i=1}^S (p_i \ln p_i)$$

Where, H' = observed species diversity, S = the number of species; p_i = the proportion of individuals or the abundance of the i^{th} species expressed as a proportion of total cover; \ln = log base _{e} .

Data analysis

All data were calculated using Excel for Windows 7 for the relative density, relative frequency, relative dominance, Importance value index and diversity index (H').

RESULTS AND DISCUSSION

Plant diversity in pre-mining area

The pre-mining area was formerly a production forest dominated by *Acacia mangium* and *Cassia siamea*. After the change of management from production forest to mining company, no activities were conducted in the area. The land was abandoned and vegetation grew naturally for 16 years without any human interference. In this area, we found 185 plant species, consisting of 133 tree species and 52 herbaceous plant species (Table 1 and 2). Among the

tree species, there were 132 local species and one non-local species (*Acacia mangium*) which was present since the area was still a production forest. The importance value indices of plant species in this area varied from 0.64% to 46.08%. Fifteen species of tree species had importance value indices higher than 20%, while 15 species of herbaceous plants had indices higher than 5% (Figure 3). Specifically, tree species were dominated by *Macaranga gigantea* (46.08 %), *Eusideroxylon zwageri* (40.96%), *Cananga odorata* (39.04 %), *Euodia speciosa* (36.48 %) and *Dillenia excelsa* (33.92 %); while the herbaceous understorey was dominated by *Blumea balsamifera* (29.04 %), *Pandanus* sp. (23.05%), *Donax cannaeformis* (19.27%), *Phrynium placenterium* (19.10 %) and *Selaginella plana* (17.76 %).

Plant diversity in reclamation site

In the first year of reclamation, 48 non-local species were planted, consisting of pioneer plants and fast growing trees. Three years later, 52 local tree species were planted. After 16 years of reclamation, we found 104 species which consist of 76 tree species and 28 herbaceous species (Table 1). The importance value indices of plant species in this area varied from 0.75% to 66.75%. The top 15 species with highest importance value indices (higher than 4.35% for tree species and more than 7.51% for the herbaceous plants) are presented in Figure 4. tree species were dominated by *Acacia mangium* as shown by the highest importance value index of 37.75%, followed by *Cassia siamea* (35.08%), *Paraserianthes falcataria* (28.28%), *Dryobalanops aromatic* (18.85%) and *Samanea saman* (15.23%). Herbaceous species were dominated by *Diplazium esculentum* with importance value index of 66.75%, followed by *Blumea balsamifera* (30.01%), *Ageratum conyzoides* (18.75%), *Acmella oleracea* (14.25%) and *Merremia peltata* (12.75%).

Table 1. Diversity of tree species at pre-mining and reclamation site (for present or × for absent)

Tree species	Pre-mining	Reclamation
<i>Acacia mangium</i> Willd.		
<i>Actinodaphne diversifolia</i> Merr.		×
<i>Actinodaphne glomerata</i> (Bl.) Nees.		
<i>Aglaia grandis</i> Korth.ex Miq.	×	
<i>Aglaia tomentosa</i> Teijsm.& Binn.	×	
<i>Aleurites moluccana</i> (L.) Willd.		
<i>Alstonia angustiloba</i> Miq.		
<i>Anthocephalus chinensis</i> (Roxb) Bosser.		
<i>Aquilaria malaccensis</i> Lam.		
<i>Archidendron havilandii</i> (Ridl.) I.C.Nielsen.		
<i>Artocarpus altilis</i> (Parkinson) Fosberg.		
<i>Artocarpus champeden</i> (Lour.) Stokes.		
<i>Artocarpus rigidus</i> Blume, Bijdr.		
<i>Averrhoa carambola</i> L.		
<i>Baccaurea macrocarpa</i> (Miq.) Müll.Arg.	×	
<i>Baccaurea</i> sp.		
<i>Baccaurea stipulata</i> J.J.Sm.	×	
<i>Barringtonia sarcostachys</i> (Blume) Miq.	×	
<i>Bauhinia</i> sp.		
<i>Beilschmiedia rivularis</i> Kosterm.	×	
<i>Bischofia javanica</i> Blume.		
<i>Callicarpa pentandra</i> Roxb.		
<i>Cananga odorata</i> (Lam.) Hook.f. & Thomson		
<i>Canarium odontophyllum</i> Miq.	×	
<i>Canthium confertum</i> (Burm.f.) Alston	×	
<i>Casuarina equisetifolia</i> L.		
<i>Cleistanthus myrianthus</i> (Hassk.) Kurz.		
<i>Clerodendrum confusum</i> Hallier f.	×	
<i>Cratoxylum arborescens</i> (Vahl) Blume.		
<i>Croton argyratus</i> Blume, Bijdr.	×	
<i>Cryptocarya</i> sp	×	
<i>Cyathea contaminans</i> (Wall. ex Hook.) Copel	×	
<i>Dacryodes rostrata</i> (Blume) H.J.Lam	×	
<i>Dehaasia incrassata</i> (Jack) Nees.	×	
<i>Dillenia excelsa</i> (Jack) Martelli ex Gilg.	×	
<i>Dillenia reticulata</i> King.	×	
<i>Dillenia sumatrana</i> Miq.	×	
<i>Dimocarpus longan</i> Lour.	×	
<i>Diospyros borneensis</i> Hiern.	×	
<i>Diospyros macrophylla</i> Blume.	×	
<i>Diospyros</i> sp.	×	
<i>Dipterocarpus confertus</i> Slooten.	×	
<i>Dipterocarpus cornutus</i> Dyer.	×	
<i>Dipterocarpus humeratus</i> Slooten	×	
<i>Disepalum anomalum</i> Hook.f.	×	
<i>Dracontomelon dao</i> (Blanco) Merr. & Rolfe.		
<i>Drimycarpus luridus</i> (Hook. f.) Ding Hou.	×	
<i>Dryobalanops aromatica</i> Gaertn.f., <i>nom cons.</i>		
<i>Drypetes longifolia</i> (Blume) Pax & K. Hoffm.	×	
<i>Drypetes subcubica</i> (J.J.Sm.) Pax & K.Hoffm.	×	
<i>Duabanga moluccana</i> Blume.		
<i>Durio zibethinus</i> Rumph. ex Murray.		
<i>Dyera costulosa</i> (Miq.) Hook. film.	×	
<i>Euodia alba</i> Hook. f.	×	
<i>Euodia speciosa</i> Rchb.f. & Zoll. ex Teijsm. & Binn.		×
<i>Elmerrillia tsiampacca</i> (L.) Dandy.		×
<i>Eusideroxylon zwageri</i> Teijsm. & Binn.		
<i>Evodia latifolia</i> DC.		
<i>Ficus fistulosa</i> Reinw. ex Blume		×
<i>Ficus geocarpa</i> Teijsm. ex Miq.		×
<i>Ficus</i> sp.		
<i>Ficus uncinata</i> (King) Becc.,		×
<i>Ficus variegata</i> Blume		×
<i>Flacourtia rukam</i> Zoll. & Moritzi		×
<i>Glochidion</i> sp.		
<i>Gluta renghas</i> L.		
<i>Hibiscus similis</i> Bl.		
<i>Homalanthus populneus</i> (Giesel.) Pax.		×
<i>Horsfieldia grandis</i> (Hk. f.) Warb.		
<i>Ilex cymosa</i> Blume		
<i>Knema conferta</i> (King) Warb.		×
<i>Knema latericia</i> Elm.		×
<i>Koompassia malaccensis</i> Maingay ex Benth.		
<i>Koordersiodendron pinnatum</i> (Blanco) Merr. Bull.		
<i>Lansium Domisticum</i> Corr.		
<i>Lansium parasiticum</i> (Osbeck) Sahni et. Bennet		
<i>Leea aculeata</i> Blume		
<i>Lepisanthes alata</i> (Blume) Leenh.		
<i>Leucaena glauca</i> (Linn.) Benth.		
<i>Lithocarpus</i> sp.		×
<i>Litsea accendens</i> (Blume) Boerl.		
<i>Litsea</i> sp.		
<i>Macaranga gigantea</i> (Reichb.f. & Zoll.) Muell.		
<i>Macaranga hypoleuca</i> (Rchb.f. & Zoll.) Müll.Arg.		×
<i>Macaranga pruinosa</i> (Miq.) Mull Arg		×
<i>Macaranga</i> sp.		×
<i>Macaranga tanarius</i> (L.) Müll.Arg.		×
<i>Macaranga triloba</i> Thunb.) Müll.Arg.		
<i>Maranthes corymbosa</i> Blume.		×
<i>Merremia peltata</i> (L.) Merr.		×
<i>Memecylon garcinoides</i> Blume.		×
<i>Mitrephora fragrans</i> Merr.		×
<i>Myristica elliptica</i> Wall.ex. Hook. f. Thoms.		×
<i>Myristica guatterifolia</i> A.DC.		×
<i>Myristica maxima</i> Warb.		
<i>Nauclea calycina</i> Bartl. ex DC		×
<i>Nauclea purpurascens</i> Korth.		×
<i>Neouvaria acuminatissima</i> (Miq.) Airy Shaw		×
<i>Nephelium cuspidatum</i> Blume.		×
<i>Nephelium lappaceum</i> L.		×
<i>Nypa fruticans</i> Wurmb		×
<i>Ochanostachys amentacea</i> Mast.		×
<i>Octomeles sumatrana</i> Miq.		×
<i>Homalanthus</i> sp.		
<i>Orophea corymbosa</i> (Blume) Miq		×
<i>Palaquium quercifolium</i> (de Vriese) Burck		
<i>Planchonia valida</i> (Blume) Blume		×
<i>Pholidocarpus</i> sp.		×
<i>Polyalthia obliqua</i> Hook.		×

<i>Polyalthia sumatrana</i> (Miq.) Kurz.	×
<i>Pternandra rostrata</i> (Cogn.) M.P.Nayar.	×
<i>Pterospermum diversifolium</i> Blume.	×
<i>Pterospermum javanicum</i> Jungh.	×
<i>Sageraea lanceolata</i> Miq.	×
<i>Sageraea glabra</i> Merr.	×
<i>Saraca hullettii</i> Prain.	×
<i>Saurauia nudiflora</i> DC.	×
<i>Schima wallichii</i> (DC.) Korth.	×
<i>Semecarpus glaucus</i> Engl.	×
<i>Shorea dispar</i> Ashton.	
<i>Shorea laevis</i> Ridl.	
<i>Shorea leprosula</i> Miq.	×
<i>Shorea pinanga</i> Scheff.	
<i>Shorea seminis</i> (de Viese) v.Slooten	×
<i>Syzygium acuminatissima</i> (Blume) Merr. & Perry.	
<i>Syzygium</i> sp. 1	×
<i>Tetramerista glabra</i> Miq.	
<i>Trema orientalis</i> (L.) Blume	×
<i>Vatica</i> sp.	×
<i>Vernonia arborea</i> Buch.-Ham.	×
<i>Vitex pubescens</i> Vahl.	
<i>Walsura pinnata</i> Hassk.	×
<i>Xanthophyllum vitellinum</i> (Blume) D.Dietr.	×

Comparison between pre-mining and reclamation sites

After 16 years of reclamation, plant diversity in the reclamation site was still lower than in the pre-mining site which was used as reference. Comparison of species diversity indices (H') between the two sites indicate a significant different in tree species, however, diversity indices for herbaceous plants were similar between the two sites. The diversity index for tree species in pre-mining area was 4.29 and in reclamation site was 3.54. Table 3 presents species richness and diversity indices in the two sites.

It is interesting to note that 16 species of tree species plants in the reclamation site grew spontaneously without being planted by the reclamation program. These were *Leucaena glauca*, *Lansium domesticum*, *Shorea laevis*, *Homalanthus populneus*, *Durio zibethinus*, *Casuarina equisetifolia*, *Averrhoa carambola*, *Artocarpus champeden*, *Bauhinia* sp, *Palaquium quercifolium*, *Horsfieldia grandis*, *Evodia latifolia*, *Cleistanthus myrianthus*, *Bischofia javanica*, *Koompassia malaccensis* and *Actinodaphne glomerata* (Figure 5). Considering the seed form and dispersal process, it is assumed that the presence of these species in the reclamation site is related to the activity of animals which are found near the reclamation site and act as dispersal agents. Both the pre-mining and reclamation sites are in fact not too far from a forest system outside the area. This forest system could be considered as a source of plant diversity, so plant colonization at the reclamation site was presumably assisted by animals from its surroundings. This phenomenon may have benefited the reclamation program, although actual plant dispersal processes need to be confirmed.

Table 3. Diversity of herbaceous species at pre-mining and reclamation site (for present or × for absent)

Herbaceous	Pre-mining	Reclamation
<i>Acmella oleracea</i> (L.) R.K.Jansen,		
<i>Ageratum conyzoides</i> L.		
<i>Amomum maximum</i> Roxb		
<i>Asplenium</i> sp.		×
<i>Begonia</i> sp.		×
<i>Blumea balsamifera</i> L.		
<i>Boehmeria nivea</i> (L.) Gaudich.		
<i>Borreria</i> sp.		×
<i>Calathea bachemiana</i> E.Morren.		×
<i>Callicarpa longifolia</i> Lamarck.		
<i>Colocasia esculenta</i> (L.) Schott;		
<i>Colocasia</i> sp.		×
<i>Commelina</i> sp.		×
<i>Curculigo latifolia</i> Dryand.		
<i>Cyperus</i> sp.		
<i>Diplazium esculentum</i> (Retz.) Sw.		
<i>Donax cannaeformis</i> (G.Forst.) K.Sch.		
<i>Dryopteris</i> sp.		×
<i>Fimbristylis</i> sp.		×
<i>Gleichenia linearis</i> (Burm. f.) S.W. Clarke.		×
<i>Globba leucantha</i> Miq.		×
<i>Helminthostachys zeylanica</i> (L.) Hook.		
<i>Hornstedtia affinis</i> Riedl.		×
<i>Hornstedtia irya</i> (Gaertn.) Warb.		×
<i>Labisia pumila</i> Benth. & Hook		
<i>Lantana camara</i> L		
<i>Mapania</i> sp		×
<i>Melastoma malabathricum</i> L.		
<i>Merremia peltata</i> (L.) Merr.		
<i>Mimosa pudica</i> L.		
<i>Pandanus</i> sp.		×
<i>Pandanus tectorius</i> Parkinson.		
<i>Peperomia pellucida</i> (L.) Kunth.		
<i>Phacelophrynium maximum</i> (Blume) K. Schum		
<i>Phyllanthus urinaria</i> L.		×
<i>Phrynium hirtum</i> Riedl.		
<i>Phrynium placentarium</i> (Lour.) Merr.		
<i>Phrynium</i> sp.		×
<i>Piper betle</i> L.		
<i>Piper nigrum</i> L.		
<i>Plagiostachys albiflora</i> Riedl.		×
<i>Pleomele angustifolia</i> (Roxb.) N.E.Br.		×
<i>Pteris</i> sp.		×
<i>Rhaphidophora minor</i> Hook.f. Climber.		×
<i>Saccharum spontaneum</i> L.		
<i>Scoparia dulcis</i> L.		
<i>Selaginella plana</i> Hieron.		
<i>Stenochlaena palustris</i> (Burm.f.) Bedd.		
<i>Tectaria</i> sp.		×
<i>Tetrastigma</i> sp.		×
<i>Zingiber longipedunculatum</i> Riedl.		
<i>Zingiber</i> sp.		×

Table 3. Plant species diversity indices in pre-mining and reclamation site

Site	Pre-mining		Reclamation	
	Number	H'	Number	H'
Tree species	133	4.29	76	3.54
Herbaceous plants	52	2.67	28	2.97
Total species	185		104	




Figure 3. The importance value indices (IVI) of 15 species of tree species (a) and 15 species of herbaceous plants (b) in pre-mining area




Figure 4. The importance value indices (IVI) of 15 species of tree species plants (a) and 15 species of herbaceous plants (b) in reclamation area




Figure 5. Importance value indices (IVI) of 16 tree species that grew spontaneously at reclamation site

Discussion

The mahang tree (*Macaranga gigantea*) is a local species which dominated in the pre-mining area, as shown

by its highest importance value index. *Macaranga* is known as a pioneer and fast growing plant, with soft wood, which sprouts all year round and is able to reach a height of up to 20 meters (Zakaria et al. 2008). In addition, many species of *Macaranga* sp. favor high light intensity, indicating its relative tolerance to open areas. Several species of *Macaranga*, such as *Macaranga tanarius* and *Macaranga javanica* have been used as indicator species for disturbed forest areas (Zakaria et al. 2008). In the reclamation site, the dominant species was *Acacia mangium*. However, *Macaranga* sp. was also quite abundant although not many seedlings were planted (Komara et al., unpublished data). The high number of seedlings of local species planted in the reclamation site was one of the factors contributing to the importance value indices of these plant species. In the case of *Macaranga* sp., however, abundance may also have been supported by the natural availability of seedlings in the site. This phenomenon is in accordance with the generalization of major factors affecting succession, as stated by Noble and Slatyer (1980), i.e. that the composition of species immediately after a disturbance depends on propagules

which have either dispersed from elsewhere or have persisted through the disturbance or on vegetative resprouting of organs surviving the disturbance. Three groups of vital attributes are important to vegetation replacement, i.e., the method of arrival or persistence of a species during and after disturbance; the ability to establish and grow to maturity; and the time taken for them to reach critical stages in their life history (Noble and Slatyer 1980).

The same phenomenon was found in *Eusideroxylon zwageri*, known as the ulin tree. This local species was the second dominant species in the pre-mining site. This plant is well known for its high quality wood but slow growth rate. The high importance value index of ulin in this area indicates that this area could conserve the ulin tree in its natural habitat. In the reclamation site, *Eusideroxylon zwageri* was the sixth rank. The slow growth rate of this species seems to have contributed to its relatively low importance value in the reclamation site.

In the pre-mining site, the dominant herbaceous species was *Blumea balsamifera* with the highest importance value index (29.04%), while in the reclamation site this plant was the second dominant species with an importance value index of 30.01%. The dominant species in the reclamation site was the fern *Diplazium esculentum* with an importance value index of 66.75%. *Blumea balsamifera* has abundant leaves, grows relatively fast, and is extremely tolerant to minimum light intensity (shade plant). In contrast, *Diplazium esculentum* is tolerant to high light intensity. The presence of *Diplazium esculentum* could be related to its dispersal ability, i.e., dispersed by wind from the forest system. However, this needs to be reconfirmed.

The presence of herbaceous plants benefits the reclamation program through their role as ground cover. Plants such as *Diplazium esculentum*, *Blumea balsamifera*, *Ageratum conyzoides*, *Acmella oleracea* and *Merremia peltata* easily grow in relatively poor soil conditions under sufficient sunlight. Besides its rapid growth, *Diplazium esculentum* seems to be an adaptable species for post-mining areas. Several herbaceous species from the pre-mining area were found in the reclamation site, e.g., *Piper betel*, *Selaginella plana*, *Selaginella deoderleinii* and *Piper nigrum*.

The ability of plants to grow in the reclamation site can be attributed to seeds already present in the top soil during reclamation process, or by seed dispersal. For example, *Leucaena glauca* (5.08 %) is propagated by seeds, and is easy to grow after being cut, felled or burned. *Lansium domesticum* (4.35%), *Durio zibethinus* (4.35%), *Averrhoa carambola* (1.45%) and others are several fruits species with seeds that could be dispersed by animals. *Shorea laevis* (4.35%) seeds contain a lot of fat that is commonly eaten by animals. Dispersion of plant species depends on animal species and the distance between the reclamation site and the nearby vegetation source, e.g., the nearest forest that will affect the distribution of the species (Novianti 2013; Traveset et al. 2014).

Based on the number of plant species that successfully grew at the reclamation site, the following findings could be used to suggest further management of this study site: in terms of species richness, there were 132 local species in

the pre-mining area as reference site, while at the reclamation site there were 35 planted local species and 16 local species which grew spontaneously. In other words, a total of 51 local species successfully grew at the reclamation site after a period of 16 years. To restore plant diversity in the reclamation site to its assumed pre-mining conditions, it should be planted with 81 local species of tree species. The next step is to select the local tree species from the pre-mining area by considering the species importance value indices. In the pre-mining area, only eight local species had importance value indices higher than 20%, e.g., *Euodia speciosa* (36.48%), *Shorea pinanga* (31.80%), *Lithocarpus* sp. (25.44%), *Ficus variegata* (25.01%), *Ficus uncinata* (22.72%), *Pternandra rostrata* (22.40%), *Shorea dispar* (21.76%) and *Canarium odontophyllum* (21.76%); however the reclamation program must consider the availability of seedlings and the ability of the plant to grow successfully in reclamation conditions.

To conclude, after 16 years of reclamation, 104 plant species were found in the reclamation site, consisting of 76 tree species and 28 herbaceous species. tree species that successfully grew in this site consisted of 35 planted local species (e.g., *Dryobalanops aromatica*, *Eusideroxylon zwageri*, *Macaranga gigantea*), 25 planted non local species, and 16 local species that grew spontaneously (e.g., *Leucaena glauca*, *Lansium domesticum*, *Shorea laevis*). In comparison, data from the pre-mining area indicate the presence of 133 plant species, consisting of 132 local tree species, one non local tree species (*Acacia mangium*) and 52 herbaceous species. Tree species diversity index in the reclamation site after 16 years post mining (i.e., 3.54) was still lower than in the pre-mining area (4.29); while the diversity indices for herbaceous plants were relatively similar (2.97 and 2.67 in the reclamation and pre-mining sites respectively). The slightly higher diversity index in the reclamation site can be attributed to higher coverage per species in this site, despite lower species richness.

ACKNOWLEDGMENTS

This research was funded by the Directorate General for Higher Education of Republic Indonesia granted to the first author and Operational Budget for Higher Education through the School of Life Sciences and Technology, Institut Teknologi Bandung, West Java, Indonesia. Many thanks to PT. Indominco Mandiri for all the facilities during this study.

REFERENCES

- Alday JG, Marrs RH, Ruiz CM. 2012. Soil and vegetation development during early succession on restored coal wastes: a six-year permanent plot study. *Plant Soil*. 353(1-2):305-320
- Claassen S, Rensburg, PJJV, Maboeta MS, Rensburg LV. 2008. Soil microbial community function and structure in a post-mining chronosequence. *Water Air Soil Pollution* 194: 315-329.
- Cooke JA, Johnson MS. 2002. Ecological restoration of land with particular reference to the mining of metals and industrial minerals: A review of theory and practice. *Environ Rev* 10: 41-71.

- Dale VH, Beyeler SC. 2001. Challenges in the development and use of ecological indicators. *Ecol Indic* 1: 3-10.
- Ekka NJ, Behera N. 2011. Species composition and diversity of vegetation developing on an age series of coal mine spoil in an open cast coal field in Orissa, India. *Trop Ecol* 52 (3): 337-343.
- Graham, MH, Hayners RJ. 2004. Organic Matter Status and the size activity and metabolic diversity of the soil microflora as indicator of the success rehabilitation of mined sand dunes. *Biologi Fertil Soils* 30: 429-437.
- Hazarika P, Talukdar NC, Singh YP. 2006. Natural colonization of plant species on coal mine spoils at Tikak Colliery, Assam. *International Soc Trop Ecol* 47 (1): 37-46.
- Hutchinson TF, Boerner REJ, Iverson LR, Sutherland S, Sutherland EK. 1999. Landscape patterns of understory composition and richness across a cross a moisture and nitrogen mineralization gradient in Ohio (USA.) *Quercus* forests. *Plant Ecol* 144: 177-189.
- Kent M, Coker P. 1992. *Vegetation Description and Analysis. A practical approach.* John Wiley and Sons, New York.
- Ludwig JA, Hindley N, Barnett G. 2003. Indicators for monitoring mine site rehabilitation: trends on waste-rock dumps, northern Australia. *Ecol Indic* 3: 143-153.
- Makineci E, Gungor BS, Kumbasli M. 2011. Natural plant revegetation on reclaimed coal mine landscapes in Agacli-Istanbul. *African J Biotechnol* 10 (16): 3248-3259.
- Malakar S, Gupta H, Kumar ML. 2015. Species composition and some psycho-chemical properties of an age series of overburden dumps in Raniganj Coalfields, West Bengal, India. *Intl J Sci Res Environ Sci* 3 (7): 0239-0247.
- Mansur I. 2010. *Silviculture technic for post mining area reclamation.* Seameo Biotrop. Bogor. Indonesian.
- Noble IR, Slatyer RO. 1980. The use of vital attributes to predict successional changes in plant communities subject to recurrent disturbances. *Plant Ecol* 43 (1): 5-21.
- Novianti V. 2013. *Process of Primary Succession and on Previously Mined Coal Mine Areas.* [Dissertation]. Institut Teknologi Bandung, Bandung. [Indonesian]
- Nugroho AW, Adman B. 2011. Growth of local plant species on land reclamation in Tenggara Seberang, East Kalimantan. In: *Proceedings of the Research Results Seminar of BPTKSDA; Balikpapan, November 3, 2011.* pp 211-217.
- Perrow MR, Davy AJ. 2002. *Handbook of Ecological Restoration, Vol. 1. Principles of Restoration.* Cambridge University Press, Cambridge, UK.
- Phillips BA. 1959. *Methods of Vegetation Survey.* Henry Holt and Co. Inc., New York.
- Regulation of State Minister for The Environment no. 4/2012 about Environment Friendly Indicators for Business and/or Activity of Open Coal Mining. [Indonesian]
- Sheoran V, Sheoran AS, Poonia P. 2010. Soil Reclamation of Abandoned Mine Land by Revegetation: A Review. *Intl J Soil Sediment Watter* 13 (2): 13.
- Sheoran V, Sheoran AS. 2009. Reclamation of abandoned mine Land. *J Mining Metallurg* 45 (1): 13-32.
- Shrestha RK, Lal R. 2011. Changes in physical and chemical properties of soil after surface mining and reclamation. *Geoderma* 11: 268-276
- Soerianegara I, Indrawan A. 1998. *Forest Ecology of Indonesia.* Forestry Faculty. Institut Pertanian Bogor, Bogor.
- Traveset A, Heleno R, Nogales M. 2014. *Seeds: The Ecology of Regeneration in Plant Communities.* 3rd ed. CAB International, London.
- Turner IM. 2004. *The Ecology of Trees in the Tropical Rain Forest.* Cambridge University, UK.
- Wardana W. 2008. *Evaluation of soil potention and vegetation biomass in reclamation area of PT. Kaltim Prima Coal. Sengata. East Kutai.* [Thesis]. Universitas Mulawarman, Samarinda. [Indonesia]
- WWF. 2005. *Borneo: Treasure Island at Risk. Status of Forest, Wildlife and related Threats on the Island of Borneo.* WWF Germany, Frankfurt.
- Zakaria R, Rosely NFN, Mansor M, Zakaria Y. 2008. The distribution of *Macaranga* Genus (Family Euphorbiaceae) In Penang island, Peninsular Malaysia. *J Biosci* 19 (2): 91-99.