

# A mutant variant of *Nix* gene on $\gamma$ -irradiated *Aedes aegypti* as a male-determination factor

BENI ERNAWAN<sup>1</sup>, HARRY NUGROHO EKO SURNIYANTORO<sup>2,\*</sup>, IRAWAN SUGORO<sup>1</sup>,  
USMAN SUMO FRIEND TAMBUNAN<sup>3</sup>

<sup>1</sup>Center for Isotopes and Radiation Application, National Nuclear Energy Agency of Indonesia. Jl. Lebak Bulus Raya 49, Jakarta 12070, Indonesia

<sup>2</sup>Center for Technology of Safety and Radiation Metrology, National Nuclear Energy Agency of Indonesia. Jl. Lebak Bulus Raya 49, Jakarta 12070, Indonesia. Tel.: +62-21-7513906, Fax.: +62-21-7657950, \*email: harry\_nes@batan.go.id

<sup>3</sup>Department of Chemistry, Faculty of Mathematics and Sciences, Universitas Indonesia. Jl. Lingkar UI, Depok 16424, West Java, Indonesia

Manuscript received: 17 January 2019. Revision accepted: 27 February 2019.

**Abstract.** *Ernawan B, Surniyantoro HNE, Sugoro I, Tambunan USF. 2019. A mutant variant of Nix gene on  $\gamma$ -irradiated Aedes aegypti as a male-determination factor. Biodiversitas 20: 893-899.* This study was aimed to examine the effect of  $\gamma$ -irradiation on the *Nix* gene sequence as a male-determination factor on *Ae. aegypti*. The novelty of the present study is the investigation on the gene mutation level of the deoxyribonucleic acid as an impact of ionizing radiation exposure. Males *Ae. aegypti* at the pupal stage were sterilized by applying 70 Gy  $\gamma$ -rays in varies dose-rates, i.e., 300, 600, 900, 1200 and 1500 Gy/h utilizing panoramic irradiator and males group without  $\gamma$ -irradiation as controls. Adult males that emerged from the pupal stage were assessed for the *Nix* gene sequence using a Real-Time Polymerase Chain Reaction method and were sequenced using Clustal Omega and Bioedit software, respectively. The results indicated that mutation occurred in *Nix* gene sequences were categorized as a point mutation. The change of *Nix* gene sequences was occurred between the nitrogenous base of guanine to adenine (G>A), thymine to adenine (T>A) and cytosine to adenine (C>A) in certain sequence numbers. There were six and nineteen on the forward-strand and reverse-strand of the *Nix* gene, respectively, due to  $\gamma$ -irradiation treatment. We conclude that  $\gamma$ -irradiation exposure causes point mutation on the *Nix* gene of males *Ae. aegypti*.

**Keywords:** *Ae. aegypti*, *Nix* gene, point mutation, Real-Time Polymerase Chain Reaction,  $\gamma$ -irradiation

**Abbreviations:** DNA: deoxyribonucleic acid; WHO: World Health Organization; DDT: Dichloro-diphenyl-trichloroethane; SIT: Sterile insect technique; bp: base pairs; Co-60: Cobalt 60 radioisotope; RT-PCR: Real-time Polymerase chain reaction; IR: Ionizing radiation; Gy: Gray (Unit of ionizing radiation dose); FISH: Fluorescence in situ hybridization;  $\gamma$ : Gamma

## INTRODUCTION

*Aedes aegypti* is a very popular mosquito species because of its role as a vector of several viral diseases in tropical countries, including dengue fever, chikungunya, yellow fever and Zika (Vorou 2016). Historically, *Ae. aegypti* originated from Africa, then spread to various continents, especially in tropical areas, including Central America, Central Asia to Southeast Asia and Northern Australia (Jansen and Beebe 2010). Distribution of *Ae. aegypti* to various areas is influenced by various factors, including the species ability to adapt in different conditions or climates, geographical conditions, mobility and population migration (Harrington et al. 2005; Chadee and Martinez 2016).

The incidence rate of dengue fever has grown rapidly around the world in recent decades. Over 2.5 billion people (40% of the world's population) are now at risk from dengue fever. The most significant dengue epidemics region in recent years have occurred in the Americas, Western Pacific, Africa, and Southeast Asia, with approximately 390 million dengue infections occur around the world per year. Of these, 500,000 cases develop into dengue hemorrhagic fever, a more severe form of the disease, which results in up to 25,000 deaths annually worldwide (WHO 2011). The Indonesian Ministry of

Health noted that the number of dengue fever in Indonesia from January to February 2016 was 8,487 patients with 108 deaths. The highest number at the ages of 5-14 years reached 43.44% and 15-44 years reached 33.25% (Kemenkes RI 2016).

So far, the control of the *Ae. aegypti* mosquito has been focused on using pesticides such as DDT, temephos, malathion, and, more recently, pyrethroids (Maestre-Serrano et al. 2014). Unfortunately, most pesticides were failed and resulted in the development of mosquito resistance to insecticides. Metabolic resistance involved a large family of multi-gene enzymes: cytochrome P450s, glutathione S-transferases, and carboxylesterase. P450 cytochrome, in particular, has been involved in providing resistance to mosquitoes (Ngoagouni et al. 2016; Moyes et al. 2017). This encourages scientists to find new methods to control the spread of *Ae. aegypti*, one of which is by utilizing nuclear techniques for sterile insect technique (SIT). Until now, information related to the effect of gamma irradiation dose rates on the quality parameters of sterile *Ae. aegypti* mosquitoes in SIT is not sufficient. The information is needed to optimize sterilization process using gamma irradiation as one of the main components of SIT.

Utilization of nuclear techniques has been widely applied in various fields. Besides being used in sterile


insect techniques (Dyck et al. 2005; Ernawan et al. 2017; Ernawan et al. 2018), nuclear techniques can also be used in attenuating rodent malaria parasites of *Plasmodium berghei* in blood of mice (Syaifudin et al. 2013) and inhibition of *Plasmodium falciparum* multiplication supplemented with tritium-labeled hypoxanthine (Surniyantoro et al. 2016). *Ae. aegypti* has 1,376 Mbp of a genome and three chromosomes (Nene et al. 2007). The *Nix* gene is one of the target genes that control the masculinity of *Ae. aegypti*. This gene is located at the M locus on chromosome 1 (denoted 1.q.2.1), meaning it is located on chromosome 1, arm q, region 2 and band 1 as shown in Figure 1 (Hall et al. 2015; Fontaine et al. 2017). Gene mutation in the *Nix* gene can cause damage to the structure and function of the *Nix* protein. This study was aimed to examine the effect of gamma irradiation on the *Nix* gene sequence as a male-determination factor on *Ae. aegypti*.

## MATERIALS AND METHODS

### Procedures

#### *Aedes aegypti* strain rearing

*Aedes aegypti* used in this study was collected from South Tangerang, Banten Province, Indonesia in 2014 and then maintained in the Entomology Laboratory, Center for Isotopes and Radiation Application, National Nuclear Energy Agency of Indonesia, Jakarta. The mosquito colonies used in the study were the fourteenth generation from the field collection. The mosquito strain was maintained in a laboratory with controlled climate parameters, including temperatures of  $26\pm 1^\circ\text{C}$ , relative humidity  $70\pm 10\%$  and photoperiod (12: 12 hours). Adult mosquitoes were maintained in the Bugdorm-1® mosquito cage (30x30x30 cm) and supplied with 10% (w/v) sugar solution, while female mosquitoes are supplied blood feed every week for egg maturation.



**Figure 1.** The position of *Nix* gene on M locus of *Aedes aegypti*'s chromosome 1; A. A representation of the relative position of the transgenes with respect to the M-locus and the m-locus; B. A karyotype of mitotic chromosomes from the khw strain of *Aedes aegypti* in males using Fluorescence in situ hybridization (FISH) technique (Hall et al. 2014)

#### Gamma irradiation procedure

Male pupa aged less than 24 hours were inserted into transparent plastic tubes and then were irradiated with a 70 Gy gamma irradiation in varies dose-rate, which are 300, 600, 900, 1200 and 1500 Gy/h (interval 300 Gy/h). The gamma irradiation process was carried out using a panoramic irradiator (Babha Atomic Research Center, India) at the Center for Isotopes and Radiation Application, National Nuclear Energy Agency of Indonesia, Jakarta. Radioisotope used as sources of gamma irradiation was Co-60. Control pupa was treated the same as experimental treatments, but without gamma irradiation. After the irradiation process, a plastic vial tube containing pupa is filled with water and placed into Bugdorm-1® until the pupa develops into an adult mosquito for further analysis.

#### Genomic deoxyribonucleic acid (DNA) isolation

The genomic DNA was isolated from 20 mg of males *Ae. aegypti* in each treatment using the GeneJET Genomic DNA Purification kit (ThermoFisher Scientific, Cat. No. 0721) and according to the manufacturer's instructions. The obtained DNA was stored at  $-20^\circ\text{C}$ .

#### Amplification of *Nix* gene and sequence analysis

Amplification of *Nix* gene was performed using Real-Time Polymerase Chain Reaction (RT-PCR) technique, as described previously by Hall et al. (2015) with the forward primer was 5'-GATTTTTGTTTTTGTCGTGCAA-3' and the reverse primer was 5'-AATGCAAGTATCATAGGTCAGCA-3'. The PCR reactions were carried out with an enzyme activation of  $95^\circ\text{C}$  for 3 minutes, followed by denaturation of  $95^\circ\text{C}$  for 1-3 seconds and annealing of  $60^\circ\text{C}$  for 20 seconds. The amplified products were sequenced in the Integrated DNA Technologies Pte. Ltd., Singapore. The sequencing results were analyzed for *Nix* gene mutation using Clustal Omega and Bioedit software.

## RESULTS AND DISCUSSION

In this study, the effect of gamma irradiation on the *Nix* gene sequence as a male-determination factor on *Ae. aegypti* was investigated. Gamma irradiation exposure in *Ae. aegypti* males pupa were carried out using a panoramic irradiator with Co-60 as a source of gamma irradiation. There was a 70 Gy exposure at various dose-rate treatments, 0 (control), 300, 600, 900, 1200 and 1500 Gy/h. The range and time of exposure-setting in each treatment are shown in Table 1, while the result of *Nix* gene amplification is shown in Figure 2.

Figure 3 demonstrates the sequencing results of *Nix* gene in each of treatment and control. They are divided into the forward sequence (3A) and reverse sequence (3B). The results indicated that mutation occurred in *Nix* gene sequences were categorized as a point mutation. The change of *Nix* gene sequences was occurred between the nitrogenous base of guanine to adenine (G>A), thymine to adenine (T>A) and cytosine to adenine (C>A) in certain sequence numbers. There were six points mutation on the



Confirmation of the DNA amplification results by real-time PCR was carried out using 0.8% agarose gel electrophoresis. The results of the visualization of agarose gel are shown in Figure 2A. Number 1-6 is a sample of males *Ae. aegypti*'s *Nix* gene at a dose-rate of 0 Gy/h (control) up to 1500 Gy/h, respectively. Number 7 is an un-template control (without DNA template), while the last is a DNA ladder with an interval of 100 bp. Agarose gel electrophoresis results showed that the *Nix* gene DNA fragments in all gamma irradiation treatments were approximately 320 bp. It also indicated that a total length of the *Nix* gene in *Ae. aegypti* is 320 bp. Figure 2B demonstrates that the *Nix* gene in all of the gamma irradiation treatments amplified at 80°C. Controls are shown in a dark green graph. The treatments of gamma irradiation dose-rates of 300, 600, 900, 1200 and 1500 Gy/h, respectively, are shown in the light green, dark purple, light purple, blue and orange.

The amplification results were sequenced to determine the sequence of nucleotide bases in the *Ae. aegypti*'s *Nix* gene. The DNA sequence in each gamma irradiation treatment is aligned to analyze the mutations that occur. The *Nix* gene DNA sequences are aligned using the Clustal Omega software from the European Molecular Biology Laboratory - The European Bioinformatics Institute (EMBL-EBI) and BioEdit software. Clustal Omega software has some advantages, including high accuracy, fast and can be used for large scale up to 190,000 sequences using one processor (Sievers et al. 2011).

## Discussion

The strength and novelty of the present study is the investigation on the *Nix* gene mutation level of the deoxyribonucleic acid as an impact of ionizing radiation exposure. To our best knowledge, this research has not been done in Indonesia. This study can be used to develop the sterile insect technique (SIT) and to control the spread of *Ae. aegypti* by utilizing nuclear techniques. Until now, the development of SIT applications for controlling insect populations, especially mosquitoes, has undergone many developments. The method of mass mosquito breeding in the laboratory, the method of releasing mosquitoes to the target area and monitoring methods have been validated and standardized (Balestrino et al. 2014; Hapairai et al. 2014; Bourtzis et al. 2016; Zheng et al. 2015). However, research to evaluate the quality parameters of sterile mosquitoes and applications to target areas is still being carried out by researchers for the success of the program mosquito population control as a vector of disease. It can also be used as a pilot study to determine various kinds of genetic mutation information on *Ae. aegypti*'s genes. This study is the first experiment using Co-60 as a source of gamma irradiation exposure to make a genetic mutation in the *Nix* gene of *Ae. aegypti*'s DNA fragment. The selection gamma irradiation dose refers to the previous studies by Sasmita et al. (2014) that stated the optimal dose of gamma irradiation in *Ae. aegypti* was 70 Gy. This approach offers an opportunity to complete a preventing program of *Ae. aegypti* spread in the population and reducing the incidence of dengue fever in human.

As the disease vector, *Ae. aegypti*, viruses are also reported to spread in tropical countries. Dengue, a common viral disease transmitted by *Ae. aegypti* has infected 96 million people globally in 2010 (Bhatt et al. 2013). Asia contributes the highest number of infected (70%), followed by Africa (16%), America (14%) and Oceania (<0.2%). Chikungunya, first isolated in Tanzania, Africa in 1953. The virus then spread to Central and Southeast Asia (Weaver and Reisen 2010). In 2007, clinical cases of chikungunya were reported to be detected in southern Europe, namely in Italy (Rezza et al. 2007). It was also reported to be detected in the Caribbean Islands, South America in 2013-2014 (Van Bortel et al. 2014). Yellow fever, according to history as evidenced by phylogenetic studies originating in Africa and spread from the 16<sup>th</sup> century during the slave trade. Until the 20<sup>th</sup> century, yellow fever epidemics were reported in North and South America (Gubler 2004). The Zika virus was first isolated in 1947 in the Zika forest, Uganda, Africa (Li et al. 2012). Furthermore, the Zika virus spread to Asia, America, the Mediterranean, and Micronesia. In recent years, clinical cases of Zika virus have been reported in South America and are of concern to the world (Vorou 2016; Wiwanitkit 2016).

The results of our study showed that mutation in the *Nix* gene had occurred due to gamma irradiation treatment at some points. There were six point mutations on the forward-strand, base number 7 (G>A), 8 (G>A), 12 (T>A), 16 (A>T), 311 (A>T) and 312 (T>A) and nineteen on the reverse-strand, base number 5 (A>G), 6 (T>A), 7 (T>A), 9 (T>G), 10 (G>A), 13 (A>G), 14 (deletion), 15 (G>C), 16 (A>C), 18 (C>A), 19 (T>C), 21 (A>T), 22 (G>A), 23 (T>G), 24 (G>T), and 26 (T>G), 27 (A>T), 28 (G>A), 29 (A>G). Generally, the mechanisms of DNA mutation are substitution, deletion, and insertion (Sagermann et al. 2006). DNA mutation due to induction of gamma irradiation is random, cannot be targeted at DNA with a particular sequence number. The change in the DNA sequence also affects both secondary and tertiary structures. Changes in the structure of the DNA fragments can affect the genetic information encoded. The negative impact is chromosome aberration and instability of the genome (Koturbash et al. 2006). The point mutation that occurs in the *Nix* gene will be translated into different *Nix* protein than normal. There are three possibilities. Firstly, a silent mutation, which is a change in a base pair in the gene (at position 3 of the codon) which causes a change in one genetic code but does not result in changes of the encoded amino acid. Secondly, a missense mutation, which is a change in genetic code (generally at positions 1 and 2 at the codon). There is a change in the amino acids that are different from the natural amino acids making up the *Nix* protein. The last, nonsense mutation, which is a change in certain amino acid codons to be a stop codon. Almost all meaningless mutations lead to the inactivity of a protein resulting in a mutant phenotype. Most of the nonsense mutation will produce a protein that has an amino acid chain shorter than normal, so the protein produced will not working properly (Arodz et al. 2012). It should be done in further research to find out the changes in amino acids that occur due to a point mutation in the *Nix* gene.

The genetic factor plays an important role in any system of living organisms. Almost all structures and functions are controlled by genes, including diseases. In human, changes in genes both mutations and polymorphisms have an impact on their ability to survive. Most of the research showed the link between genetic mutations to risk factors of disease. For examples, the results of an epigenetic study demonstrated that Ala55Val and 45 bp I/D UCP2 polymorphisms play a role in increasing the risk of obesity in human (Surniyantoro et al. 2018), the other previous studies by Surniyantoro et al. (2018) stated that heterozygous allele 194Trp of XRCC1 exon 6 polymorphism showed a significantly higher micronuclei frequency in human exposure to ionizing radiation than controls. Micronuclei act as a biomarker of DNA damage. It is also demonstrated that ionizing radiation exposure, such as gamma radiation, can influence and damage the DNA fragments. Whereas, in *Ae. aegypti*, an F1552/C1552 point mutation was linked with permethrin resistance in the *Ae. aegypti* voltage-gated sodium channel. There was a single-base substitution (T>G) at the second position of codon 1552 resulting in the replacement of Phenylalanine by Cysteine in segment 6 domain III (Yanola et al. 2010). Saavedra-Rodriguez et al. (2007) also stated that a transition in codon 1011 encodes a Val replacement while a transition in codon 1016 encodes an Iso replacement. These mutations cause pyrethroid resistance in Latin American *Ae. aegypti*. All of the supporting studies above showed the importance of genetic factors in living organisms. Ionizing radiation (IR) exposure is a mutagenic agent, in addition to chemicals, which are widely studied, that produce various types of DNA damage, from single-strand breaks, modified nucleotides to double-strand breaks (Adewoye et al. 2015). Each of DNA damage has its own repair system, depending on the type of damage (Cooper 2000). The level of DNA damage is affected by the dose and dose-rate. A 70 Gy of the dose was the optimal dose for attenuating *Ae. aegypti* in the sterile insect technique (SIT) (Nurhayati et al. 2009). In human DNA, a previous study stated an equivalent dose has a significant correlation with micronuclei frequency among radiation-exposed workers, increasing the micronuclei frequency by 16.3 per 1 mGy of equivalent dose (Surniyantoro et al. 2018). This proves that the DNA damage tended to rise with equivalent dose, but various results in the different level of dose-rates.

Gamma irradiation at the pupa stage can also reduce sperm quantity. While in the mating competitiveness parameters, gamma irradiation at high doses can reduce the competitiveness of the mosquito species mating. Bellini et al. (2013) reported a semi-field test of the competitiveness of *Ae. albopictus* mosquito mating after gamma 30-60 Gray (Gy) dose irradiation. The results showed that competitiveness decreased with increasing gamma irradiation doses. Oliva et al. (2013) reported the effects of gamma irradiation on the age of *Ae. albopictus* mosquitoes in a semi-field trial on Reunion Island, one of the French regions in the Indian Ocean. The results showed that the age of mosquitoes in gamma irradiation was lower compared to field mosquitoes.

Some studies also report on the effects of gamma irradiation on *Ae. aegypti* mosquito species. Nurhayati et al. (2009) reported the effect of gamma irradiation on sterility. The results of the study stated that gamma irradiation had an effect on these two parameters. The optimum dose of gamma irradiation in *Ae. aegypti* mosquitoes ranges from 65-70 Gy. Sasmita et al. (2015) reported the effect of gamma irradiation on several quality parameters of *Ae. aegypti* mosquitoes, including the percentage of the appearance of adult mosquitoes, age of mosquitoes, sterility and mating competitiveness. The results of the study stated that gamma irradiation had no effect on the percentage of the appearance of adult mosquitoes. Gamma irradiation doses affect the age of mosquitoes, sterility and mating competitiveness. The results also provide information on the optimum dose of gamma irradiation in *Ae. aegypti* mosquitoes ranges from 60-80 Gy. Shetty et al. (2016) reported the effect of gamma irradiation on doses of 0-50 Gy on fertility, egg hatching, appearance and age of *Ae. aegypti* mosquitoes. The result of the study stated that gamma irradiation at low doses (< 2 Gy) had no effect on the parameters of the study.

In the context of SIT, gamma irradiation is used to induce sterility in the reproductive system of insects. gamma irradiation can damage chromosomes in gonadal cells, so chromosome fragmentation can occur and cause lethal dominant mutations, translocations and other chromosome aberrations (Robinson 2002). Chromosome damage results in unbalanced production of the gamete and subsequent zygote or embryo deaths occur (Klassen 2005; Robinson 2005). The impact of gamma irradiation on somatic cells is expressed in the development of abnormal organs, decreased ability to obtain nutrients, life skills, ability to fly, and ability to mate in these insects (Bakri et al. 2005).

In conclusion, our study reported that there are six and nineteen point mutations on the forward-strand and reverse-strand, respectively, in the *Nix* gene of males *Ae. aegypti* caused by gamma irradiation exposure. This result can be used as a pilot study to determine various kinds of genetic mutation information on *Ae. aegypti*'s genes. In the subsequent studies, it is necessary to examine the changes in amino acids that occur due to a point mutation in the *Nix* gene and the gene expression which can be used to develop the dengue vector *Ae. aegypti* decreasing program.

## ACKNOWLEDGEMENTS

This work was fully supported by Center for Isotopes and Radiation Application (PAIR), National Nuclear Energy Agency of Indonesia (BATAN), Jakarta. We are also grateful to Mrs. Devita Tetriana, M.Si. for assistance with Real-Time PCR technique and our co-workers in Entomology Laboratory, PAIR and Molecular Biology Laboratory, PTKMR, BATAN, Jakarta for technical support.

## REFERENCES

- Adeyoye AB, Lindsay SJ, Dubrova YE, Hurlles ME. 2015. The genome-wide effects of ionizing radiation on mutation induction in the mammalian germline. *Nat Commun* 6 (1): 6684
- Arodz T, Płonka PM. 2012. Effects of point mutations on protein structure are nonexponentially distributed. *Proteins* 80 (7): 1780-1790.
- Bakri A, Mehta K, Lance DR. 2005. Sterilizing insects with ionizing radiation. In Dyck VA, Robinson AS. (eds.) *Sterile Insect Technique*. Springer, Netherland.
- Balestrino F, Puggioli A, Gilles JRL, Bellini R. 2014. Validation of a new larval rearing unit for *Aedes albopictus* (Diptera: Culicidae) mass rearing. *PLoS ONE* 9 (3): e91914. DOI: 10.1371/journal.pone.0091914
- Bellini R, Balestrino F, Medici A, Gentile G, Veronesi R, Carrieri M. 2013. Mating competitiveness of *Aedes albopictus* radio-sterilized males in large enclosures exposed to natural conditions. *J Med Entomol* 50 (1): 94-102.
- Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, Drake JM, Brownstein JS, Hoen AG, Sankoh O, Myers MF, George DB, Jaenisch T, Wint GRW, Simmons CP, Scott TW, Farrar JJ, Hay SI. 2013. The global distribution and burden of dengue. *Nature* 496 (7446): 504-507.
- Bourtzis K, Lees RS, Hendrichs J, Vreysen MJB. 2016. More than one rabbit out of the hat: Radiation, transgenic and symbiont-based approaches for sustainable management of mosquito and tsetse fly populations. *Acta Trop* 157: 115-130.
- Cooper, GM. 2000. *The cell: a molecular approach*. Sinauer Associates, Sunderland.
- Chadee DD, Martinez R. 2016. *Aedes aegypti* (L.) in Latin American and Caribbean region: With growing evidence for vector adaptation to climate change? *Acta Trop* 156: 137-143.
- Dyck VA. 2005. *Sterile Insect Technique: Principles and Practice in Area-Wide Integrated Pest Management*. Springer, Dordrecht.
- Ernawan B, Tambunan USF, Sugoro I, Sasmita HI. 2017. Effects of gamma irradiation dose-rate on sterile male *Aedes aegypti*. AIP Conference Proceedings. Surabaya, June 26, 2017 [Indonesian].
- Ernawan B, Sasmita HI, Parikesit AA. 2018. Sterility of male *Aedes aegypti* post  $\gamma$ -ray sterilization. *J Anim Plant Sci* 28 (4): 973-977
- Fontaine A, Filipovic I, Fansiri T, Hoffmann AA, Cheng C, Kirkpatrick M, Rasic G, Lambrechts L. 2017. Extensive genetic differentiation between homomorphic sex chromosomes in the mosquito vector, *Aedes aegypti*. *Genome Biol Evol* 9 (9): 2322-35.
- Gubler DJ. 2004. The changing epidemiology of yellow fever and dengue, 1900 to 2003: full circle? *Comp Immunol Microbiol Infect Dis* 27 (5): 319-330.
- Hall AB, Basu S, Jiang X, Qi Y, Timoshevskiy VA, Biedler JK, Sharakhova MV, Elahi R, Anderson MAE, Chen X, Sharakov IV, Tu Z. 2015. A male-determining factor in the mosquito *Aedes aegypti*. *Science* 348 (6240): 1268-1270
- Hall AB, Timoshevskiy VA, Sharakhova MV, Jiang X, Basu S, Anderson MAE, Hu W, Sharakhov IV, Adelman ZN, Tu Z. 2014. Insights into the preservation of the homomorphic sex-determining chromosome of *Aedes aegypti* from the discovery of a male-biased gene tightly linked to the M-locus. *Genome Biol Evol* 6 (1): 179-191.
- Hapairai LK, Marie J, Sinkins SP, Bossin HC. 2014. Effect of temperature and larval density on *Aedes polynesiensis* (Diptera: Culicidae) laboratory rearing productivity and male characteristics. *Acta Trop* 132 (1): 108-115.
- Harrington LC, Scott TW, Lerdthusnee K, Coleman RC, Costero A, Clark GG, Jones JJ, Kitthawee S, Kittayapong P, Sithiprasasna R, Edman JD. 2005. Dispersal of the dengue vector *Aedes aegypti* within and between rural communities. *Am J Trop Med Hyg* 72 (2): 209-220.
- Jansen, CC, Beebe NW. 2010. The dengue vector *Aedes aegypti*: what comes next. *Microbes Infect* 12 (4): 272-279.
- Kemenkes RI. 2016. Wilayah KLB ada di 11 Provinsi. Kemenkes RI, Jakarta. [Indonesian]
- Klassen W. 2005. Area-wide integrated pest management and the sterile insect technique. In: Dyck VA, Robinson AS. (eds.) *Sterile Insect Technique*. Springer, Netherland.
- Koturbash I, Rugo RE, Hendricks CA, Loree J, Thibault B, Kutanzi K, Pogribny I, Yanch JC, Engelward BP, Kovalchuk O. 2006. Irradiation induces DNA damage and modulates epigenetic effectors in distant bystander tissue in vivo. *Oncogene* 25 (31): 4267-75.
- Li MI, Wong PSJ, Ng LC, Tan CH. 2012. Oral susceptibility of Singapore *Aedes (Stegomyia) aegypti* (Linnaeus) to Zika Virus. *PLoS Negl Trop Dis* 6 (8): e1792. DOI: 10.1371/journal.pntd.0001792.
- Maestre-Serrano R, Gomez-Camargo D, Ponce-Garcia G, Flores AE. 2014. Susceptibility to insecticides and resistance mechanisms in *Aedes aegypti* from the Colombian Caribbean Region. *Pestic Biochem Physiol* 116: 63-73.
- Moyes CL, Vontas J, Martins AJ, Ng LC, Koou SY, Dusfour I, Raghavendra K, Pinto J, Corbel V, David JP, Weetman D. 2017. Contemporary status of insecticide resistance in the major *Aedes* vectors of arboviruses infecting humans. *PLoS Negl Trop Dis* 11 (7): e0005625. DOI: 10.1371/journal.pntd.0005625.
- Nene V, Wortman JR, Lawson D, Haas B, Kodira C, Tu ZJ, Loftus B, Xi Z, Megy K, Grabherr M, Ren Q, et al. 2007. Genome sequence of *Aedes aegypti*, a major arbovirus vector. *Science* 316 (5832): 1718-1723.
- Ngoagouni C, Kamgang B, Brengues C, Yahouedo G, Paupy C, Nakoune E, Kazanji M, Chandre F. 2016. Susceptibility profile and metabolic mechanisms involved in *Aedes aegypti* and *Aedes albopictus* resistant to DDT and deltamethrin in the Central African Republic. *Parasit Vectors* 9 (1): 599.
- Nurhayati S, Santoso B, Rahayu A, Tetriana D. 2009. Effect of gamma rays radiation on the competitiveness of *Aedes aegypti* mosquitoes as a dengue hemorrhagic fever vector. SNKKL V Proceeding, University of Indonesia, Depok, October 14, 2009. [Indonesian]
- Oliva CF, Maier MJ, Gilles J, Jacquet M, Lemperiere G, Quilici S, Vreysen MJB, Schooneman F, Chadee DD, Boyer S. 2013. Effects of irradiation, presence of females, and sugar supply on the longevity of sterile males *Aedes albopictus* (Skuse) under semi-field conditions on Reunion Island. *Acta Trop* 125 (3): 287-293.
- Rezza G, Nicoletti L, Angelini R, Romi R, Finarelli AC, Panning M, Cordioli P, Fortuna C, Boros S, Magurano F, Silvi G, Angelini P, Dottori M, Ciufolini MG, Majori GC, Cassone A. 2007. Infection with chikungunya virus in Italy: an outbreak in a temperate region. *The Lancet* 370 (9602): 1840-1846.
- Robinson AS. 2002. Mutations and their use in insect control. *Mutat Res* 511 (2): 113-132.
- Robinson AS. 2005. Genetic basis of the sterile insect technique. In Dyck VA, Robinson AS. (eds.) *Sterile Insect Technique*. Springer, Netherland.
- Saavedra-Rodriguez K, Urdaneta-Marquez L, Rajatileka S, Moulton M, Flores AE, Fernandez-Salas I, Bisset J, Rodriguez M et al. 2007. A mutation in the voltage-gated sodium channel gene associated with pyrethroid resistance in Latin American *Aedes aegypti*. *Insect Mol Biol* 16 (6): 785-98.
- Sagermann M, Baase WA, Matthews BW. 2006. Sequential reorganization of  $\beta$ -sheet topology by insertion of a single strand. *Protein Sci* 15 (5): 1085-1092.
- Sasmita HI, Ernawan B. 2015. Quality of Gamma-Sterilized Male of *Aedes aegypti* L.: The Effect of Irradiating on Pupal and Adult Stage. *J. Appl Isotopes Rad* 10 (2): 149-58.
- Shetty V, Shetty NJ, Harini BP, Ananthanarayana SR, Jha SK, Chaubey RC. 2016. Effect of gamma radiation on life history traits of *Aedes aegypti* (L.). *Parasite Epidemiol Control*. 1 (2): 26-35.
- Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Soding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* 7: 539. DOI: 10.1038/msb.2011.75.
- Surniyantoro HNE, Sadewa AH, Hastuti P. 2018. Uncoupling Protein 2 (UCP2) as Genetic Risk Factor for Obesity in Indonesia is Different in Gender Stratification. *Kobe J Med Sci* 64 (2): 64-72.
- Surniyantoro HNE, Darlina, Nurhayati S, Tetriana D, Syaifudin M. 2016. Gamma ray irradiation inhibits *Plasmodium falciparum* multiplication in in vitro culture supplemented with tritium labeled hypoxanthine. *Nusantara Biosci* 8 (1): 8-13.
- Surniyantoro HNE, Lusiyanti Y, Rahardjo T, Nurhayati S, Tetriana D. 2018. Association between XRCC1 exon 10 (Arg399Gln) gene polymorphism and micronucleus as a predictor of DNA damage among radiation workers. *Biodiversitas* 19 (5): 1676-1682
- Surniyantoro HNE, Lusiyanti Y, Rahardjo T, Tetriana D, Nurhayati S, Date H. 2018. Polymorphism of XRCC1 Gene Exon 6 (Arg194Trp) in Relation to Micronucleus Frequencies in Hospital Radiation Workers. *Atom Indonesia* 44 (2): 105-111.
- Syaifudin M, Darlina, Rahardjo T, Tetriana D, Nurhayati S, Surniyantoro HNE, Kisnanto T. 2013. Effectiveness of gamma rays in attenuating

- rodent malaria parasites of *Plasmodium berghei* in blood of mice. *Atom Indonesia* 39 (1): 19-23.
- Van Bortel W, Dorleans F, Rosine J, Blateau A, Rousset D, Matheus S, Leparc-Goffart I, Flusin O, Prat CM, Césaire R, Najioullah F, Ardillon V, Balleydier E, Carvalho L, Lemaître A, Noël H, Servas V, Six C, Zurbaran M, Léon L, Guinard A, Van den Kerkhof J, Henry M, Fanoy E, Braks M, Reimerink J, Swaan C, Georges R, Brooks L, Freedman J, Sudre B, Zeller H. 2014. Chikungunya outbreak in the Caribbean region, December 2013 to March 2014, and the significance for Europe. *Euro Surveill* 19 (13): 20759.
- Vorou, R. 2016. Zika virus, vectors, reservoirs, amplifying hosts, and their potential to spread worldwide: what we know and what we should investigate urgently. *Intl J Infect Dis* 48: 85-90.
- Weaver SC, Reisen WK. 2010. Present and future arboviral threats. *Antiviral Res* 85 (2): 328-345.
- WHO. 2011. Comprehensive Guidelines for Prevention and Control of Dengue and Dengue Haemorrhagic Fever. WHO-India, New Delhi.
- Wiwanitkit V. 2016. Emerging Zika virus infection: What should we know? *Asian Pac J Trop Biomed* 6 (7): 551-553.
- Yanola J, Somboon P, Walton C, Nachaiwieng W, Prapanthadara L. 2010. A novel F1552/C1552 point mutation in the *Aedes aegypti* voltage-gated sodium channel gene associated with permethrin resistance. *Pest Biochem Physiol* 96 (3): 127-31.
- Zheng M, Zhang D, Damiens DD, Lees RS, Gilles JRL. 2015. Standard operating procedures for standardized mass rearing of the dengue and chikungunya vectors *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae)-II-Egg storage and hatching. *Parasit Vectors* 8: 348. DOI: 10.1186/s13071-015-0951-x.