Anticancer mechanism of artonin E and related prenylated flavonoids from the medicinal plant *Artocarpus elasticus*

CHRISTIAN BAILLY

OncoWitan, Scientific Consulting Office, Lille (Wasquehal), 59290, France. ✉email: christian.bailly@oncowitan.com

Manuscript received: 23 June 2021. Revision accepted: 25 July 2021.

Abstract. Bailly C. 2021. Anticancer mechanism of artonin E and related prenylated flavonoids from the medicinal plant *Artocarpus elasticus*. *Asian J Nat Prod Biochem* 19: 45-57. Plants of the *Artocarpus* genus are largely distributed throughout tropical Asia and Oceania. Species such as *A. altilis* (Parkinson) Fosberg and *A. heterophyllus* Lam. are popular trees known as breadfruit and jackfruit, respectively. They contain a large structural diversity of bioactive prenylated flavonoids. Here we have focused on the less known species *Artocarpus elasticus* Reinw. ex Blume is well distributed in southeast Asia and used in traditional medicine to treat dysentery, tuberculosis, and other diseases. Numerous prenylated flavonoids have been isolated from the leaves and bark of *A. elasticus*, such as artocarpesin, artocarpin, artelastin, and many others. They are endowed with antioxidant, anti-inflammatory, and anticancer properties. A focus is made on the subgroup of compounds designated artonins, with the derivative artonin E as a lead anticancer agent. Art-E has revealed marked anticancer effects after oral administration in vitro and in vivo. The mechanism of action is discussed to highlight the structural and functional analogy between Art-E and the antitumor natural product morusin. Both compounds trigger TRAIL-mediated apoptosis of cancer cells. They can be considered further for the development of novel anticancer agents.

Keywords: *Artocarpus*, cancer therapeutics, natural products, phytotherapy, prenylated flavonoids

Abbreviations: iNOS, inducible nitric oxide synthase; ROS, reactive oxygen species

INTRODUCTION

The genus *Artocarpus* refers to a large group of plants belonging to the Moraceae family (Zerega et al. 2010). The internet database www.theplantlist.org has recorded 68 taxonomically accepted species, while about 90 additional plant names are now in synonymy or have even been moved as synonyms to other genera. *Artocarpus* derives from the Greek *artos* for “bread” and *Karpos* for “fruit.” Some of the *Artocarpus* species are relatively well-known, such as *A. altilis* (Parkinson) Fosberg, known as the breadfruit tree, and *A. heterophyllus* Lam. known as the jackfruit tree, which are two economically important *Artocarpus* species (Sikarwar et al. 2014; Mohammed Haleel et al. 2018; Sahu et al. 2019; Buddhisuharto et al. 2021). Their fruits are used for food production, and several parts of these two trees, including fruits, leaves, and barks, have been used in traditional medicine in India and surrounding countries (Ranasinghe et al., 2019). Other *Artocarpus* species are used in traditional medicine, such as (i) decoctions prepared from the leaves of the tree *A. tonkinensis*, used in northern Vietnam to treat arthritis and backache (Adorisio et al. 2016), and (ii) extracts of *Artocarpus lacucha* (also known as Monkey jack) used in traditional Thai medicine as an anthelmintic agent (Aneklaphakij et al. 2020; Gupta et al. 2020), for examples. The ethnopharmacological use of *Artocarpus* species is widespread in subtropical and tropical regions of Asia, where the trees are largely distributed. *Artocarpus* is believed to originate from the island of Borneo, from which species dispersed and diversified in several directions

(Williams et al. 2017). *Artocarpus*-based remedies are used to treat multiple diseases and conditions, including malarial fever, diarrhea, diabetes, parasitic infections, and different inflammatory diseases (Jagtap and Bapat 2010).

These plants contain a large diversity of bioactive compounds, including many prenylated flavonoids and cyclized derivatives which have revealed interesting antimicrobial, anti-inflammatory and anticancer properties (Hakim et al., 2006; Hari et al., 2014). One particular species, *Artocarpus elasticus* Reinw. ex Blume (Figure 1) has shown marked anticancer effects against different cancer cell lines and tumor models, attributed to the presence of prenylated flavonoids isolated from the root bark or other parts of the tree. The lead compound is artonin E (Art-E), which was thoroughly investigated as an antitumor agent. Over the past ten years, the mechanism of action of Art-E has been delineated, and the activated signaling pathways at the origin of antitumor activity have been defined. These studies gave us the impetus to deeply analyze the mode of action of Art-E and structurally related compounds. The present review will address successively the plant *A. elasticus* and plant extracts, the phytochemical compounds isolated from the plant, the specific flavonoid group known as artonins, and the molecular targets and pathways activated by Art-E responsible for its pharmacological properties. The role of TRAIL in the mode of action of Art-E and related compounds is discussed, opening the door to the identification of novel anticancer molecules. The purpose of the study was to promote the knowledge of artonins and the use of artonin E as a lead anti-cancer compound.

Figure 1. *Artocarpus elasticus* Reinw. ex Blume as a source of anticancer prenylated flavonoids. Different *Artocarpus* species are used in traditional medicine and as food products, such as *A. altilis* (Parkinson ex F.A.Zorn) Fosberg, *A. heterophyllus* Lam., *A. tonkinensis* A.Chev. ex Gagnep. and *A. lacucha* Buch.-Ham. Similarly, *A. elasticus* is also a source of bioactive mono-, bis-, and tris-prenylated flavonoids endowed with anticancer properties. *A. blumei* Trécul and *A. kunstleri* King are synonyms for *A. elasticus* Reinw. ex Blume (<http://www.theplantlist.org/>)

ARTOCARPUS ELASTICUS AND ITS MEDICINAL USES

The plant *Artocarpus elasticus* Reinw. ex Blume is native to Southeast Asia and can easily be found in countries like Myanmar, Thailand, Malaysia, Indonesia, and the Philippines. In some countries like Malaysia and Indonesia, *A. elasticus* is cultivated to maintain sustainability (Susiarti et al. 2020). It is an evergreen, robust, tall tree (up to 45-65 m). Various vernacular names are used locally, such as Benda (Javanese Indonesia), Terap nasi (Peninsular Malaysia), and others (Teo and Nasution 2003).

The wood of *A. elasticus* is sought as a building material. It exudes white and thick latex, elastic or flexible upon wounding, hence the name *elasticus*. The sticky latex is used to catch birds but also to treat dysentery. Different parts of the tree are used in traditional medicines, the latex for dysentery, the bark for female contraception, and the young leaves mixed with rice to treat tuberculosis. The flesh of the fruits is eaten raw or cooked. The seeds are consumed fried or roasted (Susiarti et al. 2020).

The bark and the leaves of the plant are often used to identify new phytochemicals. In addition, an extract of the stem bark of *A. elasticus* has been used as a reducing and stabilizing agent for synthesizing silver nanoparticles (Abdullah et al., 2015). The same antibacterial silver nanoparticles have been obtained with seed extract of *A. hirsutus* (Shobana et al., 2020).

PRENYL FLAVONOIDS FROM ARTOCARPUS ELASTICUS

Artocarpus elasticus is a rich reservoir of bioactive natural products, notably prenyl flavonoids, particularly abundant in the tree's bark and the roots. There is a large structural diversity of prenylated flavonoids isolated from *A. elasticus*, with compounds bearing one, two, or three prenyl groups, a classical flavonoid skeleton, or a bulkier pentacyclic core (Nomura et al., 1998; Kijjoa et al. 1998; Cidade et al. 2001). The main isolated compounds are shown in Figure 2, except the artonins discussed below. Most of these compounds can be found in other *Artocarpus* species. Here, we will discuss only those found in *A. elasticus*.

The tetracyclic compound artelastin, with three prenyl side chains, was isolated in 1996 from the wood of *A. elasticus* together with artelastochromene, artelasticin, and artocarpesin (Kijjoa et al., 1996). Artelastin has revealed marked cytotoxic properties against different cancer cell lines *in vitro*. It was found to disturb the microtubule network of MCF7 breast cancer cells and interfere with the cell cycle progression. Artelastin caused an accumulation in S phase due to a drug-induced delay in DNA replication (Pedro et al. 2005). It is a potent inhibitor of both T- and B lymphocyte mitogen-induced proliferation and an inhibitor of cytokines production, such as interferon- γ , interleukins IL-2, -4, and -10, in stimulated splenocytes (Cerqueira et al. 2003). Artelastin displays marked antioxidant properties, inhibiting the production of reactive oxygen species (ROS) and the expression of the inducible nitric oxide synthase (iNOS) in lipopolysaccharide-stimulated macrophages (Cerqueira et al. 2008).

Figure 2. Chemical structures of prenylated flavonoids which have been isolated from *A. elasticus* Reinw. ex Blume

Artelasticin can be found in *A. elasticus* (Kijjoa et al. 1996), *A. lanceifolius* (Syah et al. 2001), and other plants, such as the Cameroon medicinal plant *Dorstenia psilurus*. It was found to activate AMPK (AMP-activated protein kinase) and stimulate glucose uptake in rat skeletal muscles. Its marked blood-glucose-lowering effect can be useful for treating type-2 diabetes (Choumessi et al., 2019). Artocarpesin was found to be moderately cytotoxic toward a panel of cancer cell lines (IC_{50} values in the 60-100 μM range), but its analog cycloartocarpesin is more potent (IC_{50} values in the 15-50 μM range) (Kuate et al. 2015). This compound can be found in many other plants. Like artelasticin, it presents marked anti-inflammatory and antioxidant properties (Fang et al. 2008). Its antioxidant activity contributes to the anti-browning effect observed with extracts of *A. heterophyllus* on fresh-cut apple slices. Remarkably, artocarpesin was found to exert a strong inhibitory activity against mushroom tyrosinase (IC_{50} = 0.52 μM), whereas the related product artocarpin was inactive against the enzyme (Zheng et al. 2008, 2009). Artocarpesin, found in the twigs and woods of *A. heterophyllus*, *A. incisus*, and *A. elasticus*, has been considered for the design of skin whitening agents (Arung et al. 2008, 2011). Artocarpin has been reported in a dozen of *Artocarpus* species and displays antioxidant, anti-inflammatory, and anti-parasitic properties, as well as skin-whitening activities (Chan et al. 2018a; Morrison et al. 2021).

The bis-prenyl flavonoid cyclocommunin has been known for a long time for its potent capacity to inhibit platelet aggregation induced by collagen or arachidonic acid (Lin et al. 1993). It was first isolated from *A. communis*, hence its name cyclocommunin (Lin and Shieh,

1992), but it is also called isocyclomulberrin (Chen et al. 1993; Ma et al. 2010). Cyclocommunin is also an activator of protein kinase C (PKC) and a compound capable of elevating the level of intracellular calcium in rat neutrophils. As a result, cyclocommunin was found to stimulate a respiratory burst in neutrophils (Wang et al. 1999), enhancing the production of superoxide anion, unlike artonin B, which inhibited superoxide anion formation in these cells (Wei et al. 2005). Cyclocommunin displays modest anti-proliferative effects against hepatocellular carcinoma cells (IC_{50} = 14-33 μM depending on cell line). In contrast, Art-A and Art-B were inactive against these HCC cell lines (Ma et al. 2010). Recently, it was reported that cyclocommunin displays a significant antimycobacterial activity against *Mycobacterium tuberculosis* (laboratory strain H37Ra) with a minimum inhibitory concentration (MIC) of 12.3 μM (Boonyaketgason et al. 2020).

Artelastocarpin and carpelastofuran were isolated from the wood of *A. elasticus*. They both revealed cytotoxic properties toward cancer cells, with an efficacy like that of artelastin, artelasticin, and artelastochromene (IC_{50} = 7-12 μM depending on cell lines) (Cidade et al. 2001). Different *Artocarpus* species have obtained a series of compounds named artoindonesianins (Musthapa et al. 2010). Most of them have been isolated from species other than *A. elasticus*, such as the cytotoxic compounds artoindonesianins Z-4 and Z-5 found in the bark of *A. lanceifolius* (Musthapa et al. 2009a) and artoindonesianins A and B found in the roots of *A. champeden* (Hakim et al. 1999) and many other artoindonesianins, as indicated in Table 1.

Table 1. Artoindonesianin compounds isolated from *Artocarpus* species

Compound	<i>Artocarpus</i> species	References
Artoindonesianin A	<i>A. champeden</i>	Hakim et al. 1999
Artoindonesianin B	<i>A. champeden</i>	Hakim et al. 1999
Artoindonesianin C	<i>A. teysmanii</i>	Makmur et al. 2000
	<i>A. rigidus</i>	Namdaung et al. 2006 ;
	<i>A. kemando</i>	Ee et al. 2011
Artoindonesianin D	<i>A. maingayi</i>	Hakim et al. 2000
Artoindonesianin E	<i>A. champeden</i>	Hakim et al. 2001
Artoindonesianin F	<i>A. heterophyllum</i>	Rao et al. 2010
Artoindonesianin G	<i>A. lanceifolius</i>	Syah et al. 2001
Artoindonesianin H	<i>A. lanceifolius</i>	Syah et al. 2001
Artoindonesianin I	<i>A. lanceifolius</i>	Syah et al. 2001
Artoindonesianin J	<i>A. bracteata</i>	Ersam et al. 2002
Artoindonesianin K	<i>A. champeden</i>	Achmad et al. 2005
Artoindonesianin L	<i>A. rotunda</i>	Suhartati et al. 2001
Artoindonesianin M	<i>A. champeden</i>	Syah et al. 2002b
Artoindonesianin N	<i>A. gomezianus</i>	Hakim et al. 2002a
Artoindonesianin O	<i>A. gomezianus</i>	Hakim et al. 2002a
Artoindonesianin P	<i>A. lanceifolius</i>	Hakim et al. 2002b
	<i>A. elasticus</i>	Jenis et al. 2019
Artoindonesianin Q	<i>A. champeden</i>	Syah et al. 2002a
Artoindonesianin R	<i>A. champeden</i>	Syah et al. 2002a
Artoindonesianin S	<i>A. champeden</i>	Syah et al. 2002a
Artoindonesianin T	<i>A. champeden</i>	Syah et al. 2002a
Artoindonesianin U	<i>A. champeden</i>	Syah et al. 2004
Artoindonesianin V	<i>A. champeden</i>	Syah et al. 2004
	<i>A. altilis</i>	Shamaun et al. 2010
Artoindonesianin W	<i>A. elasticus</i>	Jenis et al. 2019
Artoindonesianin X	<i>A. fretessi</i>	Soekamto et al. 2003
Artoindonesianin Y	<i>A. fretessi</i>	Soekamto et al. 2003
Artoindonesianin Z1	<i>A. lanceifolius</i>	Hakim et al. 2006
	<i>A. anisophyllum</i>	Noraini et al. 2013
Artoindonesianin Z2	<i>A. lanceifolius</i>	Hakim et al. 2006
Artoindonesianin Z3	<i>A. lanceifolius</i>	Hakim et al. 2006
Artoindonesianin Z4	<i>A. lanceifolius</i>	Musthapa et al. 2009b
Artoindonesianin Z5	<i>A. lanceifolius</i>	Musthapa et al. 2009b
Artoindonesianin Z4	<i>A. lanceifolius</i>	Musthapa et al. 2009b
Artoindonesianin Z5	<i>A. lanceifolius</i>	Musthapa et al. 2009b
Artoindonesianin A1	<i>A. champeden</i>	Syah et al. 2006b
Artoindonesianin A2	<i>A. champeden</i>	Syah et al. 2006a
Artoindonesianin A3	<i>A. champeden</i>	Syah et al. 2006a
Artoindonesianin B1	<i>A. altilis</i>	Syah et al. 2006b
	<i>A. heterophyllum</i>	Lang et al. 2016
Artoindonesianin E1	<i>A. elasticus</i>	Musthapa et al. 2009a

For the record, the structure of artoindonesianin C, first isolated from *A. teysmanii* (Makmur et al. 2000), was printed on a post stamp on the occasion of the international year of chemistry 2011 (Figure 3). The oxepinoflavone derivative artoindonesianin E1 was isolated from the bark of *A. elasticus*, together with the related compounds artocarpin, cycloartocarpin, and cudraflavones A and C (Figure 2). The derivative artoindonesianin A-3 was initially isolated from *A. champeden* (Syah et al. 2006a) and later re-discovered from the bark of *A. elasticus* tree collected from Alor Island in Indonesia (Kuran and Ersam 2017). Other artoindonesianins have been found in *A. elasticus*, such as artoindonesianin P, which was found to inhibit the enzyme α -glucosidase (IC_{50} = 25.4 μ M) but

with a reduced efficacy compared to Art-E and artobioxanthone (IC_{50} = 16.2 and 8.6 μ M, respectively) (Jenis et al. 2019). Recently, artoindonesianins P and W were isolated from *A. elasticus* and found to inhibit human neutrophil elastase (IC_{50} = 28.7 and 11.2 μ M, respectively), whereas artobioxanthone was slightly more active (IC_{50} = 9.8 μ M) (Ban et al. 2020).

Other prenylated flavonoids have been identified, such as artelastoheterol, artelasticinol, artelastoxanthone, artobioxanthone, cycloartelastoxanthone, cycloartobioxanthone, cycloartelastoxanthendiol, elastixanthone, and artonol A, all isolated from the root bark of *A. elasticus* (Figure 2). Artelastoxanthone has revealed modest antiproliferative activities against different cancer cell lines *in vitro* (Ko et al. 2005). Cycloartelastoxanthone, cycloartobioxanthone, and artobioxanthone have protective effects on DNA damage caused by superoxide anion radicals $O_2^{\cdot-}$ (Lin et al. 2009). Elastixanthone and cycloartobioxanthone have shown antimicrobial activities (Ramli et al., 2016). Artobioxanthone and cycloartobioxanthone have been found in many *Artocarpus* species, not limited to *A. elasticus*. The latter compound is interesting because it was found to potentially inhibit migration and invasion of H460 lung cancer cells via inhibition of the phosphorylation of focal adhesion kinase (FAK) and the expression of the cell division cycle 42 (CDC42) (Tungsukruthai et al. 2017). Cycloartobioxanthone dose-dependently reduces the proliferation of other lung cancer cell lines (H23, H292, and A549) and triggers caspase-dependent apoptosis with an efficacy comparable to that of the standard DNA-damaging anticancer drugs cisplatin and etoposide, at least *in vitro* (Losuwannarak et al. 2018). But the ROS scavenging activity of cycloartobioxanthone is much weaker than that of artobioxanthone and Art-E (Sritularak et al., 2010). A recent molecular docking analysis has predicted that artobioxanthone can bind to the active site of the enzyme transglutaminase 2 (TG2), possibly inhibiting this enzyme considered an anticancer target (Parvatikar and Madagi 2021).

Figure 3. The structure of artoindonesianin C, isolated from *A. teysmanii*. A post stamp with the structure of the compound was emitted to celebrate the international year of chemistry 2011 in Indonesia.

Here we focused mostly on prenylated flavonoids isolated from *A. elasticus*, but the structural diversity of the whole *Artocarpus* genus is huge, as previously reported in specific review articles on *Artocarpus* species (Achmad et al. 2005; Hakim et al. 2006; Hakim 2010; Veitch and Grayer 2008). A recent analysis of a root extract of *A. heterophyllus* led to the identification of 47 prenylated flavonoids (Ye et al., 2019). There must be that many in the related species *A. elasticus*. An important group of prenylated flavonoids commonly found in *Artocarpus* species is called artonin, with the lead compound Art-E. Given the importance of the lead, this family is presented hereafter.

THE ARTONIN GROUP OF PRENYL FLAVONOIDS

There are 24 compounds named artonins A to Y (Table 2). The first two compounds, Art-A and -B, were isolated

in 1989 from the root bark of *A. heterophyllus* (Hano et al. 1989), followed by Art-C and -D one year later from the same plant (Hano et al. 1990a). They can be found in a few other *Artocarpus* species and display antioxidant properties (Ko et al. 2005). Art-B is an interesting compound, much more potent than Art-A at inhibiting the proliferation of nasopharyngeal cancer KB cells and the growth of the parasite *Trypanosoma brucei brucei*, *in vitro* (Bourjot et al. 2010). Art-B can reduce the proliferation of different cancer cells, such as human CCRF-CEM leukemia cells, via the induction of mitochondria-dependent apoptosis (Lee et al. 2006). Art-A is less cytotoxic than Art-B. Nevertheless, it is a more potent inhibitor of the papain-like cysteine protease cathepsin K ($IC_{50} = 1.9$ and $9.0 \mu M$, for Art-A and Art-B, respectively), largely implicated in bone resorption (Zhai et al., 2017). Art-A also displays antimalarial activity, at least *in vitro* (Widyawaruyanti et al. 2007).

Table 2. Artonin compounds isolated from *Artocarpus* species

Compound	CID ^a	Formula (g/mol) ^b	<i>Artocarpus</i> species	References
Artonin A	14557102	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. heterophyllus</i> <i>A. styracifolius</i> <i>A. hypargyreus</i> <i>A. xanthocarpus</i> <i>A. champeden</i>	Hano et al. 1989 Bourjot et al. 2010 Qiao et al. 2011 Jin et al. 2015 Widyawaruyanti et al. 2007
Artonin B	11964501	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. heterophyllus</i> <i>A. champeden</i> <i>A. styracifolius</i>	Hano et al. 1989 Hakim et al. 1999s Bourjot et al. 2010
Artonin C	14681571	C ₄₀ H ₃₈ O ₁₀ (678.7)	<i>A. heterophyllus</i>	Hano et al. 1990a
Artonin D	14681573	C ₄₀ H ₃₆ O ₁₀ (676.7)	<i>A. heterophyllus</i>	Hano et al. 1990a
Artonin E	5481962	C ₂₅ H ₂₄ O ₇ (436.5)	<i>A. communis</i> <i>A. kemando</i> <i>A. lanceifolius</i> <i>A. chama</i> <i>A. nobilis</i> <i>A. lanceifolius</i> <i>A. gomezianus</i> <i>A. rotunda</i> <i>A. rigida</i>	Hano et al. 1990b Seo et al. 2003 Cao et al. 2003 Wang et al. 2004 Jayasinghe et al. 2008 Musthapa et al. 2009 Plaibua et al. 2013 Suhartati et al. 2001 Suhartati et al. 2018
Artonin F	14680593	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. rigidus</i> <i>A. styracifolius</i> <i>A. integer var. silvestris</i> Corner	Namdaung et al. 2006 Bourjot et al. 2010 Shaha et al. 2016
Artonin G	46887714	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. rigida</i>	Hano et al. 1990
Artonin H	21595104	C ₃₀ H ₃₄ O ₇ (506.6)	<i>A. rigida</i>	Hano et al. 1990
Artonin I	57335177	C ₄₀ H ₃₆ O ₁₁ (692.7)	<i>A. heterophyllus</i>	Hano et al. 1992
Artonin J	44258663	C ₂₅ H ₂₄ O ₇ (436.5)	<i>A. heterophyllus</i>	Aida et al. 1993
Artonin K	15340661	C ₂₁ H ₁₈ O ₇ (382.4)	<i>A. heterophyllus</i>	Aida et al. 1993
Artonin L	44258662	C ₂₂ H ₂₀ O ₇ (396.4)	<i>A. heterophyllus</i>	Aida et al. 1993
Artonin M	44258661	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. altilis</i> <i>A. rotunda</i>	Hano et al. 1993 Suhartati et al. 2001
Artonin N	44258669	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. rigida</i>	Hano et al. 1993
Artonin O	46887814	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. rigida</i> <i>A. rotunda</i>	Hano et al. 1993 Suhartati et al. 2001
Artonin P	44258658	C ₂₅ H ₂₀ O ₈ (448.4)	<i>A. rigida</i> <i>A. communis</i>	Hano et al. 1993 Chan et al. 2018b
Artonin Q	131753034	C ₃₁ H ₃₀ O ₈ (530.6)	<i>A. heterophyllus</i>	Aida et al. 1994
Artonin R	131753035	C ₃₁ H ₃₀ O ₁₀ (562.6)	<i>A. heterophyllus</i>	Aida et al. 1994
Artonin S	44258666	C ₂₆ H ₂₈ O ₇ (452.5)	<i>A. heterophyllus</i>	Aida et al. 1994
Artonin T	44258664	C ₂₆ H ₂₆ O ₇ (450.5)	<i>A. heterophyllus</i>	Aida et al. 1994
Artonin U	44258358	C ₂₁ H ₂₀ O ₅ (352.4)	<i>A. heterophyllus</i>	Aida et al. 1994
Artonin V	129687399	C ₂₅ H ₂₆ O ₇ (438.5)	<i>A. altilis</i>	Hano et al. 1994
Artonin X	(FDB021143)	C ₄₀ H ₃₈ O ₉ (662.7)	<i>A. heterophyllus</i>	Shinomiya et al. 1995
Artonin Y	15541482	C ₂₀ H ₁₈ O ₆ (354.4)	<i>A. heterophyllus</i>	Shinomiya et al. 2000

Note: ^aCompound Identity number (PubChem CID). ^bFormula and molecular weight.

The most important compound in the series is arguably Art-E (also known as 5'-hydroxymorusin), initially isolated from the stem bark of *A. communis* Forst., together with Art-F (Hano et al. 1990b). Later, it was found in several other *Artocarpus* species, including *A. elasticus* (Ramli et al. 2016). The compound, bearing an isoprenoid group at position C-3, was initially characterized as a potent and selective inhibitor of arachidonate 5-lipoxygenase (IC_{50} = 0.36 μ M) expressed on cells involved in the regulation of immune responses (Reddy et al., 1991). Art-E has revealed a very modest capacity to inhibit ADP-induced platelet aggregation *in vitro*, with an IC_{50} of 192 μ M (Jantan et al. 2010). Later, Art-E was found to exert potent antiproliferative activities against different cancer cell lines, notably against breast adenocarcinoma MCF-7 and MDA-MB-231 cells (ED_{50} = 2.2 and 3.0 μ g/ml, respectively) (Wang et al., 2004). It is slightly less active against other types of cancer cells derived from solid tumors but also exerts a potent cytotoxic action against P-388 leukemia cells (Musthapa et al. 2009b). Art-E is the lead product in the series, with interesting anticancer and antimicrobial properties (see below). The anticancer effects have been characterized using different experimental models *in vitro* and *in vivo*. In particular, the compound has revealed a robust dose-dependent effect in a 4T1 breast carcinoma xenograft model. The oral administration of Art-E at 25, 50, and 100 mg/kg drastically reduced tumor growth in mice and reduced the appearance of metastasis without any apparent toxicity (Etti et al. 2017a). The compound is well tolerated; it can be administered orally for up to 10 consecutive days at 30 mg/kg, without significantly affecting cholesterol, creatinine, and blood urea nitrogen levels (Fukai et al. 2003).

Modest antibacterial effects have been reported with Art-E against *Escherichia coli* and *Bacillus subtilis* (Suhartati et al. 2008) and Art-O against *B. subtilis* (Suhartati et al. 2016). Mild antibacterial effects have been noted with Art-E against *Pseudomonas aeruginosa* strain PA01 (MIC = 32 μ g/ml). In contrast, the compound was essentially inactive against the microorganisms *Providencia stuartii*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, *Salmonella typhi*, and *Escherichia coli* (Kuate et al., 2011). Art-E showed little activity against *S. aureus* strain ATCC25922 (MIC = 256 μ g/ml) (Kuate et al. 2011) whereas a significant antibacterial effect has been reported with the strain ATCC 25923 (MIC = 3.9 μ g/ml) and the methicillin-resistant strain ATCC BAA-1720 (MIC = 13.3 μ g/ml) (Zajmi et al. 2015). A structural analysis using transmission electron microscopy revealed that Art-E disrupted the architecture of the bacterial cell to facilitate the penetration of antibiotics (Zajmi et al., 2015). Despite being structurally distinct from Art-E, a similar type of effect on the bacterial cell wall has been reported with Art-I. Importantly, the membrane destabilizing effects of Art-E are reminiscent of those reported with morusin, which can disrupt the cell membrane architecture and inhibit the phosphatidic acid biosynthesis pathway of *S.*

aureus (Pang et al., 2019). In many ways, morusin and Art-E behave similarly.

The flavonoid Art-F displays little cytotoxic effects against cancer cells but has shown antimycobacterial activity against *Mycobacterium tuberculosis* (MIC = 6.25 μ g/ml) (Namdaung et al. 2006). It has also revealed significant anti-plasmodial activity against the chloroquine-resistant strain FcB1, being slightly less active than Art-B and more active than Art-A (IC_{50} = 4.9, 1.56, 2.20 μ M for Art-A, B, and F, respectively) (Bourjot et al. 2010). Art-F, isolated from *A. integer* var. *silvestris* Corner, has revealed a weak capacity to inhibit 15-lipoxygenase (Shaha et al. 2016). Art-F can be found in different *Artocarpus* species, including a hybrid between *A. heterophyllus* (jackfruit) and *A. integer* (champedak), which has been specifically developed in Thailand to produce tastier and larger fruits, and for disease resistance (Panthong et al. 2013).

The isolation and structural characterization of several artonins have been reported. Still, no specific effect has been pointed out, such as with Art-G and -H (Hano et al. 1990b), Art-J, -K and -L (Aida et al. 1993), and Art-M, -N, O, -P (Hano et al. 1993). Art-O, found in *A. rigida*, has shown a modest anti-proliferative action against HT-29 human colon cancer cells (ED_{50} = 3.2 μ M) (Ren et al. 2010).

Art-I is much more interesting because the compound has revealed activities against multidrug-resistant (MDR) *Staphylococcus aureus* strains. It is a complex molecule (Figure 4) initially isolated from *A. heterophyllus* (Hano et al. 1992) but also from the leaves of *Morus mesozygia* (Fozing et al. 2012). Art-I has been found to inhibit the enzyme phosphodiesterase I *in vitro* (IC_{50} = 15.4 μ M) (Fozing et al. 2012) and function as an efflux pump inhibitor and a generator of ROS in *S. aureus*. Interestingly, the compound revealed an unanticipated capacity to considerably promote the activity of antibiotics (reducing the MIC 500- to 1000 fold) via Art-I-induced cell membrane damages (Farooq et al. 2014). It could be a useful natural product to treat multi-drug-resistant *Staphylococcus* infections. The total synthesis of these complex molecules has been reported recently, opening the door to the design of simpler, more active analogs (Liu et al., 2020).

Art-V has been first isolated from *A. altilis* (Hano et al. 1994). This plant, known as the paparahua tree, is largely used as a traditional medicine in Amazonian Ecuador and other Oceania and South American countries. Many flavones and terpenoids have been isolated from this plant (Luzuriaga-Quichimbo et al., 2019). Art-X is a rare compound isolated from *A. heterophyllus*, analogous to the diarylheptanoid kuwanon R, which has been characterized as an inhibitor of protein tyrosine phosphatase 1B (PTP1B) (Shinomiya et al., 1995; Hoang et al. 2009).

Figure 4. Chemical structures of artonins

ANTICANCER MECHANISM AND MOLECULAR TARGETS OF ARTONIN-E

The mechanism of action of Art-E has been thoroughly investigated, at least at the cellular level. The compound triggers a down-regulation of the anti-apoptotic protein MCL1 (myeloid leukemia cell sequence-1) in human A549 and H292 non-small cell lung cancer cells. It sensitizes them to anoikis (detachment-induced apoptosis) (Wongpankam et al. 2012). In these NSCLC cells, Art-E down-regulates several proteins involved in cell migration and invasion, such as FAK (focal adhesion kinase) and CDC42 (Cell division cycle-42). At non-cytotoxic low concentrations (0.05-05 $\mu\text{g/ml}$), the compound markedly reduced the migration of H460 NSCLC cells *in vitro* (Plaibua et al., 2013). But the main characteristic of Art-E is certainly its capacity to overcome the resistance of cancer cells induced by the ligand TRAIL (tumor necrosis factor-related apoptosis-inducing ligand). In gastric cancer cells, the effect is coupled with an Art-E-induced up-regulation of the death receptor 5 (DR5) and tumor suppressor p53 (Toume et al. 2015). A similar enhancement of TRAIL-induced apoptosis by Art-E has been reported recently with LoVo colorectal cancer cells. The compound down-regulated DR5 while up-regulated the

other major regulator of TRAIL-induced apoptosis called cFLIP (cellular FADD-like-IL-1 β -converting enzyme-inhibitory protein) on LoVo cells (Sophonnithiprasert et al., 2019). Remarkably, Art-E can trigger cell cycle arrest (S phase) and massive mitochondria-dependent apoptosis in SKOV-3 ovarian cancer cells (Rahman et al., 2016). Art-E induced apoptosis in HCT116 colon cancer cells through upregulation of p-ERK1/2 (Nimnuan-ngam et al. 2020).

The capacity of Art-E to overcome TRAIL resistance in cancer cells echoes the effects reported many years ago with Art-B and three other prenylated flavonoids isolated from *A. champeden*. Art-B and the structural analog heterophyllin were found to potently overcome TRAIL resistance in human gastric adenocarcinoma cells and enhance the expression of DR5 (Minakawa et al. 2014). Many other bioactive natural products have been found to suppress TRAIL resistance or enhance TRAIL-induced apoptosis via the death receptor pathway (Dai et al., 2015; Ahmed and Ishibashi, 2016; Shahwar et al., 2019), but the potency of Art-E and Art-B is noticeably high. A similar capacity to induce TRAIL sensitization by regulating DR5 has been observed with morusin in glioblastoma cells (Park et al. 2016), suggesting thus a class effect. TRAIL emerges as a master element of the antitumor action of Art-E and its analogs (Figure 5).

Figure 5. A schematic illustration of the anticancer mechanism of action common to morusine and artonin E (5-hydroxy-morusin). Both compounds trigger an up-regulation of the tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) and a down-regulation of the death receptor 5 (DR5). Upon trimerization of TRAIL and binding to DR5, the complex promotes the sequential recruitment of Fadd (Fas-associated protein with death domain) and caspase-8 (Casp-8) to form the death-inducing signaling complex (DISC). Processing of Bid (a BH3 domain-containing proapoptotic Bcl2 protein family member) by Casp-8 mediates mitochondrial damage. Activation of Casp-8 also directly induced Casp-3 activation, which concurs with the amplification of apoptosis.

Art-E has revealed marked activities against breast cancer cell lines. The compound inhibited the growth of estrogen receptor (ER) positive breast cancer cells MCF7 with an efficacy superior to that of the reference estrogen receptor modulator tamoxifen, at least *in vitro* ($IC_{50} = 3.8$ and $18.9 \mu M$ for Art-E and tamoxifen, at 72 hours, respectively). Molecular modeling predicted that Art-E could interact directly with the ligand-binding domain of hER α . The docking analysis indicated that Art-E could form more stable complexes with hER α compared to its analogs Art-U, -L, -S, and -T, and artoelastin (Etti et al. 2016). The analog isocyclomorusin is predicted to exhibit an even better binding capacity to hER α than Art-E (Fitriah et al., 2018). Art-E induces caspase-dependent apoptosis of MCF7 cells, associated with elevated production of ROS and an up-regulation of the cyclin-dependent protein kinase inhibitor p21 (Etti et al. 2017b). Similar results have been reported using MDA-MB 231 triple-negative breast cancer cells: up-regulation of p21, G2/M cell cycle arrest, and caspase-dependent apoptosis (Etti et al. 2017c). The binding of Art-E to hER α is entirely plausible, but this computational hypothesis has not been validated experimentally. Other targets have been proposed for Art-E, based on *in silico* modelings, such as the inflammatory cytokine-induced ubiquitin-like modifier FAT10, which is known to target hundreds of proteins for degradation by the 26S proteasome. A molecular modeling analysis has predicted that Art-E could bind and inhibit FAT10 implicated in hepatic carcinoma, but here again, there is no experimental validation of this hypothesis (Chaturvedi 2015). In brief, from a mechanical point of view, Art-E can be considered a TRAIL regulator in cancer cells, but more work is needed to better evidence the upstream molecular targets. Proteins such as hER α and FAT10 may provide a receptor for the compound, but more work is needed to validate these molecular modeling proposals.

DISCUSSION

Artocarpus species have been largely investigated to characterize their pharmacological properties, in line with the traditional medicinal uses of these plants. Thus far, most phytochemical studies have been centered around *A. altilis* and *A. heterophyllus* (Nomura et al. 1998; Sikarwar et al. 2014; Mohammed Haleel et al. 2018; Mainasara and Abu Bakar 2019; Buddhisuharto et al. 2021). The species *A. elasticus* has been less investigated. However, this bark-fiber-producing plant is traditionally used to treat inflammatory conditions and design crafts, clothes, ropes, and building materials (Veriyan et al., 2019). The phytochemical analysis indicates that, like its congeners, *A. elasticus* is rich in prenylated flavones, in particular compounds with a C-3 prenyl side chain, sometimes cyclized into a 5 or 6 membered ring, and occasionally with an additional prenyl chain on ring A or B. These types of C-prenylated flavonoids are frequent in *Artocarpus* species (Šmejkal et al. 2014; Molčanová et al. 2019).

The present analysis sheds light on the artonin group of prenylated compounds and points out the prenylflavone

Art-E as an interesting anticancer natural product. Art-E displays a range of pharmacological activities, chiefly antioxidant and anti-inflammatory effects, similar to those reported with the analogous isoprene flavonoid morusin (Choi et al., 2020). It is a potent anticancer agent, inspiring the design and development of novel anticancer agents. The structural analogy between morusin and Art-E deserves further attention. Morusin has been much more studied than Art-E as an anticancer agent but also for its antiviral and antidiabetic effects (Choi et al., 2020; Kim et al., 2021). The *in vitro* and *in vivo* anticancer effects of morusin have been extensively characterized, and the compound appeared promising against cancer stem cells (Zoofishan et al., 2018). We have a lot to learn from morusin to understand the mode of action of Art-E better.

Art-E and related prenylated flavonoids represent interesting anticancer agents, potentially useful for designing novel bioactive compounds. Promising data have been reported with morusin-loaded nanoparticles for treating glioblastoma (Agarwal et al., 2019; Zheng et al., 2021) and with artocarpin (Tzeng et al., 2016). A similar potency could be anticipated with Art-E using nanoparticle delivery systems. This compound deserves further attention.

CONCLUDING REMARK

In conclusion, this review exposes the potential health benefit of *Artocarpus elasticus* Reinw. ex Blume is used as a traditional medicine in Asia and presents the complete series of artonin bioactive natural products for the first time. An extensive analysis of the scientific information led us to underline the value of the lead compound artonin E for the design of novel anticancer agents. The anticancer properties of Art-E have been well-established, demonstrating its capacity to trigger apoptosis of different types of cancer cells. Still, additional research efforts are required to characterize its molecular targets better. Another takeaway message is that Art-E can be considered as a morusin derivative, endowed with similar, if not superior, anticancer properties. This isoprene flavonoid warrants further studies.

FUNDING AND DECLARATION OF COMPETING INTEREST

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The author declares no conflict of interest associated with this publication, and there has been no significant financial support for this work that could have influenced its outcome.

REFERENCES

- Abdullah NI, Ahmad MB, Shameli K 2015. Biosynthesis of silver nanoparticles using *Artocarpus elasticus* stem bark extract. Chem Cent J 9: 61.

- Achmad SA, Hakim EH, Juliawaty LD, Makmur L, Syah YM. 2005. Indonesian rainforest plants – Chemodiversity and bioactivity. *Malaysian J Sci* 24: 7-16.
- Adorisio S, Fierabracci A, Rossetto A, Muscari I, Nardicchi V, Liberati AM, Riccardi C, Van Sung T, Thuy TT, Delfino DV. 2016. Integration of Traditional and Western Medicine in Vietnamese Populations: A Review of Health Perceptions and Therapies. *Nat Prod Commun* 11: 1409-1416.
- Agarwal S, Mohamed MS, Mizuki T, Maekawa T, Sakthi Kumar D. 2019. Chlorotoxin modified morusin-PLGA nanoparticles for targeted glioblastoma therapy. *J Mater Chem B* 7: 5896-5919.
- Ahmed F, Ishibashi M. 2016. Bio-active Natural Products with TRAIL-Resistance Overcoming Activity. *Chem Pharm Bull (Tokyo)* 64: 119-127.
- Aida M, Shinomiya K, Hano Y, Nomura T. 1993. Artonins J, K, and L, three new isoprenylated flavones from the root bark of *Artocarpus heterophyllus* Lamk. *Heterocycles* 36: 575-583.
- Aida M, Shinomiya K, Matsuzawa K, Hano Y, Nomura T. 1994. Artonins Q, R, S, T, and U, Five New Isoprenylated Phenols from the Bark of *Artocarpus heterophyllus* Lamk. *Heterocycles* 37: 847-858.
- Aneklaphakij C, Bunsupa S, Sirichamorn Y, Bongcheewin B, Satitpatipan V. 2020. Taxonomic Notes on the 'Mahat' (*Artocarpus lacucha* and *A. thailandicus*, Moraceae) Species Complex in Thailand. *Plants (Basel)* 9: 391.
- Arung ET, Muladi S, Sukaton E, Shimizu K, Kondo R. 2008. Artocarpin, A Promising Compound as Whitening Agent and Anti-skin Cancer. *J Tropic Wood Sci Technol* 6: 33-36.
- Arung ET, Shimizu K, Kondo R. 2011. *Artocarpus* plants as a potential source of skin whitening agents. *Nat Prod Commun* 6: 1397-1402.
- Ban YJ, Baiseitova A, Nafiah MA, Kim JY, Park KH. 2020. Human neutrophil elastase inhibitory dihydrobenzoxanthones and alkylated flavones from the *Artocarpus elasticus* root barks. *Appl Biol Chem* 63: 63.
- Boonyaketgoston S, Du Y, Valenciano Murillo AL, Cassera MB, Kingston DGI, Trisuwan K. 2020. Flavanones from the Twigs and Barks of *Artocarpus lakoocha* Having Antiplasmodial and Anti-TB Activities. *Chem Pharm Bull (Tokyo)* 68: 671-674.
- Bourjot M, Apel C, Martin MT, Grelhier P, Nguyen VH, Guéritte F, Litaudon M. 2010. Antiplasmodial, antitrypanosomal, and cytotoxic activities of prenylated flavonoids isolated from the stem bark of *Artocarpus styracifolius*. *Planta Med* 76: 1600-1604.
- Buddhisuharto AK, Pramastya H, Insanu M, Fidriani I. 2021. An Updated Review of Phytochemical Compounds and Pharmacology Activities of *Artocarpus* genus. *Biointerface Res Applied Chem* 11: 14898-14905.
- Cao S, Butler MS, Buss AD. 2003. Flavonoids from *Artocarpus lanceifolius*. *Nat Prod Res* 17: 79-81.
- Cerqueira F, Cidade H, van Ufford L, Beukelman C, Kijjoa A, Nascimento MS. 2008. The natural prenylated flavone artelastin is an inhibitor of ROS and NO production. *Int Immunopharmacol* 8: 597-602.
- Cerqueira F, Cordeiro-da-Silva A, Araújo N, Cidade H, Kijjoa A, Nascimento MS. 2003. Inhibition of lymphocyte proliferation by prenylated flavones: artelastin as a potent inhibitor. *Life Sci* 73: 2321-2334.
- Chan EWC, Wong SK, Tangah J, Chan HT. 2018a. Chemistry and Pharmacology of Artocarpin: An Isoprenyl Flavone from *Artocarpus* Species. *Sys Rev Pharm* 9: 58-63.
- Chan STS, Popplewell WL, Bokes HR, McKee TC, Gustafson KR. 2018b. Five New Stilbenes from the Stem Bark of *Artocarpus communis*. *Nat Prod Sci* 24: 266-271.
- Chaturvedi I. 2015. A Molecular Docking study to find Natural Inhibitor Against FAT10 Protein for curing Hepatic Carcinoma. *J Sci* 1: 1-9.
- Chen CC, Huang YL, Ou JC, Lin CF, Pan TM. 1993. Three new prenylflavones from *Artocarpus altilis*. *J Nat Prod* 56: 1594-1597.
- Choi DW, Cho SW, Lee SG, Choi CY. 2020. The Beneficial Effects of Morusin, an Isoprene Flavonoid Isolated from the Root Bark of *Morus*. *Int J Mol Sci* 21: 6541.
- Choumessi AT, Johanns M, Beaufay C, Herent MF, Stroobant V, Vertommen D, Corbet C, Jacobs R, Herinckx G, Steinberg GR, Feron O, Quetin-Leclercq J, Rider MH. 2019. Two isoprenylated flavonoids from *Dorstenia psilurus* activate AMPK, stimulate glucose uptake, inhibit glucose production and lower glycemia. *Biochem J* 476: 3687-3704.
- Cidade HM, Nascimento MS, Pinto MM, Kijjoa A, Silva AM, Herz W. 2001. Artelastocarpin and carpelastofuran, two new flavones, and cytotoxicities of prenyl flavonoids from *Artocarpus elasticus* against three cancer cell lines. *Planta Med* 67: 867-870.
- Dai X, Zhang J, Arfuso F, Chinnathambi A, Zayed ME, Alharbi SA, Kumar AP, Ahn KS, Sethi G. 2015. Targeting TNF-related apoptosis-inducing ligand (TRAIL) receptor by natural products as a potential therapeutic approach for cancer therapy. *Exp Biol Med (Maywood)* 240: 760-773.
- Ee GC, Teo SH, Rahmani M, Lim CK, Lim YM, Go R. 2011. Artomandin, a new xanthone from *Artocarpus kemando* (Moraceae). *Nat Prod Res* 25: 995-1003.
- Ersam T, Achmad SA, Ghisalberty E, Hakim EH, Makmur L, Syah YM. 2002. A New Isoprenylated Chalcone, Artoindonesianin J, from the Root and Tree Bark of *Artocarpus bracteata*. *J Chem Res* 2002: 186-187.
- Etti I, Abdullah R, Hashim NM, Kadir A, Abdul AB, Etti C, Malami I, Waziri P, How CW. 2016. Artonin E and Structural Analogs from *Artocarpus* Species Abrogates Estrogen Receptor Signaling in Breast Cancer. *Molecules* 21: 839.
- Etti IC, Abdullah R, Kadir A, Hashim NM, Yeap SK, Imam MU, Ramli F, Malami I, Lam KL, Etti U, Waziri P, Rahman M. 2017c. The molecular mechanism of the anticancer effect of Artonin E in MDA-MB 231 triple-negative breast cancer cells. *PLoS One* 12: e0182357.
- Etti IC, Abdullah RB, Hashim NM, Arifah K, Yeap SK, Sani D, Ramli F, Malami I, Waziri P. 2017a. Reduction of breast tumor burden in mice by a prenylated flavonoid, Artonin E. *Australian Med J* 10: 681-693.
- Etti IC, Rasheed A, Hashim NM, Abdul AB, Kadir A, Yeap SK, Waziri P, Malami I, Lim KL, Etti CJ. 2017b. Artonin E induces p53-independent G1 cell cycle arrest and apoptosis through ROS-mediated mitochondrial pathway and living suppression in MCF-7 cells. *Drug Des Devel Ther* 11: 865-879.
- Fang SC, Hsu CL, Yen GC. 2008. Anti-inflammatory effects of phenolic compounds isolated from the fruits of *Artocarpus heterophyllus*. *J Agric Food Chem* 56: 4463-4468.
- Farooq S, Wahab AT, Fozing CD, Rahman AU, Choudhary MI. 2014. Artonin I inhibits multidrug resistance in *Staphylococcus aureus* and potentiates the action of inactive antibiotics in vitro. *J Appl Microbiol* 117: 996-1011.
- Fitriah A, Holil K, Syarifah U, Fitriyah F, Utomo DH. 2018. In silico Approach for Revealing the Anti-Breast Cancer and Estrogen Receptor Alpha Inhibitory Activity of *Artocarpus altilis*. *AIP Conference Proceedings* 2021: 070003.
- Fozing CD, Ali Z, Ngadjui BT, Choudhary MI, Kapche GD, Abegaz BM, Khan IA. 2012. Phosphodiesterase I-inhibiting Diels-Alder adducts from the leaves of *Morus mesozygia*. *Planta Med* 78: 154-159.
- Fukai T, Satoh K, Nomura T, Sakagami H. 2003. Antinephritis and radical scavenging activity of prenylflavonoids. *Fitoterapia* 74: 720-724.
- Gupta AK, Rather MA, Kumar Jha A, Shashank A, Singhal S, Sharma M, Pathak U, Sharma D, Mastinu A. 2020. *Artocarpus lakoocha* Roxb. and *Artocarpus heterophyllus* Lam. Flowers: New Sources of Bioactive Compounds. *Plants (Basel)* 9: 1329.
- Hakim A. Diversity of secondary metabolites from Genus *Artocarpus* (Moraceae). 2010. *Nusantara Biosci* 2: 146-156.
- Hakim EH, Achmad SA, Juliawaty LD, Makmur L, Syah YM, Aimi N, Kitajima M, Takayama H, Ghisalberty EL. 2006. Prenylated flavonoids and related compounds of the Indonesian *Artocarpus* (Moraceae). *J Nat Med* 60: 161-184.
- Hakim EH, Afrida A, Eliza E, Achmad SA, Aimi N, Kitajima N, Makmur L, Mujahidin D, Syah YM, Takayama H. 2000. Artoindonesianin-D: senyawa baru bioaktif turunan piranoflavon dan chaplasin tumbuhan *Artocarpus mainayi*. *Proc ITB* 32, 13-19.
- Hakim EH, Aripin A, Achmad SA, Aimi N, Kitajima N, Makmur L, Mujahidin D, Syah YM, Takayama H. 2001. Artoindonesianin-E suatu senyawa baru turunan flavanon dari tumbuhan *Artocarpus champeden*. *Proc ITB* 33: 69-73.
- Hakim EH, Asnizar, Yurnawilis, Aimi N, Kitajima M, Takayama H. 2002b. Artoindonesianin P, a new prenylated flavone with cytotoxic activity from *Artocarpus lanceifolius*. *Fitoterapia* 73: 668-673.
- Hakim EH, Fahriyati A, Kau MS, Achmad SA, Makmur L, Ghisalberty EL, Nomura T. 1999. Artoindonesianins A and B, two new prenylated flavones from the root of *Artocarpus champeden*. *J Nat Prod* 62: 613-615.
- Hakim EH, Ulinuha UZ, Syah YM, Ghisalberty EL. 2002a. Artoindonesianins N and O, new prenylated stilbene and prenylated arylbenzofuran derivatives from *Artocarpus gomezianus*. *Fitoterapia* 73: 597-603.

- Hano Y, Aida M, Nomura T, Ueda S. 1992. A novel way of determining the structure of artonin I, an optically active Diels–Alder type adduct, with the aid of an enzyme system of *Morus alba* cell cultures. *J Chem Soc. Chem Commun* 17: 1177–1178.
- Hano Y, Aida M, Nomura T. 1990a. Two new natural Diels–Alder-type adducts from the root bark of *Artocarpus heterophyllus*. *Heterocycles* 53: 391–395.
- Hano Y, Aida M, Shiina M, Nomura T, Kawai T, Ohe H, Kagei K. 1989. Artonins A and B, Two New Prenylflavones from the Root Bark of *Artocarpus heterophyllus* Lamk. *Heterocycles* 29: 1447–1453.
- Hano Y, Inami R, Nomura T. 1994. Constituents of Moraceae plants. 20. a novel flavone, artonin V, from the root bark of *Artocarpus altilis*. *J Chem Res Syn* 9: 348–349.
- Hano Y, Inami R, Nomura T. 1990b. Components of the Bark of *Artocarpus rigida* Bl. 1. Structures of Two New Isoprenylated Flavones, Artonins G and H. *Heterocycles* 31: 2173–2179.
- Hano Y, Inami R, Nomura T. 1993. Components of the Bark of *Artocarpus rigida* BL.2. Structures of Four New Isoprenylated Flavone Derivatives Artonins M, N, O and P. *Heterocycles* 35: 1341–1350.
- Hano Y, Yamagami Y, Kobayashi M, Isohata R, Nomura T. 1990b. Artonins E and F, two new prenylflavones from the bark of *Artocarpus communis* Forst. *Heterocycles* 31: 877–882.
- Hari A, Revikumar KG, Divya D. 2014. *Artocarpus*: A review of its phytochemistry and pharmacology. *J Pharma Search* 9: 7–12.
- Hoang DM, Ngoc TM, Dat NT, Ha do T, Kim YH, Luong HV, Ahn JS, Bae K. 2009. Protein tyrosine phosphatase 1B inhibitors isolated from *Morus bombycis*. *Bioorg Med Chem Lett* 19: 6759–6761.
- Jagtap UB, Bapat VA. 2010. *Artocarpus*: a review of its traditional uses, phytochemistry and pharmacology. *J Ethnopharmacol* 129: 142–166.
- Jantan I, Mohd Yasin YH, Jamil S, Sirat H, Basar N. 2010. Effect of prenylated flavonoids and chalcones isolated from *Artocarpus* species on platelet aggregation in human whole blood. *J Nat Med* 64: 365–369.
- Jayasinghe UL, Samarakoon TB, Kumarihamy BM, Hara N, Fujimoto Y. 2008. Four new prenylated flavonoids and xanthenes from the root bark of *Artocarpus nobilis*. *Fitoterapia* 79: 37–41.
- Jenis J, Baiseitova A, Yoon SH, Park C, Kim JY, Li ZP, Lee KW, Park KH. 2019. Competitive α -glucosidase inhibitors, dihydrobenzoxanthenes, from the barks of *Artocarpus elasticus*. *J Enzyme Inhib Med Chem* 34: 1623–1632.
- Jin YJ, Lin CC, Lu TM, Li JH, Chen IS, Kuo YH, Ko HH. 2015. Chemical constituents derived from *Artocarpus xanthocarpus* as inhibitors of melanin biosynthesis. *Phytochemistry* 117: 424–435.
- Kijjoa A, Cidade H, Pinto M, Gonzalez MJTG, Anantachoke C, Gedris TE, Herz W. 1996. Prenylflavonoids from *Artocarpus elasticus*. *Phytochemistry* 43: 691–694.
- Kijjoa A, Cidade HM, Gonzalez MJTG. 1998. Further prenylflavonoids from *Artocarpus elasticus*. *Phytochemistry* 47: 875–878.
- Kim TI, Kwon EB, Oh YC, Go Y, Choi JG. 2021. Mori ramulus and its Major Component Morusin Inhibit Herpes Simplex Virus Type 1 Replication and the Virus-Induced Reactive Oxygen Species. *Am J Chin Med* 49: 163–179.
- Ko HH, Lu YH, Yang SZ, Won SJ, Lin CN. 2005. Cytotoxic prenylflavonoids from *Artocarpus elasticus*. *J Nat Prod* 68: 1692–1695.
- Kuete V, Ango PY, Fotso GW, Kapche GD, Dzoyem JP, Wouking AG, Ngadjui BT, Abegaz BM. 2011. Antimicrobial activities of the methanol extract and compounds from *Artocarpus communis* (Moraceae). *BMC Complement Altern Med* 11: 42.
- Kuete V, Mbaveng AT, Zeino M, Fozing CD, Ngameni B, Kapche GD, Ngadjui BT, Efferth T. 2015. Cytotoxicity of three naturally occurring flavonoid-derived compounds (artocarpesin, cycloartocarpesin and isobavachalcone) towards multi-factorial drug-resistant cancer cells. *Phytomedicine* 22: 1096–102.
- Kuran RY, Ersam T. 2017. Prenylated Flavone of the Bark of *Artocarpus elasticus* from Alor Island-NTT Indonesia. *J Applied Chem Sci* 4: 322–324.
- Lang L, Dong N, Wu D, Yao X, Lu W, Zhang C, Ouyang P, Zhu J, Tang Y, Wang W, Li J, Huang J. 2016. 2-Arylbenzo[b]furan derivatives as potent human lipoxygenase inhibitors. *J Enzyme Inhib Med Chem* 31: 98–105.
- Lee CC, Lin CN, Jow GM. 2006. Cytotoxic and apoptotic effects of prenylflavonoid artonin B in human acute lymphoblastic leukemia cells. *Acta Pharmacol Sin* 27: 1165–1174.
- Lin CN, Shieh WL, Ko FN, Teng CM. 1993. Antiplatelet activity of some prenylflavonoids. *Biochem Pharmacol* 45: 509–5012.
- Lin CN, Shieh WL. 1992. Pyranoflavonoids from *Artocarpus communis*. *Phytochemistry* 31: 2922–2924.
- Lin KW, Liu CH, Tu HY, Ko HH, Wei BL. 2009. Antioxidant prenylflavonoids from *Artocarpus communis* and *Artocarpus elasticus*. *Food Chem* 115: 558–562.
- Liu X, Yang J, Gao L, Zhang L, Lei X. 2020. Chemoenzymatic Total Syntheses of Artonin I with an Intermolecular Diels–Alderase. *Biotechnol J* 15: e2000119.
- Losuwanarak N, Sritularak B, Chanvorachote P. 2018. Cycloartobioxanthone Induces Human Lung Cancer Cell Apoptosis via Mitochondria-dependent Apoptotic Pathway. *In Vivo* 32: 71–78.
- Luzuriaga-Quichimbo CX, Blanco-Salas J, Cerón-Martínez CE, Ruiz-Téllez T. 2019. Providing added value to local uses of paparahua (*Artocarpus altilis*) in Amazonian Ecuador by phytochemical data review. *Revista Brasileira de Farmacognosia* 29: 62–68.
- Ma JP, Qiao X, Pan S, Shen H, Zhu GF, Hou AJ. 2010. New isoprenylated flavonoids and cytotoxic constituents from *Artocarpus tonkinensis*. *J Asian Nat Prod Res* 12: 586–592.
- Mainasara MM, Abu Bakar MF. 2019. Phytochemical constituents, antioxidant and antiproliferative properties of *Artocarpus altilis* (sukun) from Endau Rompin, Johor, Malaysia. *Sci World J* 14: 124–135.
- Makmur L, Syamsurizal S, Tukiran T, Achmad SA, Aimi N, Hakim EH, Kitajima M, Takayama H. 2000. Artoindonesianin C, a new xanthone derivative from *Artocarpus teysmanii*. *J Nat Prod* 63: 243–244.
- Minakawa T, Toume K, Arai MA, Koyano T, Kowithyakorn T, Ishibashi M. 2013. Prenylflavonoids isolated from *Artocarpus champeden* with TRAIL-resistance overcoming activity. *Phytochemistry* 96: 299–304. Corrigendum in *Phytochemistry* 2014, 106: 207.
- Mohammed Haleel PM, Rashid K, Senthil Kumar S. 2018. *Artocarpus heterophyllus*: Review Study on Potential Activities. *Res J Pharmacol Pharmacodyn* 10: 24–28.
- Molčanová L, Janošíková D, Dall’Acqua S, Smejkal KS. 2019. C-prenylated flavonoids with potential cytotoxic activity against solid tumor cell lines. *Phytochem Rev* 18: 1051–1100.
- Morrison JJ, Zhang J, Lin J, Murray JE, Porter R, Langat MK, Sadgrove NJ, Barker J, Zhang G, Delgoda R. 2021. Potential chemopreventive, anticancer and anti-inflammatory properties of a refined artocarpin-rich wood extract of *Artocarpus heterophyllus* Lam. *Sci Rep* 11: 6854.
- Musthapa I, Hakim EH, Juliawaty LD, Syah YM, Achmad SA. 2010. Prenylated flavones from some Indonesian *Artocarpus* and their antimalarial properties. *Med Plant* 2: 157–160.
- Musthapa I, Juliawaty LD, Syah YM, Hakim EH, Latip J, Ghisalberti EL. 2009a. An oxepinoflavone from *Artocarpus elasticus* with cytotoxic activity against P-388 cells. *Arch Pharm Res* 32: 191–194.
- Musthapa I, Latip J, Takayama H, Juliawaty LD, Hakim EH, Syah YM. 2009b. Prenylated flavones from *Artocarpus lanceifolius* and their cytotoxic properties against P-388 cells. *Nat Prod Commun* 4: 927–930.
- Namdaung U, Aroonrerk N, Suksamrarn S, Danwisetkanjana K, Saenboonrueng J, Arjchomphu W, Suksamrarn A. 2006. Bioactive constituents of the root bark of *Artocarpus rigidus* subsp. *rigidus*. *Chem Pharm Bull (Tokyo)* 54: 1433–1436.
- Nimnuan-ngam S, Yangnok K, Innajak S, Watanapokasin R. 2020. Effect of artonin E on apoptosis cell death induction in colon cancer cells. *Cancer Epidemiol Biomarker Prev* 29, B061.
- Nomura T, Hano Y, Aida M. 1998. Isoprenoid-substituted flavonoids from *Artocarpus* plants (Moraceae). *Heterocycles* 47: 1179–1205.
- Noraini J. Phytochemicals and bioactivities of *Artocarpus anisophyllus* MIQ. 2013. Masters thesis, Universiti Teknologi Malaysia, Faculty of Science (<http://eprints.utm.my/id/eprint/33249/>).
- Pang D, Liao S, Wang W, Mu L, Li E, Shen W, Liu F, Zou Y. 2019. Destruction of the cell membrane and inhibition of cell phosphatidic acid biosynthesis in *Staphylococcus aureus*: an explanation for the antibacterial mechanism of morusin. *Food Funct* 10: 6438–6446.
- Panthong K, Tohdee K, Towatana NH, Voravuthikunchaib SP, Chusrib S. 2013. Prenylated Flavone from Roots of a Hybrid between *Artocarpus heterophyllus* and *Artocarpus integer* and its Biological Activities. *J Braz Chem Soc* 24: 1656–1661.
- Park D, Ha JJ, Park SY, Choi M, Lim SL, Kim SH, Lee JH, Ahn KS, Yun M, Lee SG. 2016. Morusin Induces TRAIL Sensitization by

- regulating EGFR and DR5 in Human Glioblastoma Cells. *J Nat Prod* 79: 317-323.
- Parvatikar PP, Madagi SB. 2021. Molecular Docking Analysis: Interaction Studies of Natural Compounds with Human TG2 Protein. In S.-I. Ao et al. (Eds.): WCECS 2018, *Transactions on Engineering Technologies*. Springer Nature Singapore Pte Ltd., Singapore
- Pedro M, Ferreira MM, Cidade H, Kijjoa A, Bronze-da-Rocha E, Nascimento MS. 2005. Artelastin is a cytotoxic prenylated flavone that disturbs microtubules and interferes with DNA replication in MCF-7 human breast cancer cells. *Life Sci* 77: 293-311.
- Plaibua K, Pongrakhananon V, Chunhacha P, Sritularak B, Chanvorachote P. 2013. Effects of artonin E on migration and invasion capabilities of human lung cancer cells. *Anticancer Res* 33: 3079-3088.
- Qiao X, Zhao T, Wang M, Lei C, Hou A. 2011. [Flavonoids from *Artocarpus hypargyreus*]. *Zhongguo Zhong Yao Za Zhi* 36: 2975-2979.
- Rahman MA, Ramli F, Karimian H, Dehghan F, Nordin N, Ali HM, Mohan S, Hashim NM. 2016. Artonin E Induces Apoptosis via Mitochondrial Dysregulation in SKOV-3 Ovarian Cancer Cells. *PLoS One* 11: e0151466.
- Ramli F, Rahmani M, Ismail IS, Sukari MA, Abd Rahman M, Zajmi A, Akim AM, Hashim NM, Go R. 2016. A New Bioactive Secondary Metabolite from *Artocarpus elasticus*. *Nat Prod Commun* 11: 1103-1106.
- Ranasinghe RASN, Maduwanthi SDT, Marapana RAUJ. 2019. Nutritional and Health Benefits of Jackfruit (*Artocarpus heterophyllus* Lam.): A Review. *Int J Food Sci* 2019: 4327183.
- Rao GV, Mukhopadhyay T, Radhakrishnan N. 2010. Artoindonesianin F, a potent tyrosinase inhibitor from the roots of *Artocarpus heterophyllus* Lam. *Indian J Chem* 49B: 1264-1266.
- Reddy GR, Ueda N, Hada T, Sackeyfio AC, Yamamoto S, Hano Y, Aida M, Nomura T. 1991. A prenylflavone, artonin E, as arachidonate 5-lipoxygenase inhibitor. *Biochem Pharmacol* 41: 115-118.
- Ren Y, Kardono LB, Riswan S, Chai H, Farnsworth NR, Soejarto DD, Carcache de Blanco EJ, Kinghorn AD. 2010. Cytotoxic and NF-kappaB inhibitory constituents of *Artocarpus rigida*. *J Nat Prod* 73: 949-955.
- Sahu SK, Liu M, Yssel A, Kariba R, Muthemba S, Jiang S, Song B, Hendre PS, Muchugi A, Jamnadass R, Kao SM, Featherston J, Zerega NJC, Xu X, Yang H, Deynze AV, Peer YV, Liu X, Liu H. 2019. Draft Genomes of Two *Artocarpus* Plants, Jackfruit (*A. heterophyllus*) and Breadfruit (*A. altilis*). *Genes (Basel)* 11: 27.
- Seo EK, Lee D, Shin YG, Chai HB, Navarro HA, Kardono LB, Rahman I, Cordell GA, Farnsworth NR, Pezzuto JM, Kinghorn AD, Wani MC, Wall ME. 2003. Bioactive prenylated flavonoids from the stem bark of *Artocarpus kemo*. *Arch Pharm Res* 26: 124-127.
- Shaha MKK, Sirata HM, Jamil S, Jalil J. 2016. Flavonoids from the Bark of *Artocarpus integer* var. *silvestris* and their Anti-inflammatory Properties. *Nat Prod Commun* 11: 1275-1278.
- Shahwar D, Iqbal MJ, Nisa MU, Todorovska M, Attar R, Sabitaliyevich UY, Farooqi AA, Ahmad A, Xu B. 2019. Natural Product Mediated Regulation of Death Receptors and Intracellular Machinery: Fresh from the Pipeline about TRAIL-Mediated Signaling and Natural TRAIL Sensitizers. *Int J Mol Sci* 20: 2010.
- Shamaun SS, Rahmani M, Hashim NM, Ismail HB, Sukari MA, Lian GE, Go R. 2010. Prenylated flavones from *Artocarpus altilis*. *J Nat Med* 64: 478-81.
- Shinomiya K, Aida M, Hano Y, Nomura T. 1995. A diels-alder-type adduct from *Artocarpus heterophyllus*. *Phytochemistry* 40: 1317-1319.
- Shinomiya K, Hano Y, Nomura T. 2000. Mechanism on one-sided Wessely-Moser rearrangement reaction. *Heterocycles* 53: 877-886.
- Shobana S, Veena S, Sameer SSM, Swarnalakshmi K, Vishal LA. 2020. Green Synthesis of Silver Nanoparticles Using *Artocarpus hirsutus* Seed Extract and its Antibacterial Activity. *Curr Pharm Biotechnol* 21: 980-989.
- Sikarwar MS, Hui BJ, Subramaniam K, Valeisamy BD, Yean LK, Balaji K. 2014. A Review on *Artocarpus altilis* (Parkinson) Fosberg (breadfruit). *J Applied Pharm Sci* 4: 91-97.
- Šmejkal K. 2014. Cytotoxic potential of C-prenylated flavonoids. *Phytochem Rev* 13: 245-275.
- Soekanto NH, Achmad SA, Ghisalberti EL, Hakim EH, Syah YM. 2003. Artoindonesianins X and Y, two isoprenylated 2-arylbenzofurans, from *Artocarpus fretessi* (Moraceae). *Phytochemistry* 64: 831-834.
- Sophonnihiprasert T, Mahabusarakam W, Watanapokasin R. 2019. Artonin E sensitizes TRAIL-induced apoptosis by DR5 upregulation and cFLIP downregulation in TRAIL-refractory colorectal cancer LoVo cells. *J Gastrointest Oncol* 10: 209-217.
- Sritularak B, Tantituvanont A, Chanvorachote P, Meksawan K, Miyamoto T, Kohno Y, Likhitwitayawuid K. 2010. Flavonoids with free radical scavenging activity and nitric oxide inhibitory effect from the stem bark of *Artocarpus gomezianus*. *J Med Plants Res* 4: 387-392.
- Suhartati T, Achmad SA, Aimi N, Hakim EH, Kitajima M, Takayama H, Takeya K. 2001. Artoindonesianin L, a new prenylated flavone with cytotoxic activity from *Artocarpus rotunda*. *Fitoterapia* 72: 912-918.
- Suhartati T, Hernawan H, Suwandi JF, Yandri Y, Hadi S. 2018. Isolation of Artonin E from the root bark of *Artocarpus rigida*, synthesis of Artonin E acetate and evaluation of anticancer activity. *Macedonian J Chem & Chem Engineer* 37: 35-42.
- Suhartati T, Wulandari A, Suwandi JF, Yandri Y, Hadi S. 2016. Artonin O, a Xanthone Compound from Root Wood of *Artocarpus rigida*. *Orient J Chem* 32: 2777-2784.
- Suhartati T, Yandri Y, Hadi S. 2008. The Bioactivity Test of Artonin E from the Bark of *Artocarpus Rigida* Blume. *Eur J Sci Res* 23: 330-337.
- Susiarti S, Rahayu M, Kuncari ES, Astuti IP. 2020. Utilization of "Benda" (*Artocarpus elasticus* Reinw. ex Blume) in Bogor, West Java, Indonesia: An Ethnobotanical Case Study. *J Trop Biol Conserv* 17: 297-307.
- Syah YM, Achmad SA, Ghisalberti EL, Hakim EH, Makmur L, Mujahidin D. 2001. Artoindonesianins G-I, three new isoprenylated flavones from *Artocarpus lanceifolius*. *Fitoterapia* 72: 765-73.
- Syah YM, Achmad SA, Ghisalberti EL, Hakim EH, Makmur L, Mujahidin D. 2002b. Artoindonesianin M, a new prenylated flavone from *Artocarpus champeden*. *Bull Soc Nat Prod Chem (Indonesia)* 2: 31-36.
- Syah YM, Achmad SA, Ghisalberti EL, Hakim EH, Makmur L, Mujahidin D. 2002a. Artoindonesianins Q-T, four isoprenylated flavones from *Artocarpus champeden* Spreng. (Moraceae). *Phytochemistry* 61: 949-953.
- Syah YM, Achmad SA, Ghisalberti EL, Hakim EH, Mujahidin D. 2004. Two new cytotoxic isoprenylated flavones, artoindonesianins U and V, from the heartwood of *Artocarpus champeden*. *Fitoterapia* 75: 134-140.
- Syah YM, Juliawaty LD, Achmad SA, Hakim EH, Ghisalberti E. 2006a. Cytotoxic prenylated flavones from *Artocarpus champeden*. *J Nat Med* 60: 308-312.
- Syah YM, Hakim EH, Makmur L, Kurdi VA, Ghisalberti E, Aimi N, Achmad SA. 2006b. Prenylated 2-arylbenzofurans from two Species of *Artocarpus*. *Nat Product Commun* 1: 549-552.
- Teo SP, Nasution RE. 2003. *Artocarpus elasticus* Reinw. ex Blume. In: Brink M, Escobin RP. (eds.). *Plant Resources of South-East Asia No 17: Fibre plants*. PROSEA Foundation, Bogor, Indonesia. prota4u.org/prosea
- Toume K, Habu T, Arai MA, Koyano T, Kowithayakorn T, Ishibashi M. 2015. Prenylated flavonoids and resveratrol derivatives isolated from *Artocarpus communis* with the ability to overcome TRAIL resistance. *J Nat Prod* 78: 103-110. Erratum in *J Nat Prod* 78: 968.
- Tungsukruthai S, Sritularak B, Chanvorachote P. 2017. Cycloartobioxanthone Inhibits Migration and Invasion of Lung Cancer Cells. *Anticancer Res* 37: 6311-6319.
- Tzeng CW, Tzeng WS, Lin LT, Lee CW, Yen FL, Lin CC. 2016. Enhanced autophagic activity of artocarpin in human hepatocellular carcinoma cells through improving its solubility by a nanoparticle system. *Phytomedicine* 23: 528-540.
- Veitch NC, Grayer RJ. 2008. Flavonoids and their glycosides, including anthocyanins. *Nat Prod Rep* 25: 555-611.
- Veriyan A, Raffidinal R, Linda R. 2019. Kajian Etnobotani Serat Kulit Kayu Kepauk (*Artocarpus elasticus* Reinw. ex Blume) Pada Suku Dayak Desa'di Desa Kunyai Kecamatan Sungai Tebelian Kabupaten Sintang. *Protobiont* 8: 41-45.
- Wang JP, Tsao LT, Raung SL, Lin PL, Lin CN. 1999. Stimulation of respiratory burst by cyclocommunin in rat neutrophils is associated with the increase in cellular Ca²⁺ and protein kinase C activity. *Free Radic Biol Med* 26: 580-588.
- Wang YH, Hou AJ, Chen L, Chen DF, Sun HD, Zhao QS, Bastow KF, Nakanish Y, Wang XH, Lee KH. 2004. New isoprenylated flavones, artochamins A-E, and cytotoxic principles from *Artocarpus chama*. *J Nat Prod* 67: 757-761.
- Wei BL, Weng JR, Chiu PH, Hung CF, Wang JP, Lin CN. 2005. Anti-inflammatory flavonoids from *Artocarpus heterophyllus* and *Artocarpus communis*. *J Agric Food Chem* 53: 3867-3871.

- Widyawaruyanti A, Subehan S, Kalauni SK, Awale S, Nindatu M, Zaini NC, Syafruddin D, Asih PBS, Tezuka Y, Kadota S. 2007. New prenylated flavones from *Artocarpus champeden*, and their antimalarial activity in vitro. *J Nat Med* 61: 410-413.
- Williams EW, Gardner EM, Harris R 3rd, Chaveerach A, Pereira JT, Zerega NJ. 2017. Out of Borneo: biogeography, phylogeny and divergence date estimates of *Artocarpus* (Moraceae). *Ann Bot* 119: 611-627.
- Wongpankam E, Chunhacha P, Pongrakhananon V, Sritularak B, Chanvorachote P. 2012. Artonin E mediates MCL1 down-regulation and sensitizes lung cancer cells to anoikis. *Anticancer Res* 32: 5343-5351.
- Ye JB, Ren G, Li WY, Zhong GY, Zhang M, Yuan JB, Lu T. 2019. Characterization and Identification of Prenylated Flavonoids from *Artocarpus heterophyllus* Lam. Roots by Quadrupole Time-Of-Flight and Linear Trap Quadrupole Orbitrap Mass Spectrometry. *Molecules* 24: 4591.
- Zajmi A, Mohd Hashim N, Noordin MI, Khalifa SA, Ramli F, Mohd Ali H, El-Seedi HR. 2015. Ultrastructural Study on the Antibacterial Activity of Artonin E versus Streptomycin against *Staphylococcus aureus* Strains. *PLoS One* 10: e0128157.
- Zerega NJC, Nur Supardi MN, Motley TJ. 2010. Phylogeny and Recircumscription of Artocarpeae (Moraceae) with a Focus on *Artocarpus*. *Systematic Bot* 35: 766-782.
- Zhai XX, Xiao CY, Jiang W, Chen YL, Ding YQ, Ren G. 2017. Isoprenylated Phenolics from Roots of *Artocarpus heterophyllus*. *Nat Prod Commun* 12: 921-924.
- Zheng R, Zhao L, Chen X, Liu L, Liu Y, Chen X, Wang C, Yu X, Cheng H, Li S. 2021. Metal-coordinated nanomedicine for combined tumor therapy by inducing paraptosis and apoptosis. *J Control Release* 336: 159-168.
- Zheng ZP, Chen S, Wang S, Wang XC, Cheng KW, Wu JJ, Yang D, Wang M. 2009. Chemical components and tyrosinase inhibitors from the twigs of *Artocarpus heterophyllus*. *J Agric Food Chem* 57: 6649-6655.
- Zheng ZP, Cheng KW, To JT, Li H, Wang M. 2008. Isolation of tyrosinase inhibitors from *Artocarpus heterophyllus* and use of its extract as anti-browning agent. *Mol Nutr Food Res* 52: 1530-1538.
- Zoofishan Z, Hohmann J, Hunyadi A. 2018. Phenolic antioxidants of *Morus nigra* roots, and antitumor potential of morusin. *Phytochemistry Rev* 17: 1031-1045.