

## Short communication: An overview of the trematodes fauna of pool frog *Pelophylax lessonae* (Camerano, 1882) in the Volga Basin, Russia: 1. Adult stages

IGOR V. CHIKHLYAEV<sup>1</sup>, ALEXANDER B. RUCHIN<sup>2,\*</sup>, ALEXANDER I. FAYZULIN<sup>1</sup>

<sup>1</sup> Institute of Ecology of the Volga Basin, Russian Academy of Sciences, Togliatti, Russia

<sup>2</sup> Joint Directorate of the Mordovia State Nature Reserve and National Park "Smolny", Saransk, Russia  
\*email: sasha\_ruchin@rambler.ru, ruchin\_mgpz@mail.ru. Tel: +50-8001010010; Fax: +50-7001010062.

Manuscript received: 16 November 2018. Revision accepted: 27 November 2018.

**Abstract.** Chikhlyayev IV, Ruchin AB, Fayzulin AI. 2018. Short communication: An overview of the trematodes fauna of pool frog *Pelophylax lessonae* (Camerano, 1882) in the Volga Basin, Russia: 1. Adult stages. *Nusantara Bioscience* 10: 256-262. The paper presents data on fauna of trematodes of a pool frog *Pelophylax lessonae* (Camerano, 1882) from 13 regions of the Volga basin (Russia). It consolidates data from different authors over the past 80 years, supplemented by our own research results. There are authentically known findings of 19 trematodes species at an adult stage of development. Three species of trematodes make the basis of helminth fauna: *Pneumonoeces variegatus*, *Opisthioglyphe ranae* and *Diplodiscus subclavatus*. By all species of helminths the following data are provided: taxonomic position, localization, area of detection, biology, definitive hosts, geographic distribution and the degree host-specificity.

**Keywords:** Adult stages, *Pelophylax lessonae*, pool frog, trematodes, Volga basin

### INTRODUCTION

The pool frog *Pelophylax lessonae* (Camerano, 1882) (Amphibia: Anura) is a common in Europe from Southern France in the west to Tatarstan and Bashkortostan (Russia) in the east (Borkin et al. 1987; Hotz et al. 2008; Hofman et al. 2012; Dedukh et al. 2015; Fayzulin et al. 2018; Zeisset, Hoogesteger 2018). It inhabits deciduous and mixed forests, where it prefers shallow, overgrown standing reservoirs: lakes, ponds, oxbow lakes, swamps; it rarely occurs in shallow waters of rivers and streams, in floodplains and islands. The pool frog penetrates into the forest-steppe and steppe zones passing thickets of riverine shrubs and afforested floodplains of rivers. It is common in anthropogenic landscapes, where it inhabits temporary and drying out reservoirs: ruts, roadside pits, ditches, and quarries with water. Pool frogs are carnivorous animals, eating mainly different classes of terrestrial and aquatic invertebrates (insects, spiders, centipedes, crustaceans, gastropods, oligochaetes); rarely - vertebrates (fish fry, tadpoles and amphibian yearlings) animals (Kuzmin 2012). A wide range of food, intra- and inter-species cannibalism affect the pool frog infected by various species of the trematodes at the adult stage of development (marita). The trematodes are known to develop with an obligate change of hosts and therefore can serve as bioindicators of the trophic relationships of amphibians in the biocoenosis.

This paper continues a series of publications devoted to the modern characterization of the helminth fauna of amphibians in the Volga basin (Ruchin et al. 2009, 2016; Reshetnikov et al. 2013; Chikhlyayev, Ruchin 2014;

Chikhlyayev et al. 2016b, 2018). This study aimed to present a review on fauna of adult stages of trematodes of a pool frog from Volga basin based on our researches and literature.

### MATERIALS AND METHODS

In the period from 1936 to 2016, many authors had examined 1,460 specimens of pool frog from 13 regions of the Volga basin, Russia Federation among which are: Kaluga, Moscow, Ivanovo, Kostroma, Nizhny Novgorod, Tambov, Ulyanovsk and Samara regions, the Republics of Mordovia, Chuvashia, Mari El, Tatarstan and Bashkortostan. For species determination of trematodes, we used reports of K.M. Ryzhikov et al. (1980). At distribution on taxons adhered to the modern data on a systematics of trematodes (Keys to the Trematoda, 2002, 2008) and given the website "Fauna Europaea" (<http://www.fauna-eu.org>).

### RESULTS AND DISCUSSION

In total in the pool frog on the territory of Volga basin there are registered 19 species of trematodes at an adult stage of development, belonging to 13 genera, 7 families and 3 orders (Table 1). Of these, 11 species are broadly specific, polyhostal parasites of anurans and 8 are specific, oligohostal for Ranidae family. Among all trematodes, 15 species use frogs as obligatory definitive (final) hosts.

Other 4 species (*G. vitelliloba*, *H. cylindracea*, *D. rastellus*, *O. ranae*) combine different stages of development in one individual or individuals of different ages, and use amphibians as amphyxenic, and the latter two species, also as post-cyclic hosts.

The annotated list of pool frog trematodes species with an indication of their systematic position, localization, areas of detection, biology, degree of parasites host-specificity and geographic distribution is given below. Also, the list of their definitive hosts in Russia is indicated, corrected according to literary analysis (Ryzhikov et al. 1980; Kostyunin 2010; Chikhlyayev et al. 2012a, b; Kirillov et al. 2012, 2018; Kuzmin 2012). The sign “?” marked the single finds of helminths at final hosts demanding confirmation.

Class: Trematoda Rudolphi, 1808  
Order: Hemiurida Skrjabin et Guschanskaja, 1956  
Family: Derogenidae Nicoll, 1910

*Halipegus ovocaudatus* (Vulpian, 1859)

Localization: oral cavity.

Areas of detection: Moscow, Nizhny Novgorod and Samara regions, the Republic of Tatarstan.

Biology: Widely specific parasite of anurans. Tetraxenic life cycle. Intermediate host - gastropod mollusks genus *Planorbis*; first additional hosts - cyclops genus *Macrocyclus*; second additional hosts - dragonflies of different genera (Kechemir 1976).

Definitive hosts: *Pelophylax ridibundus* (Pallas, 1771), *Pelophylax lessonae* (Camerano, 1882), *Rana arvalis* Nilsson, 1842, *Rana temporaria* Linnaeus, 1758, *Bombina bombina* (Linnaeus, 1761), *Triturus cristatus* (Laurenti, 1768) (rarely) (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Europe.

Order: Paramphistomida Skrjabin et Schulz, 1937  
Family: Diplodiscidae Cohn, 1904

*Diplodiscus subclavatus* (Pallas, 1760)

Localization: rectum, small intestine.

Areas of detection: Moscow, Ivanovo, Nizhny Novgorod, Tambov and Samara regions, the Republics of Mordovia, Chuvashia, Mari El, Tatarstan and Bashkortostan.

Biology: Widely specific parasite of amphibians. Dixenic life cycle. Intermediate hosts - gastropods genus *Planorbis*; less often - genera *Anisus*, *Viviparus* and *Segmentina*. Additional hosts are absent (Skryabin 1949).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *Pelophylax esculentus* (Linnaeus, 1758), *Pelophylax nigromaculatus* (Hallowell, 1861), *R. arvalis*, *R. temporaria*, *Rana macrocnemis* Boulenger, 1885, *Rana amurensis* Boulenger, 1886, *Rana dybowskii* Guenther, 1876, *B. bombina*, *Pelobates fuscus* (Laurenti, 1768), *Hyla arborea* (Linnaeus, 1758), *Hyla orientalis* Bedriaga, 1890, *Bufo bufo* (Linnaeus, 1758), *Bufo viridis* (Laurenti, 1768), *Lissotriton vulgaris* (Linnaeus, 1758), *T. cristatus*

(Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: cosmopolite.

Order: Plagiorchiida La Rue, 1957  
Family: Gorgoderidae Looss, 1899

*Gorgodera cygnoides* (Zeder, 1800)

Localization: bladder.

Areas of detection: Ivanovo and Nizhny Novgorod regions, the Republics of Mordovia, Tatarstan and Bashkortostan.

Biology: Widely specific parasite of anurans. Trixenic life cycle. Intermediate hosts - bivalve molluscs of genera *Cyclas*, *Pisidium* and *Sphaerium*; additional hosts - dragonfly of different species, cyclops genus *Mesocyclops* (Pigulevsky 1952).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *R. arvalis*, *R. temporaria*, *R. macrocnemis*, *B. bombina*, *P. fuscus*, *Bufo calamita* Laurenti, 1768, *B. viridis*, *H. arborea*, *H. orientalis* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

*Gorgodera loossi* Sinitzin, 1905

Localization: bladder.

Area of detection: Republic of Bashkortostan.

Biology: Specific parasite of water frogs. Trixenic life cycle. Intermediate hosts - bivalve mollusks genus *Sphaerium*; additional hosts - dragonflies of genera *Agriion* and *Epitheca* (Pigulevsky 1952).

Definitive hosts: *P. ridibundus*, *P. lessonae* and *R. temporaria* (?) (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Europe.

*Gorgodera microovata* Fuhrmann, 1924

Localization: bladder.

Areas of detection: Nizhny Novgorod region, the Republics of Mordovia and Tatarstan.

Biology: Specific parasite of frogs. The life cycle has not been studied.

Definitive hosts: *P. ridibundus*, *P. lessonae*, *P. esculentus*, *R. arvalis*, *R. temporaria*, *Rana asiatica* Bedriaga, 1898 (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Europe.

*Gorgodera pagenstecheri* Sinitzin, 1905

Localization: bladder.

Areas of detection: Nizhny Novgorod and Samara regions, the Republics of Mari El and Tatarstan.

Biology: Specific parasite of frogs. Trixenic life cycle. Intermediate hosts - bivalve mollusks genera *Sphaerium* and *Pisidium*; additional hosts - dragonfly of different species, caddisflies genus *Limnophilus* (Pigulevsky 1952).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *R. arvalis*, *R. temporaria*, *R. macrocnemis* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

*Gorgodera varsoviensis* Sinitzin, 1905

Localization: bladder.

Areas of detection: Moscow, Kostroma, Nizhny Novgorod and Samara regions, the Republic of Tatarstan.

Biology: Specific parasite of frogs. Trixenic life cycle. Intermediate host - bivalve mollusks genus *Sphaerium*; additional hosts - dragonfly of different species, caddisflies genus *Limnophilus* (Pigulevsky 1952).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *P. esculentus*, *R. arvalis*, *R. temporaria* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Europe.

*Gorgoderina vitelliloba* (Olsson, 1876)

Localization: bladder.

Areas of detection: Nizhny Novgorod and Samara regions.

Biology: Widely specific parasite of anurans. Trixenic life cycle. Intermediate hosts - bivalve mollusks genera *Sphaerium*, *Pisidium* and *Musculium*; additional hosts - the tadpoles, alderflies genus *Sialis* (Pigulevsky 1953).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *P. esculentus*, *R. arvalis*, *R. temporaria*, *R. macrocnemis*, *B. bombina* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

Family: Plagiorchiidae Luhe, 1901

*Haplometra cylindracea* (Zeder, 1800)

Localization: lungs.

Areas of detection: Moscow, Ivanovo, Kostroma and Nizhny Novgorod regions.

Biology: Specific parasite of frogs. Trixenic life cycle. Intermediate hosts - gastropod mollusks genus *Lymnaea*; additional hosts - tadpoles (Grabda-Kazubskaya 1970; Dobrovolsky, Raikhel 1973).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *R. arvalis*, *R. temporaria*, *R. macrocnemis*, *R. amurensis*, *B. bombina* (?), *B. viridis* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

Family: Telorchidae Looss, 1899

*Dolichosaccus rastellus* (Olsson, 1876)

Localization: small intestine.

Area of detection: Kostroma region.

Biology: Specific parasite of frogs. Trixenic life cycle. Intermediate hosts - gastropod mollusks genus *Lymnaea*; additional hosts - tadpoles of frogs (Grabda-Kazubskaya 1969; Kalabekov 1976).

Definitive hosts: *P. ridibundus* (?), *P. lessonae* (?), *R. arvalis*, *R. temporaria*, *R. macrocnemis*, *R. asiatica*, *R. amurensis*, *R. dybowskii* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

*Opisthioglyphe ranae* (Frohlich, 1791)

Localization: small intestine.

Areas of detection: Kaluga, Moscow, Ivanovo, Kostroma, Nizhny Novgorod, Tambov and Samara regions, the Republics of Mordovia, Chuvashia and Tatarstan.

Biology: Widely specific parasite of amphibians. Trixenic life cycle. Intermediate hosts - gastropod mollusks genus *Lymnaea*; additional hosts - gastropods family Lymnaeidae and juvenile of amphibians (Dobrovolsky 1965a; Grabda-Kazubskaya 1969).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *P. esculentus*, *R. arvalis*, *R. temporaria*, *R. macrocnemis*, *R. amurensis*, *B. bombina*, *P. fuscus*, *H. arborea*, *H. orientalis*, *B. bufo*, *B. viridis*, *T. cristatus*, *L. vulgaris*, *Lissotriton montandoni* (Boulenger, 1880), *Salamandrella keyserlingii* Dybowski, 1870 (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

Family: Haematoloecidae Freitas et Lent, 1939

*Pneumonoeces variegatus* (Rudolphi, 1819)

Syn.: *Haematoloechus variegatus* (Rudolphi, 1819)

Localization: lungs.

Areas of detection: Kaluga, Moscow, Ivanovo, Kostroma, Nizhny Novgorod, Tambov and Samara regions, the Republics of Mordovia, Chuvashia, Mari El, Tatarstan and Bashkortostan.

Biology: Widely specific parasite of anurans. Trixenic life cycle. Intermediate hosts - gastropod mollusks of different species; additional hosts - dipterans genera *Anopheles* and *Culex* (Skryabin, Antipin 1962).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *P. esculentus*, *R. arvalis*, *R. temporaria*, *R. asiatica*, *R. amurensis*, *B. bombina*, *P. fuscus* (rarely), *B. bufo*, *B. viridis* (rarely) (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

*Pneumonoeces asper* (Looss, 1899)

Syn.: *Haematoloechus asper* Looss, 1899

Localization: lungs.

Areas of detection: Moscow, Ivanovo, Kostroma, Nizhny Novgorod, Ulyanovsk and Samara regions, the Republics of Chuvashia and Tatarstan.

Biology: Specific parasite of frogs. Trixenic life cycle. Intermediate host - gastropod mollusks genus *Planorbis*; additional hosts - dragonflies genera *Calopteryx*, *Agrion* and *Lestes* (Dobrovolsky 1965b).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *P. esculentus*, *R. arvalis*, *R. temporaria*, *B. bombina* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Europe.

*Skrjabinoeces similis* (Looss, 1899)

Syn.: *Skrjabinoeces volgensis* Sudarikov, 1950

Localization: lungs.

Areas of detection: Kaluga, Moscow, Ivanovo, Kostroma, Nizhny Novgorod and Samara regions, the Republics of Mordovia, Tatarstan and Bashkortostan.

Biology: Specific parasite of frogs. Trixenic life cycle. Intermediate hosts - gastropod mollusks genera *Planorbis*, *Anisus* and *Coretus*; additional hosts - dragonfly of different species (Grabda 1960).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *P. esculentus*, *R. arvalis*, *R. temporaria* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

*Skrjabinoeces breviansa* Sudarikov, 1950

Localization: lungs.

Areas of detection: Nizhny Novgorod, Ulyanovsk and Samara regions.

Biology: Specific parasite of frogs. The life cycle has not been studied.

Definitive hosts: *P. ridibundus*, *P. lessonae* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Europe.

Family: Pleurogenidae Looss, 1899

*Pleurogenes claviger* (Rudolphi, 1819)

Localization: small intestine.

Areas of detection: Moscow, Ivanovo, Kostroma, Nizhny Novgorod, Tambov and Samara regions, the Republics of Mordovia and Tatarstan.

Biology: Widely specific parasite of amphibians. Trixenic life cycle. Intermediate hosts - gastropod mollusks genus *Bithynia*; additional hosts - dragonflies, caddisflies, beetles, mayflies, alderflies, as well isopod and amphipod crustaceans genera *Asellus*, *Gammarus* and *Pontogammarus* (Khotenovsky 1970; Grabda-Kazubskaya 1971).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *P. esculentus*, *R. arvalis*, *R. temporaria*, *R. macrocnemis*, *B. bombina*, *P. fuscus*, *H. arborea*, *H. orientalis*, *B. bufo*, *B. viridis*, *L. vulgaris*, *T. cristatus*, *L. montandoni* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: cosmopolite.

*Brandesia turgida* (Brandes, 1888)

Localization: pouch-like herniations (diverticules) in the wall of the duodenum.

Areas of detection: Nizhny Novgorod and Samara regions, the Republics of Mordovia, Chuvashia and Tatarstan.

Biology: Specific parasite of frogs. The life cycle has not been studied.

Definitive hosts: *P. ridibundus*, *P. lessonae*, *R. arvalis* (rarely), *R. temporaria* (rarely) (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

*Pleurogenoides medians* (Olsson, 1876)

Localization: small intestine.

Areas of detection: Kaluga, Moscow, Kostroma, Nizhny Novgorod and Samara regions, the Republics of Mordovia, Chuvashia, Tatarstan and Bashkortostan.

Biology: Widely specific parasite of anurans. Trixenic life cycle. Intermediate hosts - gastropod mollusks genera

*Bithynia*, *Lymnaea* and *Planorbarius*; additional hosts - aquatic arthropods: dragonflies, caddisflies, mayflies, beetles, alderflies, dipterans of different species, amphipods and isopods crustaceans genus *Asellus* (Khotenovsky 1970).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *P. esculentus*, *R. arvalis*, *R. temporaria*, *R. amurensis*, *B. bombina*, *P. fuscus*, *H. arborea*, *H. orientalis*, *B. bufo*, *B. viridis*, *L. vulgaris*, *T. cristatus* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

*Prosotocus confusus* (Looss, 1894)

Localization: stomach, small intestine.

Areas of detection: Moscow, Ivanovo, Kostroma, Nizhny Novgorod, Tambov and Samara regions, the Republics of Mordovia, Chuvashia and Tatarstan.

Biology: Widely specific parasite of anurans. Trixenic life cycle. Intermediate hosts - gastropod mollusks genera *Bithynia* and *Codiella*; additional hosts - aquatic arthropods: dragonflies, beetles, caddisflies, mayflies, alderflies of different species and amphipods genus *Gammarus* (Shevchenko, Vergun 1961; Khotenovsky 1970).

Definitive hosts: *P. ridibundus*, *P. lessonae*, *P. esculentus*, *R. arvalis*, *R. temporaria*, *B. bombina*, *P. fuscus*, *B. bufo*, *B. viridis* (Vojtkova, Roca 1994; Chikhlyayev et al. 2012a, b; Kuzmin 2012).

Distribution: Palearctic.

The largest species diversity of trematodes was recorded in pool frogs in Nizhny Novgorod (17 species) and Samara (14) regions, the Republic of Tatarstan (14); less diverse species composition registered in Moscow (11), Kostroma (10) and Ivanovo (9) regions, the Republics of Mordovia (10); the minimum - in the Republic of Chuvashia (7) and Bashkortostan (6), Tambov (5) and Kaluga (4) regions, the Republic of Mari El (3) and Ulyanovsk (2) region (Table 1). These distinctions have biotopical character, that is, depend on a variety of conditions of dwelling in different a biotope with the individual complex abiotic (type and character of reservoir) and biotic (composition of flora, invertebrates and vertebrates) factors. Also, they can be bound to the geographic location, level of anthropogenic influence and a difference in volumes of frogs samplings.

The trematodes composition of the pool frog strongly differs in the Volga basin. Of the recorded 19 species any is not noted in all regions. Other species the trematode of *P. variegatus* found in 12 regions of 13 examined meets more often. Two more species (*D. subclavatus*, *O. ranae*) were detected at 10 regions, and 5 species (*P. asper*, *S. similis*, *P. claviger*, *P. medians*, *P. confusus*) at 8-9 regions. Other species in the habitat range of this host were less common, and their findings were of a sporadic nature. The rest of the trematodes, in particular, *G. loossi*, *G. vitelliloba* and *D. rastellus*, were found locally and they were observed in 1-2 regions (Table 1).

The development of the trematodes species registered at the marita stage in the pool frog in the biocoenosis of the

Volga basin proceeds on 6 types and 2 subtypes from 3 groups of life cycles. The greatest part circulates on complex - trixenic (15 species); a smaller one - on dixenic (1) and tetra-xenic (1) - cycles. The development cycles of two trematodes species (*S. breviansa*, *B. turgida*) are unknown. Species with known life cycles develop with involvement of gastropods (11 species) and bivalve (6) mollusks as intermediate hosts; insects (12), crustaceans (4) and young (tadpoles and yearlings) amphibians (4) - in the role of additional hosts.

The helminth fauna of amphibians depends on their way of life, the nature of a biotope, stay duration in water

and food spectrum. The pool frog helminths composition in the Volga basin for 50% is presented by adult stages of trematodes. Another 50% are necessary on nematodes, monogeneans and larval stages of trematodes. The reason for this is the semi-aquatic life style and consumption of aquatic invertebrates - additional hosts of trematodes. Invermination indices with many of them are usually relatively low, due to the narrow biotopic specialization of the amphibians inhabiting forest reservoirs. In these conditions, separate species (*P. variegatus*, *O. ranae*, *D. subclavatus*), can often occur and be ordinary (background) parasites of the pool frog.

**Table 1.** Adult stages of trematodes of a pool frog *Pelophylax lessonae* in the Volga basin regions

Trematodes species	KL	MS	IV	KS	NN	TM	UL	SM	MR	CH	ME	TT	BS
Class Trematoda Rudolphi, 1808													
Order Hemiurida Skrjabin et Guschanskaja, 1956													
Family Derogenidae Nicoll, 1910													
<i>Halipegus ovocaudatus</i>		+			+			+				+	
Order Paramphistomida Skrjabin et Schulz, 1937													
Family Diplodiscidae Cohn, 1904													
<i>Diplodiscus subclavatus</i>		+	+		+	+		+	+	+	+	+	+
Order Plagiorchiida La Rue, 1957													
Family Gorgoderidae Looss, 1899													
<i>Gorgodera cygnoides</i>			+		+				+			+	+
<i>Gorgodera loossi</i>													+
<i>Gorgodera microovata</i>					+				+			+	
<i>Gorgodera pagenstecheri</i>					+			+			+	+	
<i>Gorgodera varsoviensis</i>		+		+	+			+				+	
<i>Gorgoderina vitelliloba</i>					+			+					
Family Plagiorchiidae Luhe, 1901													
<i>Haplometra cylindracea</i>		+	+	+	+								
Family Telorchidae Looss, 1899													
<i>Dolichosaccus rastellus</i>					+								
<i>Opisthioglyphe ranae</i>	+	+	+	+	+	+		+	+	+		+	
Family Haematoloechidae Freitas et Lent, 1939													
<i>Pneumonoeces variegatus</i>	+	+	+	+	+	+		+	+	+	+	+	+
<i>Pneumonoeces asper</i>		+	+	+	+		+	+		+		+	
<i>Skrjabinoeces similis</i>	+	+	+	+	+			+	+			+	+
<i>Skrjabinoeces breviansa</i>					+		+	+					
Family Pleurogenidae Looss, 1899													
<i>Brandesia turgida</i>					+			+	+	+		+	
<i>Pleurogenes claviger</i>		+	+	+	+	+		+	+			+	
<i>Pleurogenoides medians</i>	+	+		+	+			+	+	+		+	+
<i>Prosotocus confusus</i>		+	+	+	+	+		+	+	+		+	
Species in total	4	11	9	10	17	5	2	14	10	7	3	14	6
	22	121	132	134	418	40	17	203	53	38	8	257	17
Examined, specimens													

Note: KL: Kaluga region (Chikhlyayev et al. 2016a); MS: Moscow region (Kotova 1936); IV: Ivanovo region (Kirillova 2002; Kirillova, Egorov 2002); KS: Kostroma region (Radchenko, Budalova 1980); NN: Nizhny Novgorod region (Sudarikov 1950, 1951; Borisova 1988; Nosova 1983, 1985, 1990, 1993; our data); TM: Tambov region (Rezvantseva, Chikhlyayev 2005; Kolodina et al. 2016); UL: Ulyanovsk region (Indiryakova et al. 2008); SM: Samara region (Evlanov et al. 2001, 2002; Chikhlyayev 2004, 2009, 2017; Fayzulin et al. 2013; Kirillov et al., 2018; our data); MR: Republic of Mordovia (Chikhlyayev et al. 2015; Ruchin et al. 2016; our data); CH: Republic of Chuvashia (Chikhlyayev, Fayzulin 2015; our data); ME: Republic of Mari El (our data); TT: Republic of Tatarstan (Smirnova 1968, 1970; Smirnova, Sizova 1978; Smirnova et al. 1987; Shal'dybin 1974, 1977; our data); BS: Republic of Bashkortostan (Ayupov et al. 1974; Bayanov 1992; Yumagulova 2000)

## ACKNOWLEDGMENTS

The study was conducted with the support of a grant RFBR 14-04-97031 r\_povolzhje\_a.

## REFERENCES

- Ayupov KV, Valiullin SM, Khaziev GZ, Bayanov MG, Kazadaev VI, Antonov PP. 1974. Helminths of animals, humans and plants in the Bashkir ASSR. In: Helminths of Animals, Humans and Plants in the Southern Urals. Issue 1. Institute of Biology of the Bashkir branch Academy of Sciences of the USSR, Ufa.
- Bayanov MG. 1992. Helminths of amphibians in Bashkortostan. In: Problems of Animal Ecology of Southern Ural. Vol 5. Publishing House of the Bashkir State University, Ufa. Chief department of VINITI, No. 587-B92.
- Borisova VI. 1988. Helminths of amphibians geographical distribution study. *Parazitologiya* 22 (6): 471-475.
- Borkin LJ, Vinogradov AE, Rozanov IM, Tsaune IA. 1987. Hemiclonal inheritance in the *Rana esculenta* hybridogenetic complex: proof by the method of flow DNA cytometry. *Reports of Academy of Sciences of the USSR* 295 (5): 1261-1264.
- Chikhlyayev IV. 2004. Helminths of amphibians of the Middle Volga region (fauna, ecology) [PhD Thesis]. Institute of Parasitology RAS, Moscow [Russia].
- Chikhlyayev IV. 2009. On the helminths of the Pool frog *Rana lessonae* Camerano, 1882 in Samara city. *Bulletin of Mordovia University. Ser Biol Sci* 1: 96-98.
- Chikhlyayev IV. 2017. Materials on the helminth fauna of the Pool frog *Pelophylax lessonae* (Camerano, 1882) in Zhiguli Nature Reserve. Samarskaya Luka: Problems of Regional and Global Ecology 26 (4): 244-248.
- Chikhlyayev IV, Fayzulin AI. 2015. Materials on the helminth fauna of the Pool frog *Pelophylax lessonae* (Camerano, 1882) in the Chuvash Republic. *Scientific Papers of Priskursky State Nature Reserve* 30 (1): 276-279.
- Chikhlyayev IV, Kirillov AA, Kirillova NY. 2012a. Trematodes (Trematoda) of amphibians (Amphibia) of the Middle Volga region. 1. Orders Fasciolida, Hemiurida, Paramphistomida and Strigeida. *Parazitologiya* 46 (3): 171-192.
- Chikhlyayev IV, Kirillov AA, Kirillova NY. 2012b. Trematodes (Trematoda) of amphibians (Amphibia) of the Middle Volga region. 2. Order Plagiorchiida. *Parazitologiya* 46 (4): 290-313.
- Chikhlyayev IV, Kirillova NY, Kirillov AA. 2018. Ecological analysis of trematodes (Trematoda) of Marsh frog *Pelophylax ridibundus* (Ranidae, Anura) from various habitats of the National Park «Samarskaya Luka» (Russia). *Nature Conservation Research* 3 (Suppl.1): 36-50. DOI: 10.24189/ncr.2018.039
- Chikhlyayev IV, Korzikov VA, Fayzulin AI. 2016a. Materials for the helminth fauna of the Pool frog *Pelophylax lessonae* and the Common toad *Bufo bufo* (Amphibia, Anura) in Kaluga region. *Bulletin of Samara Scientific Center of the Russian Academy of Sciences* 5 (2): 377-381.
- Chikhlyayev IV, Ruchin AB. 2014. The helminth fauna study of European common brown frog (*Rana temporaria* Linnaeus, 1758) in the Volga basin. *Acta Parasitol* 59 (3): 459-471. DOI: 10.2478/s11686-014-0268-5
- Chikhlyayev IV, Ruchin AB, Fayzulin AI. 2015. Helminths of tailless amphibians (Anura, Amphibia) of the Mordovia reserve. *Proceedings of Mordovia State Nature Reserve* 14: 376-388.
- Chikhlyayev IV, Ruchin AB, Fayzulin AI. 2016b. The helminth fauna study of European common toad in the Volga basin. *Nature, Environment and Pollution Technology* 15 (3): 1103-1109.
- Dedukh D, Litvinchuk S, Rosanov J, Mazepa G, Saifitdinova A, Shabanov D, Krasikova A. 2015. Optional endoreplication and selective elimination of parental genomes during oogenesis in diploid and triploid hybrid European water frogs. *PLoS ONE* 10 (4): e0123304. DOI: 10.1371/journal.pone.0123304
- Dobrovolsky AA. 1965a. Some information about the life cycle of the trematode *Opisthioglyphe ranae* (Froelich, 1791) (Plagiorchiidae). *Helminthologia* 3: 205-221.
- Dobrovolsky AA. 1965b. The life cycle of *Pneumonoeces asper* Looss, 1899 (Plagiorchiidae, Pneumonoecinae). *Proceedings of the All-Union Society of Helminthologists Conference. Part 4. Moscow. USSR Academy of Sciences.* pp 59-64.
- Dobrovolsky AA, Raikhel AS. 1973. The life cycle of *Haplometra cylindracea* Zeder, 1800 (Trematoda, Plagiorchiidae). *Bulletin of Leningrad University* 3: 5-13.
- Evlanov IA, Kirillov AA, Chikhlyayev IA, Guzova NY, Zhiltsova LV. 2001. Parasites of vertebrates of the Samara region. Part 1: A systematic catalog. Institute of Ecology of the Volga River Basin RAS, Togliatti.
- Evlanov IA, Kirillov AA, Chikhlyayev IA, Guzova NY, Zhiltsova LV. 2002. Parasites of vertebrates of the Samara region. Part 2: Distribution of parasites by hosts. Institute of Ecology of the Volga River Basin RAS, Togliatti.
- Fayzulin AI, Chikhlyayev IV, Kuzovenko AE. 2013. Amphibians of the Samara region: Monograph. Cassandra, Togliatti.
- Fayzulin AI, Zamaletdinov RI, Litvinchuk SN, Rosanov JM, Borkin LJ, Ermakov OA, Ruchin AB, Lada GA, Svinin AO, Bashinsky IV, Chikhlyayev IV. 2018. Species composition and distributional peculiarities of green frogs (*Pelophylax esculentus* complex) in Protected Areas of the Middle Volga Region (Russia). *Nature Conservation Research* 3 (Suppl.1): 1-16. DOI: 10.24189/ncr.2018.056
- Grabda B. 1960. Life cycle of *Haematoloechus similis* (Looss, 1899) (Trematoda: Plagiorchiidae). *Acta Parasitol Polon* 8: 357-366.
- Grabda-Kazubska B. 1969. Studies on abbreviation of the life-cycle in *Opisthioglyphe ranae* (Froelich, 1791) and *O. rastellus* (Olsson, 1876) (Trematoda, Plagiorchiidae). *Acta Parasitol Polon* 16: 20-27.
- Grabda-Kazubska B. 1970. Studies on the life-cycle of *Haplometra cylindracea* (Zeder, 1800) (Trematoda, Plagiorchiidae). *Acta Parasitol Polon* 18: 497-512.
- Grabda-Kazubska B. 1971. Life cycle of *Pleurogenes claviger* (Rudolphi, 1819) (Trematoda: Pleurogenidae). *Acta Parasitol Polon* 19: 337-348.
- Hofman S, Pabijan M, Dziewulska-Szwajkowska D, Szymuraa JM. 2012. Mitochondrial genome organization and divergence in hybridizing central European waterfrogs of the *Pelophylax esculentus* complex (Anura, Ranidae). *Gene* 491 (1): 71-80. <https://doi.org/10.1016/j.gene.2011.08.004>
- Hotz H, Guex G.-D, Beerli P, Semlitsch RD, Pruvost NBM. 2008. Hemiclone diversity in the hybridogenetic frog *Rana esculenta* outside the area of clone formation: the view from protein electrophoresis. *J Zool Syst Evol Res* 46 (1): 56-62. DOI: 10.1111/j.1439-0469.2007.00430.x
- Indiryakova TA, Romanova EM, Matveeva EA. 2008. Helminth fauna species diversity of amphibians in the Ulyanovsk region. *Proceedings OGAU* 1 (17): 172-176.
- Kalabekov AL. 1976. Development cycles of some trematodes of Longlegged Wood frog (*Rana macrocnemis* Boul.). Ecology and biology issues of animals of the northern slopes of the Central Caucasus: Collection of Zoological papers. 3-42.
- Kechemir N. 1976. Cycle a quatre hotes obligatoires du trematode hemiuride *Halipegus ovocaudatus*. *Bull Soc Zool France* 101 (5): 1061-1062.
- Keys to the Trematoda (Eds Gibson DI, Jones A, Bray RA). 2002. Vol 1. CABI Publishing and London The Natural History Museum, Wallingford, UK.
- Keys to the Trematoda (Eds Bray RA, Gibson DI, Jones A). 2008. Vol 3. CABI Publishing and London The Natural History Museum, Wallingford, UK.
- Khotenovsky IA. 1970. The family Pleurogenidae Looss, 1899. In: Skryabin KI (ed). Trematodes of Animals and Human. Bases of Trematodology. Vol 23. Nauka, Moscow.
- Kirillov AA, Kirillova NY, Chikhlyayev IV. Trematodes of terrestrial vertebrates of the Middle Volga region. 2012. Cassandra, Togliatti.
- Kirillov AA, Kirillova NY, Chikhlyayev IV. Parasites of vertebrate animals of the Samara region: Monograph. 2018. Poliar, Togliatti.
- Kirillova YA. 2002. Helminth fauna of tailless amphibians of the order Anura in Central Non-Chernozem Zone of Russian Federation [PhD Thesis]. Ivanovo State University, Ivanovo [Russia].
- Kirillova YA, Egorov SV. 2002. Helminth fauna of tailless amphibians of the Ivanovo region. In: Theory and Practice of Fight Against Parasitic Diseases (Zoonoses); Materials of a Scientific Conference. Vol 3. Moscow [Russia].

- Kolodina AS, Pyatova MV, Ravkovskaya EA, Lada GA. 2016. To the question of nutrition and helminths of the Pool frog (*Pelophylax lessonae*) in the conditions of Tambov region. Bulletin of Tambov University. Ser Natur and Techn Sci 21(5): 1791-1796.
- Kostyunin VM. 2010. Helminth fauna of terrestrial vertebrates of the Middle Povolzhie: Monograph. NGPU Publishing, Nizhny Novgorod.
- Kotova EN. 1936. Parasitic worms of fish and amphibians of the Klyazma river in the area of the Bolshevo Biological Station. Notes of the Bolshevo Biological Station 9: 139-140.
- Kuzmin SL. 2012. Amphibians of the former USSR (2nd edit.). Association of Scientific Publications KMK, Moscow.
- Nosova KF. 1983. Helminths of tailless amphibians in green space of Gorky city. In: Fauna, Taxonomy, Biology and Ecology of Helminths and Their Intermediate Hosts; Intercollege Collection of Scientific Papers. Gorky State Pedagogical University, Gorky [USSR].
- Nosova KF. 1985. To the study of lakes origin and zonality influence of the on the helminth fauna of the Pool frog. Helminths and Their Intermediate Hosts; Intercollege Collection of Scientific Papers. Gorky State Pedagogical University, Gorky [USSR].
- Nosova KF. 1990. Helminths of anurans in lake ecosystems of the middle belt of the European part of the RSFSR [PhD Thesis]. Gorky State Pedagogical University, Gorky [USSR].
- Nosova KF. 1993. Helminth fauna of the Pool frog in Nizhny Novgorod region. Publishing House of NSPI, Nizhny Novgorod. Chief department of VINITI, No. 2038-B93.
- Pigulevsky SV. 1952. The family Gorgoderidae Looss, 1901. In: Skryabin KI (ed). Trematodes of Animals and Human. Bases of Trematodology. Vol 7. Part 1. Publishing House of the USSR Academy of Sciences, Moscow.
- Pigulevsky SV. 1953. The family Gorgoderidae Looss, 1901. In: Skryabin KI (ed). Trematodes of Animals and Human. Bases of Trematodology. Vol 8. Part 2. Publishing House of the USSR Academy of Sciences, Moscow.
- Radchenko NM, Budalova TM. 1980. Helminths of amphibians in Kostroma region. In: IX Conference of Ukrainian Parasitological Society; Abstracts of Scientific Conference. Part 3. Naukova Dumka, Kiev.
- Reshetnikov AN, Sokolov SG, Chikhlyayev IV, Fayzulin AI, Kirillov AA, Kuzovenko AE, Protasova EN, Skomorokhov MO. 2013. Direct and indirect interactions between an invasive alien fish (*Perccottus glenii*) and two native semi-aquatic snakes. Copeia 1: 103-110.
- Rezvantseva MV, Chikhlyayev IV. 2005. About helminths of green frogs in Tambov region. In: Actual Problems of Herpetology and Toxinology; Collection of Research Papers. Issue 8. Institute of Ecology of the Volga River Basin RAS, Togliatti [Russia].
- Ruchin AB, Chikhlyayev IV, Lukijanov SV. 2009. Analysis of helminthofauna of Common spadefoot *Pelobates fuscus* (Laurenti, 1768) and Moor frog *Rana arvalis* Nilsson, 1842 (Amphibia: Anura) at their joint habitation. Parazitologiya 43 (3): 240-247.
- Ruchin AB, Kirillov AA, Chikhlyayev IV, Kirillova NY. 2016. Parasitic worms of land vertebrates in Mordovia Reserve (annotated list of species). Moscow.
- Ryzhikov KM, Sharpilo VP, Shevchenko NN. 1980. Helminths of amphibian fauna of the USSR. Nauka, Moscow.
- Shaldybin SL. 1974. The role of anurans in coastal biogeocoenosis [PhD Abstract]. Kazan State University, Kazan [USSR].
- Shaldybin SL. 1977. To the parasitic fauna of anurans in Volga-Kama Nature Reserve. In: Questions of Herpetology. Nauka, Leningrad.
- Shevchenko NN, Vergun GI. 1961. On the life cycle of amphibian trematodes *Prosotocus confusus* (Looss, 1894) Looss, 1899. Helminthologia 3(1-4): 294-298.
- Skryabin KI. 1949. Trematodes of Animals and Humans. Bases of Trematodology. Vol 3. Suborder Paramphistomatata. Publishing House of the USSR Academy of Sciences, Moscow-Leningrad.
- Skryabin KI, Antipin DN. 1962. Superfamily Plagiorchioidea Dollfus, 1930. In: Skryabin KI (ed). Trematodes of Animals and Humans. Bases of Trematodology. Vol 20. Publishing House of the USSR Academy of Sciences, Moscow.
- Smirnova MI. 1968. To the helminth fauna of amphibians of the Kuibyshev reservoir coast. In: Natural Resources of the Volga-Kama Region. Issue 2. Publishing House of Kazan State University, Kazan.
- Smirnova MI. 1970. Biocoenotic connections of some vertebrate animals helminths on the coast of the Kuibyshev reservoir [PhD Abstract]. Kazan State University, Kazan [USSR].
- Smirnova MI, Gorshkov PK, Sizova VG. 1987. Helminth fauna of tailless amphibians in Tatarstan Republic. Institute of Biology of Academy of Sciences of the USSR, Kazan. Chief department of VINITI, № 8067-B87.
- Smirnova MI, Sizova VG. 1978. Helminth fauna of aquatic amphibians of the green zone of Kazan city. In: Natural Resources of the Volga-Kama Region. Issue 5. Publishing House of Kazan State University, Kazan.
- Sudarikov VE. 1950. To the fauna of helminths of vertebrates of the Middle Volga region. In: Papers of the Helminthological Laboratory of USSR Academy of Sciences. Issue 5. Publishing House of Academy of Sciences of the USSR, Moscow.
- Sudarikov VE. 1951. Fauna of helminths of vertebrates of the Middle Volga region. Papers of the Helminthological Laboratory of USSR Academy of Sciences. Issue 3. Publishing House of Academy of Sciences of the USSR, Moscow.
- Vojtkova L, Roca V. Parasites of the frogs and toads in Europe. Part II: Trematoda. Rev. Esp. Herp. 1994 (8): 7-18.
- Yumagulova GR. 2000. Helminths of amphibians of the Southern Urals [PhD Abstract]. Bashkir State Agricultural University, Ufa [Russia].
- Zeisset I, Hoogesteger T. 2018. A reassessment of the biogeographic range of northern clade pool frogs (*Pelophylax lessonae*). Herpetol J 28 (2): 63-72.