

# Unveiling the interplay between tree stand attributes, species diversity, and biomass carbon in Chunati Wildlife Sanctuary, Southeastern Bangladesh

MEHEDI HASAN RAKIB\*, S. M. SAKIB HOSSAIN, MOHAMMAD WASIF MURSALIN SADNAN,  
MOHD IMRAN HOSSAIN CHOWDHURY

Institute of Forestry and Environmental Sciences, University of Chittagong, Chattogram-4331, Bangladesh.  
Tel./fax.: +880-1639222696, \*email: rakib.ifescu@gmail.com

Manuscript received: 1 October 2024. Revision accepted: 3 December 2024.

**Abstract.** Rakib MH, Hossain SMS, Sadnan MWM, Chowdhury MIH. 2024. Unveiling the interplay between tree stand attributes, species diversity, and biomass carbon in Chunati Wildlife Sanctuary, Southeastern Bangladesh. *Intl J Trop Drylands* 8: 95-105. Biodiversity plays a pivotal role in maintaining ecosystem health and resilience, especially in tropical forest landscapes. This study investigates the intricate relationships between stand structure, biodiversity indices, and biomass carbon within forest ecosystems, specifically focusing on the Chunati Wildlife Sanctuary (CWS), a tropical forestland in Southeastern Bangladesh. The key structural attributes, i.e. tree height, diameter at breast height (DBH), basal area, and stand density, as well as biodiversity indices, were analyzed to understand their correlations with biomass carbon. The study exhibits the measured soil carbon levels ranging from 35.26 to 93.26 Mg C ha<sup>-1</sup> and tree biomass carbon from approximately 131.13 to 387.23 Mg C ha<sup>-1</sup>. The findings revealed a strong positive correlation between tree height and biomass carbon ( $R^2 = 0.6517$ ), indicating that taller trees generally store more carbon due to enhanced photosynthesis. Notably, DBH exhibited a robust correlation ( $R^2 = 0.8683$ ), highlighting the significance of larger trees as carbon sinks. The basal area emerged as the strongest predictor of biomass carbon, with an impressive correlation ( $R^2 = 0.9119$ ). The Shannon-Wiener and Margalef's richness indices had moderate to strong correlations with biomass carbon ( $R^2 = 0.4473$  and  $R^2 = 0.5663$ ) respectively. These results underscore the importance of maintaining diverse and structurally complex forests for effective carbon sequestration. The future study should incorporate additional ecological variables to refine models for predicting forest carbon dynamics with the vast change in tropical landscapes and inform better conservation strategies, leading to tropical biodiversity conservation and climate change mitigation efforts on tropical landscapes.

**Keywords:** Carbon stock, fossil carbon, phytosociology, tree biomass, tropical-forest

## INTRODUCTION

Protected areas are essential for conserving biodiversity and delivering ecosystem services (Mamun et al. 2022). In Bangladesh, protected area coverage increased slightly from 1.67% in 2008 to 1.80% of the total forest land area in 2015, one of the lowest globally (Islam et al. 2022). The government has designated 49 protected areas, 17 of which are under the co-management system (Uddin et al. 2020a). Among these, Chunati Wildlife Sanctuary (CWS), covering approximately 7,764 hectares in southeastern Bangladesh, plays a vital role in conserving biodiversity while supporting the livelihoods of approximately 50,000 people across 9,400 households (Hossain et al. 2020). It encompasses various forest types, including tropical evergreen, semi-evergreen, and deciduous forests, contributing to its rich biodiversity (Festus et al. 2024). However, deforestation, illegal logging, and land conversion jeopardize this biodiversity.

The success of conservation efforts in protected areas is intricately tied to the well-being of forest-dependent communities (Hossain et al. 2020). Empowering local communities living around protected areas has been identified as a key strategy for biodiversity conservation (Rahman et al. 2018). In CWS, 70% of the population is

considered poor, 19% extremely poor, 10% middle class, and just 1% rich (Rahman et al. 2016). The primary occupations include day labor and non-wage agricultural work, with an unemployment rate at 17% (Rahman et al. 2017; Rahman and Obaydullah 2020). This heavy reliance on forest resources (Akwaji and Onah 2023), coupled with high unemployment, threatens CWS's ecological integrity (Rahman and Obaydullah 2020). To mitigate forest degradation, the Bangladesh Forest Department (BFD), in collaboration with USAID, launched the Nishorgo support project in 2004, followed by the integrated protected area co-management (IPAC) initiative (Kufata et al. 2024), which promoted community co-management and alternative income activities (Chowdhury 2014). However, limited livelihood support from these projects has reduced their effectiveness in decreasing forest dependency (Roy et al. 2015).

Forests are critical for maintaining ecological balance, regulating climate, providing habitats for species, and serving as carbon sinks (Ali et al. 2022). Consequently, understanding forest ecosystems' biodiversity and carbon stock is vital for effective conservation and sustainable management (Ali et al. 2022; Islam et al. 2018, 2020). To assess species diversity, researchers use indices such as the Shannon-Wiener index, and Margalef richness index,

which provide insights into species richness, evenness, and distribution (Kessler et al. 2009; Hasan et al. 2023; Scherer et al. 2023). These metrics inform conservation planning and biodiversity management. Assessing carbon stocks, including tree biomass carbon and soil organic carbon (SOC), is equally crucial for understanding a forest's role in carbon sequestration (Alongi 2011; Shaheed and Chowdhury 2014; Ali et al. 2022).

This study aims to evaluate the biodiversity and carbon stock in the Chunati Wildlife Sanctuary, offering a comprehensive understanding of its ecological status. The primary objectives are to analyze species diversity using the Shannon-Wiener index, Simpson's index, and Pielou's evenness index, and to quantify carbon stock by measuring both tree biomass carbon and SOC. The leading question of the study: (i) How do structural attributes such as tree height, diameter at breast height (DBH), and basal area predict biomass carbon stock in the Chunati Wildlife Sanctuary; (ii) To what extent do biodiversity indices, such as the Shannon-Wiener and Margalef's richness indices, influence the carbon sequestration capacity of tropical forest ecosystems, and how can these relationships guide conservation efforts for enhancing carbon storage?


By examining the interplay between biodiversity indices, tree stand structure, and carbon stock, this research seeks to contribute to the existing knowledge base on tropical forest biodiversity and carbon sequestration in Bangladesh. Additionally, it provides baseline data for monitoring changes in CWS's ecological parameters over time, which is essential for assessing conservation measures' effectiveness and informing forest management decisions. The findings will support policy-making at both national and international levels, particularly regarding biodiversity conservation and climate change mitigation. Insights from this study will aid in the protection and sustainable management of CWS, benefiting both local

conservation initiatives and global efforts to combat climate change. By providing data on species diversity and carbon sequestration potential, this research will inform strategies to preserve biodiversity and enhance carbon storage in tropical forests.

## MATERIALS AND METHODS

### Study site

The Chunati Wildlife Sanctuary (CWS) is situated at 21°40' North and 92°07' East, in the southeastern part of Bangladesh (Figure 1). It encompasses the Banskhalhi and Lohagara Upazila of Chittagong District and the Chakoria Upazila of Cox's Bazar District. The Chunati Wildlife Sanctuary experiences a tropical moist climate with annual rainfall ranging from 3,000 to 3,500 mm. The air temperature fluctuates between 16 and 34°C, while relative humidity remains high, typically between 70 and 90%. The soil in the sanctuary is predominantly sandy loam, supporting a rich biodiversity and influencing the carbon sequestration potential of the forest. The sanctuary covers seven Union Parishads: Chunati, Adhunagar, Herbang, Puichari, Banskhalhi, Borohatia, and Toitong. CWS was established in 1986 and spans 7,764 hectares of Reserve Forest. The Wildlife and Nature Conservation Division of Chittagong oversees the administration of the sanctuary. It is divided into two Forest Ranges and seven forest beats (smallest forest management unit): The Chunati range, which covers a total area of 3,332 ha and includes Chunati, Aziznagar, and Harbang beats; and the Jaldi range, which encompasses a total area of 4,432 ha and comprises Jaldi, Chambal, Napora, and Puichari (Islam et al. 2018; Mamun et al. 2022; Rahman et al. 2022).



**Figure 1.** Map of the study area with sampling (vegetation and soil) points throughout the seven Forest Beats of the Chunati Wildlife Sanctuary. The maps are created using Arcmap10.8

**Table 1.** Equations used in the calculation for phytosociological parameters and biodiversity indices of tree species in this study

Equation	References
Basal area ( $\text{m}^2 \text{ ha}^{-1}$ ) = $\frac{\sum \pi \times D^2}{4 \times \text{Area of all quadrats}} \times 10000$	(Shukla and Chandel 2016); Chowdhury et al. (2019)
Relative density, $R_d (\%) = \frac{S}{n} \times 100$	Dallmeier et al. (1992); Das (2024)
Relative frequency, $R_f (\%) = \frac{F_i}{\sum F_i}$	Dallmeier et al. (1992); Das (2024)
Relative dominance, $RD_o (\%) = \frac{\text{Basal area of one species}}{\text{Total Basal area}} \times 100$	Hossain et al. (2013); Chowdhury et al. (2019)
Important Value Index (IVI) = $R_d + R_f + RD_o$	Hossain et al. (2013)
Shannon-Wiener index, $H' = -\sum P_i \ln P_i$	Michael (1984); Hop et al. (2023)
Margalef richness index = $\frac{S-1}{\ln N}$	Kohli et al. (1996)

Note: D: The diameter at breast height,  $F_i$ : The frequency of a species,  $P_i$ : The abundance of the species in each plot, S: the total number of species, N: The total number of individuals of all the species

**Table 2.** Equations used in the estimation of biomass, and soil organic carbon in this study

Equation	References
AGB (kg) = $(0.0673 \times (\rho (\text{DBH})^2 H)^{0.976})$	Chave et al. (2014)
Biomass C ( $\text{Mg ha}^{-1}$ ) = Biomass (dry mass, $\text{Mg ha}^{-1}$ ) $\times 0.5 \text{ Mg C}$	Sandra Brown (1997); Baul et al. (2021a)
Loss of ignition, $LOI \% = \frac{W_1}{W_2} \times 100$	Ball (1964)
Soil organic carbon, $SOC \% = 0.47 \times (\% LOI - 1.87)$	Ball (1964); Baul et al. (2021b)
$SOC \text{ stock } (\text{Mg ha}^{-1}) = SOC \% \times BD \times SD$	Pearson et al. (2005)

Note: AGB (kg): Above-ground biomass,  $\rho$  wood density ( $\text{g cm}^{-3}$ ), D: Tree DBH (cm), H: Tree height (m),  $W_1$ : the loss in mass (g),  $W_2$ : Oven-dried soil (g), BD: Bulk density of soil ( $\text{g cm}^{-3}$ ), and SD means the soil depth (cm). Living root biomass was estimated as 15% of AGB (Macdicken K. 2015). To estimate AGB, wood density ( $\text{g cm}^{-3}$ ), a required variable, was collected from the Bangladesh Forest Research Institute (Sattar et al. 1999). For those not found in BFRI publications, we used the global wood density database (Chave et al. 2009)

### Data collection

#### Vegetation survey

Vegetation survey was conducted to assess the biodiversity indices, and biomass carbon estimation of CWS between October to December 2023. A total of 70 plots (10 from each beat) with size of  $20 \times 20 \text{ m}$  were randomly located (Figure 1). In each plot, DBH (diameter at breast height; 1.3 m above the ground), the total height of all trees having dbh  $\geq 5 \text{ cm}$  were recorded using Range finder and Suunto clinometer. All the surveyed species were identified and recorded in local and scientific name. In the case of unknown species plant samples were collected to be identified by professional taxonomists from the Bangladesh Forest Research Institute (BFRI).

#### Sampling of soil

A pit of 30 cm depth, under the litter layer approximate center of the same vegetation plot, was dug using a soil auger, and soil samples were collected from three vertical layers at 0-10, 11-20, and 21-30 cm depth. Hence 12 (3 depths  $\times$  4 points) samples from each beat which makes the result of a total of 84 (12  $\times$  7 beats) samples are collected

from the CWS (Figure 1). Collected soil samples were performed under lab analysis for SOC estimation.

### Data analysis

Many indices are available worldwide that measure phytosociological parameters, biodiversity indices, biomass estimation, and soil organic carbon. In this study, different phytosociological and biodiversity indices were calculated for all the beats of CWS using the equations presented in Table 1. The estimation of biomass and soil organic carbon, the equations presented in Table 2 were used.

## RESULTS AND DISCUSSION

### Phytosociological status

The tree species survey in the CWS recorded 100 species belonging to 41 families. Among all the families Fabaceae (9 species) contained the highest number of tree species followed by Moraceae (8 species), Myrtaceae (8 species), and Eupharbiaceae (7 species). The highest basal area was occupied by *Ficus bengalensis* ( $11.10 \text{ m}^2 \text{ ha}^{-1}$ ) in

the contrary *Garcinia lanceifolia* (0.05 m<sup>2</sup> ha<sup>-1</sup>) scored the lowest (Table 3). The highest relative density was *Acacia auriculiformis* (20.61%) followed by *Dipterocarpus turbinatus* (9.10%) and *Swietenia macrophylla* (5.11%). Among all the recorded tree species *Lagerstroemia speciosa* (2.44%) and 22 others showed identical maximum relative frequency followed by *Stereospermum personatum* (2.24%) and *A. auriculiformis* (2.21%). The species with the highest relative dominance was *F. bengalensis* (8.06%) whereas *G. lanceifolia* (0.04%) scored the lowest. The most important species of CWS in terms of IVI was *A. auriculiformis* (23.25) followed by *D. turbinatus* (14.37) and *S. robusta* (10.58), while, *G. lanceifolia* (0.17) had the lowest IVI. It is a clear indication that exotic species alongside some native species mostly dominate the CWS. These exotic species are playing a role in the overall ecosystem degradation of the CWS.

### Estimation of biomass and soil organic carbon

The comparative analysis of tree biomass carbon and soil organic carbon (SOC) across the seven forest beats of Chunati Wildlife Sanctuary indicates significant variability in both tree biomass carbon and SOC. Tree biomass carbon

ranges from approximately 131.13 to 387.23 Mg C ha<sup>-1</sup>, with the highest values observed in beats dominated by larger trees and higher basal areas. In contrast, SOC ranges from 35.26 to 93.26 Mg C ha<sup>-1</sup> across different soil layers (0-10, 11-20, and 21-30 cm), with higher values in areas with denser tree cover and organic matter input (Figure 2.B).


The substantial range in tree biomass carbon suggests that forest structure, species composition, and management practices significantly influence carbon storage capacity. The mean biomass carbon (224.43 Mg C ha<sup>-1</sup>) in the study area (Figure 2.A) is considerably higher than that of other regions in Bangladesh, such as the homestead forests in Rangpur (54 Mg C ha<sup>-1</sup>) and Sundarbans (99 Mg C ha<sup>-1</sup>), indicating the sanctuary's critical role in carbon sequestration (Baul et al. 2021b; Jaman et al. 2016). The SOC values are comparable to those found in similar tropical ecosystems, with the mean SOC (53.59 Mg C ha<sup>-1</sup>) closely matching that of the Chittagong Hill Tracts (52.83 Mg C ha<sup>-1</sup>) but exceeding that of the Sal Forest (48.3 Mg C ha<sup>-1</sup>) and Sundarbans (38.8 Mg C ha<sup>-1</sup>) (Baul et al. 2021b; Mukul et al. 2014).

**Table 3.** Phytosociological attributes of vegetation in Chunati Wildlife Sanctuary, Bangladesh


Scientific name	Family	BA (m <sup>2</sup> ha <sup>-1</sup> )	R <sub>d</sub> (%)	R <sub>f</sub> (%)	RD <sub>o</sub> (%)	IVI
<i>Mangifera indica</i> L.	Anacardiaceae	1.57	4.47	2.20	1.16	7.83
<i>Lannea coromandelica</i> Merr.	Anacardiaceae	0.12	0.05	0.03	0.09	0.17
<i>Anacardium occidentale</i> L.	Anacardiaceae	1.44	0.16	0.08	1.07	1.31
<i>Mangifera sylvatica</i> Roxb.	Anacardiaceae	4.70	0.27	0.13	3.47	3.87
<i>Alstonia scholaris</i> L.	Apocynaceae	0.78	0.27	2.44	0.57	3.28
<i>Arenga pinnata</i> Merr.	Arecaceae	0.18	0.05	2.44	0.14	2.63
<i>Stereospermum personatum</i> Hassk.	Bignoniaceae	1.07	0.59	2.24	0.79	3.62
<i>Stereospermum suaveolens</i> Roxb.	Bignoniaceae	0.22	0.05	0.20	0.16	0.42
<i>Eruca vesicaria</i> L.	Brassicaceae	0.15	0.11	2.44	0.11	2.66
<i>Bursera serrata</i> Wall.	Burseraceae	1.04	1.13	2.44	0.77	4.34
<i>Bauhinia acuminata</i> L.	Caesalpiniaceae	0.48	0.16	0.73	0.36	1.25
<i>Senna siamea</i> Lam.	Caesalpiniaceae	1.39	0.16	0.73	1.03	1.92
<i>Tamarindus indica</i> L.	Caesalpiniaceae	5.17	0.22	0.98	3.82	5.01
<i>Bhesa robusta</i> Roxb.	Celastraceae	1.32	0.22	2.44	0.97	3.63
<i>Garcinia lanceifolia</i> Roxb.	Clusiaceae	0.05	0.05	0.07	0.04	0.17
<i>Garcinia cowa</i> Roxb.	Clusiaceae	0.20	1.18	1.63	0.15	2.96
<i>Garcinia speciosa</i> Wall.	Clusiaceae	0.19	0.22	0.30	0.14	0.65
<i>Mesua ferrea</i> L.	Clusiaceae	0.76	0.32	0.44	0.56	1.33
<i>Terminalia arjuna</i> DC.	Combretaceae	1.53	1.56	0.98	1.13	3.67
<i>Terminalia bellirica</i> Gaertn.	Combretaceae	1.07	0.91	0.58	0.79	2.28
<i>Terminalia chebula</i> Retz.	Combretaceae	0.29	1.29	0.81	0.21	2.32
<i>Terminalia paniculata</i> Roth.	Combretaceae	1.28	0.11	0.07	0.94	1.12
<i>Dillenia indica</i> L.	Dilleniaceae	0.83	0.11	0.81	0.61	1.53
<i>Dillenia pentagyna</i> Roxb.	Dilleniaceae	0.15	0.22	1.63	0.11	1.95
<i>Dipterocarpus costatus</i> Gaertn.	Dipterocarpaceae	2.94	2.42	0.33	2.17	4.93
<i>Shorea robusta</i> Gaertn.	Dipterocarpaceae	8.24	4.04	0.55	6.09	10.68
<i>Dipterocarpus turbinatus</i> Gaertn.	Dipterocarpaceae	5.56	9.10	1.24	4.11	14.44
<i>Hopea odorata</i> Roxb.	Dipterocarpaceae	1.47	2.31	0.32	1.09	3.72
<i>Diospyros malabarica</i> Kostel.	Ebenaceae	0.69	0.05	2.44	0.51	3.01
<i>Elaeocarpus serratus</i> L.	Elaeocarpaceae	2.43	0.48	2.44	1.80	4.72
<i>Phyllanthus emblica</i> L.	Euphorbiaceae	1.38	0.86	0.67	1.02	2.55
<i>Aporosa wallichii</i> Hook.	Euphorbiaceae	0.35	0.22	0.17	0.26	0.64
<i>Antidesma velutinum</i> Tul.	Euphorbiaceae	0.14	0.11	0.08	0.10	0.30
<i>Antidesma ghaesembilla</i> Gaertn.	Euphorbiaceae	0.17	0.16	0.13	0.13	0.41
<i>Baccaurea ramiflora</i> Lour.	Euphorbiaceae	0.10	0.22	0.17	0.07	0.45
<i>Trewia nudiflora</i> L.	Euphorbiaceae	0.86	1.51	1.18	0.63	3.32

<i>Vernicia fordii</i> Hemsl.	Euphorbiaceae	0.35	0.05	0.04	0.26	0.35
<i>Acacia auriculiformis</i> A.	Fabaceae	0.59	20.61	2.21	0.43	23.26
<i>Pongamia pinnata</i> L.	Fabaceae	0.21	0.05	0.01	0.15	0.21
<i>Castanopsis indica</i> Roxb.	Fabaceae	0.49	1.18	0.13	0.36	1.67
<i>Ormosia robusta</i> Baker	Fabaceae	0.15	0.11	0.01	0.11	0.23
<i>Acacia hybrida</i> G.Lodd.	Fabaceae	0.53	0.16	0.02	0.39	0.57
<i>Quercus dentata</i> Thunb.	Fabaceae	0.30	0.05	0.01	0.22	0.28
<i>Xylia xylocarpa</i> Roxb.	Fabaceae	3.63	0.05	0.01	2.68	2.74
<i>Erythrina variegata</i> L.	Fabaceae	1.13	0.32	0.03	0.84	1.19
<i>Butea monosperma</i> Lam.	Fabaceae	0.72	0.16	0.02	0.53	0.71
<i>Gmelina arborea</i> Roxb.	Lamiaceae	1.13	3.28	1.75	0.84	5.87
<i>Tectona grandis</i> L.	Lamiaceae	2.01	1.29	0.69	1.49	3.47
<i>Cinnamomum tamala</i> Buch.	Lauraceae	0.72	0.11	2.44	0.53	3.08
<i>Barringtonia acutangula</i> L.	Lecythidaceae	0.69	0.11	2.44	0.51	3.06
<i>Lagerstroemia speciosa</i> (L.) Pers.	Lythraceae	0.95	1.83	2.44	0.70	4.97
<i>Magnolia champaca</i> (L.) Baill. ex Pierre	Magnoliaceae	2.23	0.54	2.44	1.65	4.63
<i>Sterculia foetida</i> L.	Malvaceae	0.89	0.05	0.17	0.66	0.89
<i>Brownlowia elata</i> R. Br.	Malvaceae	1.26	0.05	0.17	0.93	1.16
<i>Bombax ceiba</i> L.	Malvaceae	0.45	0.59	1.92	0.33	2.84
<i>Abroma augustum</i> (L.) L. f.	Malvaceae	0.75	0.05	0.17	0.55	0.78
<i>Chukrasia tabularis</i> A. Juss.	Meliaceae	0.70	1.83	0.60	0.52	2.95
<i>Swietenia macrophylla</i> King	Meliaceae	4.15	5.11	1.68	3.07	9.86
<i>Azadirachta indica</i> A. Juss.	Meliaceae	2.59	0.43	0.14	1.91	2.48
<i>Aphanamixis polystachya</i> (Wall.) R. Parker	Meliaceae	0.52	0.05	0.02	0.39	0.46
<i>Albizia lebbek</i> (L.) Benth.	Mimosaceae	1.99	0.48	0.32	1.47	2.27
<i>Acacia mangium</i> Willd.	Mimosaceae	0.71	0.48	0.32	0.52	1.33
<i>Samanea saman</i> (Jacq.) Merr.	Mimosaceae	4.62	1.56	1.04	3.41	6.01
<i>Albizia richardiana</i> (Voigt) King & Prain	Mimosaceae	4.47	0.16	0.11	3.30	3.57
<i>Albizia procera</i> (Roxb.) Benth.	Mimosaceae	2.41	0.97	0.65	1.78	3.39
<i>Ficus religiosa</i> L.	Moraceae	1.38	0.38	0.12	1.02	1.52
<i>Ficus benghalensis</i> L.	Moraceae	11.10	0.32	0.11	8.21	8.64
<i>Artocarpus chama</i> Buch.-Ham.	Moraceae	0.78	1.88	0.62	0.57	3.08
<i>Artocarpus lacucha</i> Buch.-Ham.	Moraceae	1.35	0.43	0.14	1.00	1.57
<i>Ficus racemosa</i> L.	Moraceae	0.36	2.15	0.71	0.26	3.12
<i>Artocarpus heterophyllus</i> Lam.	Moraceae	0.58	1.72	0.57	0.43	2.72
<i>Ficus auriculata</i> Lour.	Moraceae	0.24	0.05	0.02	0.18	0.25
<i>Streblus asper</i> Lour.	Moraceae	1.11	0.48	0.16	0.82	1.47
<i>Muntingia calabura</i> L.	Muntingiaceae	0.26	0.48	2.44	0.20	3.12
<i>Syzygium grande</i> (Wight) Walp.	Myrtaceae	1.74	4.31	1.04	1.29	6.64
<i>Eucalyptus globulus</i> Labill.	Myrtaceae	0.45	2.10	0.51	0.34	2.94
<i>Syzygium cumini</i> (L.) Skeels	Myrtaceae	3.60	1.13	0.27	2.66	4.06
<i>Melaleuca alternifolia</i> (Maiden & Betche) Cheel	Myrtaceae	0.79	0.05	0.01	0.58	0.65
<i>Psidium guajava</i> L.	Myrtaceae	0.26	0.70	0.17	0.19	1.06
<i>Syzygium praecox</i> (Roxb.) Rathakr. & Chithra	Myrtaceae	0.11	0.22	0.05	0.08	0.35
<i>Syzygium fruticosum</i> (Roxb.) A. M. Cowan & Cowan	Myrtaceae	0.79	1.40	0.34	0.59	2.32
<i>Eucalyptus hybrida</i> (no specific authority, hybrid designation)	Myrtaceae	0.69	0.16	0.04	0.51	0.71
<i>Nyctanthes arbor-tristis</i> L.	Oleaceae	0.20	0.05	2.44	0.15	2.64
<i>Averrhoa carambola</i> L.	Oxalidaceae	0.52	0.32	2.44	0.38	3.14
<i>Glochidion lanceolarium</i> (Roxb.) Voigt	Phyllanthaceae	0.67	1.29	2.44	0.50	4.23
<i>Ziziphus mauritiana</i> Lam.	Rhamnaceae	0.21	0.16	2.44	0.16	2.76
<i>Prunus amygdalus</i> Batsch	Rosaceae	1.03	0.16	2.44	0.76	3.36
<i>Adina cordifolia</i> (Roxb.) Hook. f. ex Brandis	Rubiaceae	1.32	0.11	0.41	0.97	1.49
<i>Neolamarckia cadamba</i> (Roxb.) Bosser	Rubiaceae	5.75	0.54	2.03	4.25	6.82
<i>Zanthoxylum rhetsa</i> (Roxb.) DC.	Rutaceae	1.31	0.05	0.81	0.97	1.83
<i>Aegle marmelos</i> (L.) Corrêa	Rutaceae	0.95	0.05	0.81	0.70	1.57
<i>Citrus limetta</i> Risso	Rutaceae	0.05	0.05	0.81	0.04	0.91
<i>Litchi chinensis</i> Sonn.	Sapindaceae	0.14	0.65	2.44	0.10	3.19
<i>Madhuca longifolia</i> (J. König) J. F. Macbr.	Sapotaceae	0.38	0.05	2.44	0.28	2.78
<i>Spondias mombin</i> L.	Spondiadiaceae	3.05	0.05	2.44	2.25	4.75
<i>Sterculia villosa</i> Roxb.	Sterculiaceae	0.27	0.11	2.44	0.20	2.75
<i>Tamarix dioica</i> Roxb. ex Roth	Tamaricaceae	0.46	0.48	2.44	0.34	3.26
<i>Aquilaria malaccensis</i> Lam.	Thymelaeaceae	0.66	0.48	2.44	0.49	3.41
<i>Vitex peduncularis</i> Wall. ex Schauer	Verbenaceae	0.30	0.05	0.49	0.22	0.76
<i>Vitex pinnata</i> L.	Verbenaceae	0.83	0.22	1.95	0.61	2.78
Total		135.3	100	100	100	300

Note: basal area (BA m<sup>2</sup> ha<sup>-1</sup>), relative density (Rd %), relative frequency (Rf %), relative dominance (Rdo %), and important value index (IVI) found from the recorded tree species



**Figure 2.** A beat-wise comparison among the seven beats in Chunati Wildlife Sanctuary in term of: A. tree biomass carbon stock; B. average soil organic carbon across 3 vertical layers (0-10, 11-20, 21-30 cm). The error bar represents the standard error of the mean



**Figure 3.** The tree biomass carbon of ten tree species with the most Important Value Index (IVI) in the Chunati Wildlife Sanctuary

This figure underscores the importance of both tree biomass and soil carbon in assessing the overall carbon storage potential of tropical forests. The variation across different beats highlights the need for site-specific forest management strategies to maximize carbon sequestration while maintaining biodiversity (Rahman et al. 2018). The findings align with existing literature, which emphasizes the crucial role of forest structure in carbon dynamics and highlights the potential of tropical forest ecosystems for significant carbon sequestration (Chave et al. 2014).

Figure 3 illustrates the biomass carbon contributions of the ten most dominant tree species in Chunati Wildlife Sanctuary based on their Importance Value Index (IVI). Species like *Dipterocarpus turbinatus*, *Shorea robusta*, and *Neolamarckia cadamba* demonstrate the highest biomass carbon storage, indicating their significant role in carbon sequestration within the sanctuary. In contrast, fast-growing exotic species such as *Acacia auriculiformis* and

*Swietenia macrophylla*, despite being dominant by IVI, show relatively lower biomass carbon, suggesting that while these species thrive in number, they contribute less to carbon storage compared to native slow-growing species. This disparity highlights the ecological importance of preserving native species for maximizing carbon sequestration in tropical forests.

**Interplay between tree stands structure and biomass carbon**

The relationship between tree biomass carbon and various stand structure parameters is crucial for understanding forest carbon dynamics and for developing effective strategies for carbon sequestration in forest ecosystems. Figure 4 illustrates these relationships by examining tree height, tree diameter at breast height (DBH), basal area, and stand density, all of which are key structural attributes of forest stands (Pozo and Säumel

2018). By analyzing these parameters, we can gain insights into how forest structure influences carbon storage and, by extension, the role of forests in mitigating climate change.

#### *Tree height and biomass carbon*

Tree height is a fundamental parameter in forestry as it often correlates with the overall growth and productivity of a tree. Figure 4.A shows the relationship between tree height and biomass carbon is shown to be positively correlated. The regression equation  $y = 32.99x - 201.07$  reveals that biomass carbon increases with tree height. The coefficient of determination  $R^2 = 0.6517$  indicates that approximately 65% of the variation in biomass carbon can be explained by tree height (Ali et al. 2022; Chowdhury and Das 2024a; Chowdhury and Rakib 2024b). This strong correlation suggests that taller trees generally accumulate more biomass carbon, likely due to their greater leaf area and increased photosynthetic capacity, which leads to higher growth rates and carbon assimilation. However, the fact that 35% of the variation remains unexplained suggests that other factors (Scherer et al. 2023), such as species-specific growth patterns (Hossain et al. 2020), soil fertility (Mamun et al. 2022), and climatic conditions, also play significant roles in determining biomass carbon. This finding aligns with previous studies that have emphasized the importance of tree height in carbon storage but also highlight the need to consider additional factors that might influence carbon sequestration at different stages of tree growth (Zukswert et al. 2023).

#### *Tree DBH and biomass carbon*

Tree DBH is another critical indicator of forest stand structure and is often used in allometric equations to estimate tree biomass and carbon storage. Figure 4.B shows a robust positive correlation between tree DBH and biomass carbon, with a regression equation of  $y = 20.376x - 268.74$  and a high  $R^2 = 0.8683$ . This suggests that 87% of the variation in biomass carbon is explained by tree DBH, making DBH one of the most reliable predictors of biomass carbon in forest ecosystems. Larger DBH typically indicates older and more mature trees (Rahman et al. 2018; Mehta et al. 2021), which have had more time to accumulate biomass (Sheikh et al. 2011). This strong relationship underscores the importance of preserving large-diameter trees in forests, as they serve as significant carbon sinks. The remaining 13% of unexplained variation could be attributed to factors such as tree species composition, age, and environmental conditions, which can influence how biomass is distributed within a tree and across different trees in a stand (Mamun et al. 2022; Ali et al. 2022). This finding suggests that forest management practices should prioritize the protection and maintenance of large-diameter trees to enhance carbon sequestration efforts (Hossain et al. 2020).

#### *Basal area and biomass carbon*

Basal area (BA) is a measure of the cross-sectional area of a tree trunk and is often used as an indicator of forest density and productivity. Figure 4.C demonstrates a very strong positive correlation between basal area and biomass

carbon with the regression equation  $y = 1.7256x - 44.008$  and an impressive  $R^2 = 0.9119$ . This suggests that about 91% of the variability in biomass carbon is explained by basal area, making it the strongest predictor among the variables analyzed. Basal area is directly related to the total wood volume in a forest stand, and thus, higher basal areas typically correspond to greater biomass and carbon storage (Rahman et al. 2018; Mamun et al. 2022). The strong relationship between basal area and biomass carbon highlights the importance of managing forest stands to maintain or increase basal area, particularly in the context of forest conservation and carbon sequestration efforts. The small amount of unexplained variation (9%) might be due to differences in tree form, species diversity, and stand age, which can influence the distribution of biomass within a stand (Simon et al. 2018; Hossain et al. 2020; Miah et al. 2023).

#### *Stand density and biomass carbon*

Stand density, measured as the number of trees per hectare, is another important structural attribute of forests that can influence biomass carbon. Figure 4.D shows the relationship between stand density and biomass carbon with the regression equation  $y = 0.7044x - 399.7$  and  $R^2 = 0.6175$ . This indicates that 62% of the variation in biomass carbon is related to stand density. While this relationship is still significant, it is weaker than those observed for tree height, DBH, and basal area. Stand density can influence competition among trees for resources such as light, water, and nutrients, which in turn affects growth rates and biomass accumulation. Higher stand densities might not always lead to higher biomass carbon if competition for resources, such as light, water, and nutrients, becomes too intense among trees, leading to stunted growth, lower biomass accumulation and increased mortality (Scherer et al. 2023). Conversely, lower densities might reduce competition but also limit the total biomass that can be supported in a given area (Hossain et al. 2020). The 38% of unexplained variation suggests that stand density alone is not a sufficient predictor of biomass carbon, and that factors like species composition, stand age, and management practices must also be considered.

### **The interplay between tree diversity and biomass carbon**


Figure 5 presents the relationship between tree biomass carbon and two diversity indices: the Shannon-Wiener diversity index and the Margalef richness index. These indices are widely used to assess biodiversity within ecological communities, and their relationship with biomass carbon offers insights into how biodiversity might influence carbon storage in forest ecosystems.

#### *Diversity index and biomass carbon*


Figure 5.A illustrates the relationship between the Shannon-Wiener diversity index and biomass carbon. The regression equation  $y = 109.72x - 45.844$  and the coefficient of determination  $R^2 = 0.44$  indicate a positive correlation, with approximately 45% of the variation in biomass carbon explained by the Shannon-Wiener diversity index. This

suggests that higher biodiversity, as measured by the Shannon-Wiener index, is associated with greater biomass carbon storage. The Shannon-Wiener index accounts for both species richness and evenness, meaning that more diverse and evenly distributed species communities tend to support higher levels of biomass (Rahman et al. 2018).

However, with 55% of the variation left unexplained, it is clear that while biodiversity is an important factor, other variables such as environmental conditions, species-specific traits, and forest management practices also significantly influence biomass carbon.



**Figure 4.** Relationship between tree biomass carbon and stand structure in term of: A. Tree height; B. Tree DBH; C. Basal area; and D. Stand density



**Figure 5.** Relationship between tree biomass carbon and A. Shannon-Wiener diversity index; B. Margaleaf's richness index

### *Richness index and biomass carbon*

Figure 5.B examines the relationship between the Margalef richness index and biomass carbon, showing a regression equation of  $y = 208.46x - 209.91$  with an  $R^2 = 0.56$ . This indicates a stronger correlation compared to the Shannon-Wiener index, with about 57% of the variation in biomass carbon explained by species richness alone. The Margalef index specifically measures species richness, reflecting the number of different species within a community. The stronger relationship here suggests that species richness is a critical component of biomass carbon storage, potentially because a greater variety of species can lead to more efficient resource utilization, higher productivity, and, consequently, more biomass accumulation. However, similar to the Shannon-Wiener index, the remaining 43% of unexplained variation highlights the complexity of the factors that govern biomass carbon storage in forests (Hossain et al. 2020).

The results of this study emphasize the critical role of tree DBH (diameter at breast height) and basal area as primary predictors of biomass carbon in forest stands. These structural parameters, alongside tree height and stand density, offer significant insights into how forest composition influences carbon storage. The strong correlations observed for DBH and basal area indicate that maintaining or enhancing these attributes through sustainable forest management practices could be pivotal in maximizing carbon sequestration in forests. Nonetheless, the unexplained variation in each relationship highlights the inherent complexity of forest ecosystems and underscores the need for a comprehensive approach that considers multiple factors—including species diversity, environmental conditions, and stand dynamics—in efforts to boost forest carbon storage and mitigate climate change.

We also found that biodiversity, whether assessed through species richness or a combination of richness and evenness, plays a substantial role in determining biomass carbon levels in forest ecosystems (Rahman et al. 2018). The stronger correlation observed with the Margalef richness index suggests that species richness may have a more direct impact on biomass carbon than overall diversity captured by the Shannon-Wiener index. These findings reinforce the importance of conserving biodiversity, not only for its intrinsic value but also for its crucial role in enhancing ecosystem services like carbon sequestration (Kongsager et al. 2013). However, the considerable unexplained variation in both models indicates that biodiversity is just one component of a larger, intricate system. Future research should integrate additional ecological and environmental factors that contribute to biomass carbon storage, aiming to develop more comprehensive models capable of better-predicting carbon dynamics in diverse forest ecosystems.

This study underscores the intricate interactions between native and exotic species within the CWS, with a few dominant species significantly influencing the forest's ecological balance. These findings highlight the urgent need for targeted conservation strategies to safeguard native biodiversity and maintain the ecological integrity of the reserve (Ali et al. 2022). Future research should

prioritize long-term monitoring of these species and assess the effectiveness of management practices in enhancing biodiversity and ecosystem resilience (Mitra et al. 2023). Similarly, while structural attributes such as DBH and basal area are reliable indicators of biomass carbon, the complexity of forest ecosystems necessitates a holistic approach that incorporates various factors, including biodiversity (Scherer et al. 2023), environmental conditions (Teets et al. 2023), and stand dynamics, to optimize carbon storage and address climate change (Hossain et al. 2020). Future studies should aim to incorporate additional ecological and environmental variables to refine models predicting carbon dynamics across diverse forest ecosystems.

### ACKNOWLEDGEMENTS

We are thankful to The Divisional Forest Officer, Chattogram South Forest Division, Bangladesh for providing the necessary support during the fieldwork, and also grateful to the Institute of Forestry and Environmental Sciences for lab and library facilities to conduct the work properly.

### REFERENCES

- Akwaji PI, Onah DO. 2023. Forest inventory for sustainable forest management in cross river state, Nigeria. *Sci Rep Life Sci* 4 (3): 38-52.
- Ali S, Khan SM, Siddiq Z, Ahmad Z, Ahmad KS, Abdullah A, Hashem A, Al-Arjani ABF, Abd\_Allah EF. 2022. Carbon sequestration potential of reserve forests present in the protected Margalla Hills National Park. *J King Saud Univ Sci* 34 (4): 101978. DOI: 10.1016/j.jksus.2022.101978.
- Alongi DM. 2011. Carbon payments for mangrove conservation: Ecosystem constraints and uncertainties of sequestration potential. *Carbon payments for mangrove conservation: Ecosystem constraints and uncertainties of sequestration potential. Environ Sci Policy* 14 (4): 462-470. DOI: 10.1016/j.envsci.2011.02.004.
- Baul TK, Chakraborty A, Nandi R, Mohiuddin M, Kilpeläinen A, Sultana T. 2021a. Effects of tree species diversity and stand structure on carbon stocks of homestead forests in Maheshkhali Island, Southern Bangladesh. *Carbon Balance Manag* 16: 11. DOI: 10.1186/s13021-021-00175-6.
- Baul TK, Peuly TA, Nandi R, Schmidt LH, Karmakar S. 2021b. Carbon stocks of homestead forests have a mitigation potential to climate change in Bangladesh. *Sci Rep* 11: 9254. DOI: 10.1038/s41598-021-88775-7.
- Ball DF. 1964. Loss-on-ignition as an estimate of organic matter and organic carbon in non-calcareous soils. *J Soil Sci* 15 (1): 84-92. DOI: 10.1111/j.1365-2389.1964.tb00247.x.
- Brown S. 1997. Estimating Biomass and Biomass Change of Tropical Forests: A Primer. Food & Agriculture Org.
- Chave J, Coomes D, Jansen S, Lewis SL, Swenson NG, Zanne AE. 2009. Towards a worldwide wood economics spectrum. *Ecol Lett* 12 (4): 351-366. DOI: 10.1111/j.1461-0248.2009.01285.X.
- Chave J, Réjou-Méchain M, Búrquez A, Chidumayo E, Colgan MS, Delitti WB, Duque A, Eid T, Fearnside PM, Goodman RC, Henry M. 2014. Improved allometric models to estimate the aboveground biomass of tropical trees. *Glob Change Biol* 20 (10): 3177-3190. DOI: 10.1111/gcb.12629.
- Chowdhury MIH, Das C. 2024a. Carbon sequestration potentials for conservation of Sheikh Russel Aviary and Eco-Park, Rangunia, Chittagong. *J Environ Sci Econ* 3 (2): 90-101. DOI: 10.56556/jescae.v3i2.940.

- Chowdhury MIH, Rakib MH. 2024b. Phytosociology and carbon sequestration potential of tropical forest landscape: A case from Northwest Chittagong, Bangladesh. *Asian J For* 8: 126-135. DOI: 10.13057/asianjfor/r080203.
- Chowdhury MSH. 2014. Forest conservation in protected areas of Bangladesh. *World Forests*. Springer, Cham. DOI: 10.1007/978-3-319-08147-2.
- Chowdhury MA, Islam KN, Hafiz N, Islam K. 2019. Diversity of trees in a community managed forest: The case of Komolchori VCF, Khagrachari, Bangladesh. *Geol Ecol Landsc* 3: 95-103. DOI: 10.1080/24749508.2018.1508980.
- Dallmeier F, Kabel M, Rice R, Taylor CM. 1992. Long-term monitoring of biological diversity in tropical forest areas: Methods for establishment and inventory of permanent plots. UNESCO, Paris.
- Das N. 2024. Tree species diversity, composition and structure in the tropical moist deciduous forest of Kadigarh National Park, Mymensingh, Bangladesh. *Asian J For* 8: 41-49. DOI: 10.13057/asianjfor/r080104.
- Festus AG, Zakaria R, Rusly R, Bernard O. 2024. Ferns richness along environmental gradients in a tropical forest ecosystem. *Sci Rep Life Sci* 5 (2): 92-106. DOI: 10.5281/zenodo.11303599.
- Hasan M, Jorin T, Sayok AK. 2023. Biodiversity information systems in geospatial applications for protected area management. *Am J Geogr Inf Syst* 12 (1): 1-27. DOI: 10.5923/j.ajgis.20231201.01.
- Hop NV, Quy NV, Lam NV, Trong PT, Tinh PC. 2023. Woody plant diversity and aboveground carbon stock of Dipterocarpus chartaceus dominant forests in Binh Chau-PhuocBuu Nature Reserve, South Vietnam. *Asian J For* 7 (2): 115-125. DOI: 10.13057/asianjfor/r070206.
- Hossain MK, Alim A, Hossen S, Hossain A, Rahman A. 2020. Diversity and conservation status of tree species in Hazarikhil Wildlife Sanctuary (HWS) of Chittagong, Bangladesh. *Geol Ecol Landsc* 4 (4): 298-305. DOI: 10.1080/24749508.2019.1694131.
- Hossain MA, Hossain MK, Salam M, Rahman S. 2013. Composition and diversity of tree species in Dudhpukuria-Dhopachori Wildlife Sanctuary of Chittagong (South) Forest Division, Bangladesh. *Res J Pharm Biol Chem Sci* 4 (2): 1447-1457.
- Islam K, Jashimuddin M, Nath B, Nath TK. 2018. Land use classification and change detection by using multi-temporal remotely sensed imagery: The case of Chhunati wildlife sanctuary, Bangladesh. *Egypt J Remote Sens Space Sci* 21 (1): 37-47. DOI: 10.1016/j.EJRS.2016.12.005.
- Islam I, Cui S, Hoque MZ, Abdullah HM, Tonny KF, Ahmed M, Ferdush J, Xu L, Ding S. 2022. Dynamics of tree outside forest land cover development and ecosystem carbon storage change in Eastern Coastal Zone, Bangladesh. *Land* 11 (1): 76. DOI: 10.3390/land11010076.
- Islam KN, Rahman MM, Jashimuddin M, Islam K, Zhang Y. 2020. Impact of co-management on tree diversity and carbon sequestration in protected areas: Experiences from Bangladesh. *Trees For People* 2: 100033. DOI: 10.1016/j.tfp.2020.100033.
- Jaman M, Hossain M, Islam M, Helal M, Jamil M. 2016. Quantification of carbon stock and tree diversity of homegardens in Rangpur District, Bangladesh. *Intl J Agric For* 6 (5): 169-180.
- Kessler M, Abrahamczyk S, Bos M, Buchori D, Putra DD, Gradstein SR, Höhn P, Kluge J, Orend F, Pitopang R, Saleh S, Schulze CH, Sporn SG, Steffan-Dewenter I, Tjitrosoedirdjo SS, Tschardt T. 2009. Alpha and beta diversity of plants and animals along a tropical land-use gradient. *Ecol Appl* 19 (8): 2142-2156. DOI: 10.1890/08-1074.1.
- Kohli RK, Singh HP, Rani D. 1996. Status of floor vegetation under some monoculture and mixculture plantations in north India. *J For Res* 1 (4): 205-209. DOI: 10.1007/BF02348326.
- Kongsager R, Napier J, Mertz O. 2013. The carbon sequestration potential of tree crop plantations. *Mitigation Adapt Strategies Global Change* 18 (8): 1197-1213. DOI: 10.1007/s11027-012-9417-z.
- Kufata FH, Lemessa D, Bekele Jiru D. 2024. The perceptions of the local peoples on the trends of the illegal practices inside and outside Dhati Wolel national park in west Ethiopia. *Sci Rep Life Sci* 5 (2): 1-17. DOI: 10.5281/zenodo.11215159.
- Macdicken K. 2015. A Guide to Monitoring Carbon Storage in Forestry and Agroforestry Projects.
- Mamun A AI, Hossain MK, Hossain Md A. 2022. Ecological restoration through natural regeneration in Chhunati Wildlife Sanctuary – a protected area of South-East Bangladesh. *Ecofeminism Clim Change* 3 (1): 41-55. DOI: 10.1108/EFCC-06-2020-0019.
- Mehta B, Nagar B, Patel B, Chaklashiya P, Shah M, Verma P, Shah MB. 2021. A review on a Lesser Known Indian Mangrove: *Avicennia officinalis* L. (Family: Acanthaceae). *Intl J Green Pharm* 15 (1): 1-10.
- Miah MD, Hasnat GNT, Nath B, Saem, MGU, Rahman MM. 2023. Spatial and temporal changes in the urban green spaces and land surface temperature in the Chittagong City Corporation of Bangladesh Between 2000 and 2020. *Forestist* 73 (2): 171-182. DOI: 10.5152/forestist.2022.22013.
- Miah R, Hasan M, Parisha JT, Sayok AK. 2023. A framework on biodiversity conservation related policy analysis. *Am J Environ Eng* 13 (1): 1-12. DOI: 10.5923/j.ajee.20231301.01.
- Michael P. 1990. Ecological methods for field and laboratory investigation. Tata Mc Graw Hill, New Delhi, India.
- Mitra S, Naskar N, Lahiri S, Chaudhuri P. 2023. A study on phytochemical profiling of *Avicennia marina* mangrove leaves collected from Indian Sundarbans. *Sustain Chem Environ* 4: 100041. DOI: 10.1016/j.scenv.2023.100041.
- Mukul SA. 2019. Biodiversity Conservation and Sustainable Development in Bangladesh: An overview of the present status, management problems and future prospects. Department of Forestry and Environmental Science, Sylhet, Bangladesh. DOI: 10.13140/2.1.3573.4726.
- Pearson T, Walker S, Brown S. 2005. Sourcebook for Land use, Land-use change and forestry projects.
- Pozo P, Säumel I. 2018. How to bloom the green desert: Eucalyptus plantations and native forests in Uruguay beyond black and white perspectives. *Forests* 9 (10): 614. DOI: 10.3390/f9100614.
- Rahman Md. H, Roy B, Chowdhury GM, Hasan A, Saimun Md. SR. 2022. Medicinal plant sources and traditional healthcare practices of forest-dependent communities in and around Chhunati Wildlife Sanctuary in southeastern Bangladesh. *Environ Sustain* 5 (2): 207-241. DOI: 10.1007/S42398-022-00230-Z.
- Rahman MM, Mahmud MAAI, Ahmed FU. 2017. Restoration of degraded forest ecosystem through non-forestry livelihood supports: Experience from the Chhunati Wildlife Sanctuary in Bangladesh. *For Sci Technol* 13 (3): 109-115. DOI: 10.1080/21580103.2017.1349003.
- Rahman M, Parvin W, Sultana N, Tarek S. 2018. Ex-situ conservation of threatened forest tree species for sustainable use of forest genetic resources in Bangladesh. *J Biodiv Conserv Bioresour Manag* 4 (2): 89-98. DOI: 10.3329/jbcbm.v4i2.39855.
- Rahman S, Obaydullah AKM. 2020. Causes for degradation to Chhunati Wildlife Sanctuary and its restoration strategy. *Intl J Res Innov Soc Sci (IJRISS)* 4 (5): 90-97.
- Rakib MH, Chowdhury MIH, Das C, Tonima H. 2024. Wildlife Ecological Spectrum: unveiling alpha ( $\alpha$ ), beta ( $\beta$ ), and gamma ( $\gamma$ ) diversity of the Kaptai National Park, Bangladesh. DOI: 10.21203/rs.3.rs-4668666/v1.
- Roy M, Biswas B, Ghosh S. 2015. Trend analysis of climate change in Chittagong Station in Bangladesh. *Intl Lett Nat Sci* 47 (1): 42-53. DOI: 10.18052/www.scipress.com/ILNS.47.42.
- Sattar MA, Bhattachajee DK, Kabir MF. 1999. Physical and mechanical properties and uses of timbers of Bangladesh. Bangladesh Forest Research Institute, Chittagong.
- Scherer L, Rosa F, Sun Z, Michelsen O, De Laurentis V, Marques A, Pfister S, Verones F, Kuipers KJJ. 2023. Biodiversity impact assessment considering land use intensities and fragmentation. *Environ Sci Technol* 57 (48): 19612-19623. DOI: 10.1021/acs.est.3c04191.
- Sheikh MA, Kumar M, Bussman RW, Todaria N. 2011. Forest carbon stocks and fluxes in physiographic zones of India. *Carbon Balance Manag* 6 (1): 15. DOI: 10.1186/1750-0680-6-15.
- Shukla R, Chandel PS. 2016. A Textbook of Plant Ecology including Ethnobotany and Soil Science.
- Simon LN, Mbua RL, Roger E. 2018. Estimation of carbon stock in Bimbia Bonadikombo coastal community forest, south west region, Cameroon: An implication for climate change mitigation. *Intl J Sci Res Manag* 6 (10): FE-2018. DOI: 10.18535/ijrsm/v6i10.fe02.
- Teets A, Bailey AS, Hufkens K, Ollinger S, Schädle C, Seyedsnasrollah B, Richardson AD. 2023. Early spring onset increases carbon uptake more than late fall senescence: Modeling future phenological change in a US northern deciduous forest. *Oecologia* 201: 241-257. DOI: 10.1007/s00442-022-05296-4.
- Uddin M, Chowdhury FI, Hossain MK. 2020a. Assessment of tree species diversity, composition and structure of Medha Kachhapi National Park, Cox's Bazar, Bangladesh. *Asian J For* 4 (1): 15-21. DOI: 10.13057/asianjfor/r040104.

- Uddin MN, Hossain MM, Karim MS, Siriwong W, Boonyanuphap J. 2020b. The phytosociological attributes of village common forests in Chittagong hill tracts, Bangladesh. *Songklanakarin J Sci Technol* 42 (4): 819-829. DOI: 10.14456/sjst-psu.2020.105.
- Zhou Z, Ollinger SV, Lepine LC. 2018. Landscape variation in canopy nitrogen and carbon assimilation in a temperate mixed forest. *Oecologia* 188: 595-606. DOI: 10.1007/s00442-018-4223-2.
- Zukswert J, Vadeboncoeur MA, Yanai RD. 2023. Responses of stomatal density and carbon isotope composition of sugar maple and yellow birch foliage to N, P, and CaSiO<sub>3</sub> fertilization. *Tree Physiol* 44 (1): tpad142. DOI: 10.1093/treephys/tpad142.